



THÈSE

En vue de l'obtention du
DOCTORAT DE L'UNIVERSITÉ DE TOULOUSE
Délivré par l'Université Toulouse 3 – Paul Sabatier

Présentée et soutenue par
DENIS ROCHETTE

Le 11 juillet 2023

ASYMMETRIC CLONING IN QUANTUM INFORMATION THEORY

École doctorale : **Mathématiques, Informatique, Télécommunications de Toulouse**

Spécialité : **Mathématiques**

Unité de recherche : **Institut de Mathématiques de Toulouse** (UMR 5219)

Thèse dirigée par

M. Clément Pellegrini
M. Ion Nechita

Jury

M. Guillaume Aubrun	Rapporteur
M. Omar Fawzi	Rapporteur
M. Michael Wolf	Rapporteur
M. Antonio Acín	Examineur
M. Fabrice Gamboa	Examineur
Mme. Maria Jivulescu	Examinatrice

Acknowledgements

My first words of thanks go to my supervisors **Clément Pellegrini** and **Ion Nechita** for giving me the opportunity to do a PhD under their supervision and with their advice. Throughout these three years, their kindness and support have been indispensable to the success of this thesis. Thank you for giving me the opportunity to attend many conferences, workshops and summer schools. These invaluable experiences have greatly enriched my research and academic network, for which I am deeply grateful.

I would like to thank **Guillaume Aubrun**, **Omar Fawzi** and **Michael Wolf** who accepted to review this manuscript and did so with great care. I would also like to thank **Antonio Acín**, **Fabrice Gamboa** and **Maria Jivulescu** for accepting to be members of my thesis defence jury.

I would like to express my deep gratitude to the permanent members of the Institut de Mathématiques de Toulouse and the Probability team, with whom I have had the opportunity to discuss mathematics and other subjects. My special thanks go to **Tristan Benoist** and **Reda Chhaibi**, whose advice and encouragement have been crucial to my progress.

I would like to thank all the students who are part of our Quantum team in Toulouse or who have joined us in the past for a thesis, an internship or just a visit. Thanks to **Arnaud**, **Pierre**, **Khurshed**, **Jan-Luka**, **Linda**, **Satvik**, **Qing-Hua**, **Sang-Jun**, and all the new students who have joined us this year. Thanks to **Anna**, our first postdoc! And of course a special thanks to **Faedi**, with whom I spent many hours over a blackboard or a drink.

I would like to thank all the PhD students and postdocs at the Institut de Mathématiques de Toulouse whom I have had the pleasure to meet, and with whom I've spent some great times, in the lab, over a picnic in the park, or a drink in the Biergarten. Thank you for organising all the student activities in the lab, especially the student seminar. And of course, thank you for helping me to improve my Tarot skills. There are so many of you, so to mention just a few, thank you to **Nicolas**, **Sophia**, **Paola**, **Lucas**, **Perla**, **Alberto**, **Fanny**, **Michèle**, **Clément**, **Étienne**, **Viviana**, **Anthony**, **Virgile**, **Joachim**, **Corentin**, **Mahmoud**, **Alain**, **Fu-Hsuan**, **Javier**, **Axel**, Louis —

I would also like to thank all the PhD students and postdocs at the Theoretical Physics Laboratory for welcoming me among the physicists over the last three years. Thank you for all the pool, restaurant and cinema evenings... Thanks for all the discussions in the best coffee room and the seminars. Special thanks to **Bhupen**,

Sasank, Jeanne, Naïmo, Nathan, Noam, Quentin, Théo, Thomas ².

I would like to express my sincere gratitude to **Malika** for the invaluable support and dedication provided throughout the three years of my thesis. Your assistance, enthusiasm, and kindness have been truly exceptional and irreplaceable.

I would like to thank the Amsterdam QuSoft quantum team for inviting me for a week. In particular, I would like to thank, **Dmitry** and **Māris** for your welcome and for our joint work.

I would like to thank Quantum Munich at TUM for their kind invitation. I would especially like to thank **Cambyse**.

Finally, I'd like to thank all the people I've met during the three years of my thesis. Thank you to **Cécilia**, who was part of our Quantum team in Toulouse for a while. Thanks to **Andreas, Mehdi, Leevi, Teiko, Stephan, Yvan, Adrian, Matthias** and many others. I would especially like to thank **Anne** for giving me the opportunity to continue doing quantum science during a postdoc.

My next thanks go to my friends **Pierre** and **Herménégilde**. Congratulations to you **Julien** and **Héloïse** on your wedding, congratulations to you **Joseph** and **Shiva-chan**, and welcome to you **Léo**. Thank you **Julien, Jérémy, Tristan, Rémi, Alexandre, Steven, Johanna, Youri, Viven, Mickaël, Maëlys** and all the others for the time we spent together.

I would like to thank my family for their support and help during the three years of my thesis in Toulouse. I know the long distance has not always been easy.

Finally, I'd like to thank you, **Dr. Claire Lacouture**. Thank you for your presence, your love, and your support that made the writing of this manuscript so much easier.

2. It wasn't a footnote, just the two Thomas!

Contents

1	Schur-Weyl duality	7
1.1	Diagrammatic algebras	9
1.2	Tensor representation	20
1.3	Schur-Weyl dualities	23
2	Quantum Information Theory	29
2.1	Postulates of quantum mechanics	31
2.2	Quantum entanglement	34
2.3	Quantum channels	38
2.4	Quantum fidelity	39
2.5	Graphical calculus	40
3	Quantum Cloning	43
3.1	Quantum cloning problem	45
3.2	Upper bound	54
3.3	Universal $1 \longrightarrow 2$ quantum cloning problem	57
3.4	Universal $1 \longrightarrow N$ quantum cloning problem	65
4	Extendibility	85
4.1	Extendibility of quantum states on a graph	87
4.2	Quantum cloning: star-graph extendibility of isotropic states	88
4.3	K_N -Extendibility of isotropic states	89
A	Elements of representation theory	105
A.1	Finite groups	107
A.2	Matrix groups	140

Schur-Weyl duality

Chapter contents

1.1	Diagrammatic algebras	9
1.1.1	Symmetric group algebra \mathbb{S}_k	10
1.1.2	Partition algebra $\mathbb{P}_k(\delta)$	12
1.1.3	Others diagrammatic algebras	15
1.2	Tensor representation	20
1.3	Schur-Weyl dualities	23
1.3.1	Commutant	23
1.3.2	Schur-Weyl duality for \mathfrak{S}_n	23
1.3.3	Other Schur-Weyl dualities	26

The primary goal of this chapter consists in establishing the foundational result of Issai Schur and Hermann Weyl, known as the Schur-Weyl duality, which relates the symmetric group \mathfrak{S}_n and the complex general linear group GL_d . Moreover, this chapter explores other adaptations of this theorem for distinct groups and algebras.

Theorem (Schur-Weyl duality [Sch27; Wey46]). *The space of n -fold tensors over \mathbb{C}^d decomposes under the action of the direct product group $\mathrm{GL}_d \times \mathfrak{S}_n$ as follows:*

$$(\mathbb{C}^d)^{\otimes n} \simeq \bigoplus_{\substack{\lambda \vdash n \\ \lambda_1 \leq d}} V_\lambda^d \otimes V_\lambda.$$

Where V_λ^d is an irreducible representation space of GL_d and V_λ is an irreducible representation space of \mathfrak{S}_n .

For a comprehensive exploration of the representation theory concerning the symmetric group \mathfrak{S}_n and the complex general linear group GL_d , refer to [Appendix A](#). However, the current section aims to provide a self-contained exposition.

Subsequently, two notations for permutations and partitions of the set $\{1, \dots, n\}$ are employed. The conventional cyclic notation $(1\ 2\ 3)(4\ 5)$ denotes the permutation:

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix},$$

and the notation $1\ 2\ 3 \mid 4\ 5$ represents the partition:

$$\{\{1, 2, 3\}, \{4, 5\}\}.$$

1.1 Diagrammatic algebras


The term diagram algebras has no specific definition by axiomatic properties or other rigorous means. In the present thesis, a diagram algebra refers to a finite unital associative algebra over the complex field, where the basis consists of homotopy classes of diagrams. The multiplication operation within this algebraic structure finds its definition through the process of concatenation. For a survey on diagram algebras, see [Koe08; HJ20].

In the context of finite dimensional algebras over the algebraically closed field \mathbb{C} , the notion of **semisimplicity** is employed interchangeably with that of the direct sum of full matrix algebras, closed under matrix multiplication.

1.1.1 Symmetric group algebra \mathbb{S}_k


Define \mathfrak{S}_k as the **symmetric group**, which is the group of order $k!$ containing all the **permutations** of the set $\{1, \dots, k\}$. Given a permutation σ belonging to the symmetric group \mathfrak{S}_k , it is possible to represent this permutation as a **diagram** via a graph consisting of $2k$ vertices. These vertices are divided equally between two columns.

Interpretation of the diagram proceeds from right to left. A connection between the i -th vertex in the right column and the j -th vertex in the left column is established if and only if the relation $\sigma(i) = j$ holds. For example,




represents the permutation $(1\ 2\ 3)$ of the symmetric group \mathfrak{S}_3 .

The composition, denoted by $\sigma \circ \tau$, of two permutations σ and τ of the symmetric group \mathfrak{S}_k , is obtained by positioning the diagram of τ immediately to the right of the diagram of σ , and subsequently associating the leftmost column of τ 's diagram with the rightmost column of σ 's diagram. For example, consider the following two permutations $\sigma := (1\ 2\ 4)$ and $\tau := (1\ 2)(3)$, which are elements of the symmetric group \mathfrak{S}_3 ,




the composition $\sigma \circ \tau = (1\ 3)(2)$ becomes



Numerous generating sets for the symmetric group \mathfrak{S}_k exist, with varying cardinalities. A particularly notable generating set is the collection of adjacent **transpositions**. This set is characterized by containing the $k - 1$ permutations of the form

$(i \ i+1)$, for all $1 \leq i < k$:




Let σ be an element of the symmetric group $\mathfrak{S}k$. The **cycle type** associated with σ , denoted by λ , is defined as the l -tuple containing the lengths of the l disjoint **cycles** composing σ , arranged in non-increasing order. As a consequence, the cycle type λ corresponds to a **partition** of the integer k into l parts, denoted by $\lambda \vdash k$. This partition λ obeys the following conditions:

$$\lambda_1 \geq \cdots \geq \lambda_l \quad \text{and} \quad \sum_{i=1}^l \lambda_i = k.$$

Given a partition $\lambda \vdash k$, let λ' denote the **conjugate** partition associated with λ , defined by: λ'_i is the number of parts in λ that are greater than or equal to i . A partition $\lambda \vdash k$ may be represented as a **Young diagram**, which is a collection of k empty boxes arranged in left-justified rows such that the i -th row contains λ_i boxes. The conjugate partition λ' , is the partition corresponding to transposing the Young diagram representing λ . For example, consider the permutation $\sigma \in \mathfrak{S}_7$, expressed as a product of disjoint cycles, arranged in non-increasing order, by


$$\sigma := (1 \ 2 \ 3 \ 4)(5 \ 6 \ 7).$$

The cycle type of σ is the partition $\lambda \vdash 7$ given by $\lambda := (4, 3)$, where the i -th entry of λ denotes the length of the i -th cycle in the disjoint cycle decomposition of σ . Moreover, the partition λ can be represented using a Young diagram, which consists of 2 rows with 4 and 3 boxes, respectively



The conjugate partition $\lambda' := (2, 2, 2, 1)$ is represented using a Young diagram, which

consists of 4 rows with 2, 2, 2, and 1 boxes, respectively:



Note that λ'_1 is the length of the first column of λ . Within the context of the symmetric group \mathfrak{S}_k , the concept of cycle type plays a crucial role in characterizing **conjugacy classes**. Specifically, two permutations in \mathfrak{S}_k are said to be **conjugate** if and only if their respective cycle types are identical.

Remark. It is essential to note that the symmetric group \mathfrak{S}_k , defined as permutations of the set $\{1, \dots, k\}$ or as the above diagrams conveys the same underlying mathematical structure. These two forms are equivalent and can be employed interchangeably.

The **group algebra** of the symmetric group \mathfrak{S}_k , denoted by \mathbb{S}_k , is the complex vector space spanned by the permutations of \mathfrak{S}_k , i.e.


$$\mathbb{S}_k := \text{Span}_{\mathbb{C}} \{ \sigma \in \mathfrak{S}_k \}.$$

The multiplication in the group algebra \mathbb{S}_k , is defined on the elements of the symmetric group \mathfrak{S}_k by its group law, and is denoted $\sigma \cdot \tau$, for some σ and τ in \mathfrak{S}_k .


Remark. The symmetric group algebra \mathbb{S}_k is a finite group algebra. As a consequence, the symmetric group algebra \mathbb{S}_k is always semisimple [Ser+77; FH13].

1.1.2 Partition algebra $\mathbb{P}_k(\delta)$

The **partition monoid**, denoted \mathbb{P}_k , is a diagrammatic monoid generated by the $3 \times (k - 1)$ diagrams




for all $1 \leq i < k$, as well as the k disconnected diagrams




for all $1 \leq i \leq k$. These $4k - 3$ diagrams distributed in 4 distinct collections do not constitute a minimal generating set, as it is possible to choose a single representative diagram from each collection and subsequently use the transpositions to generate the remaining diagrams in the collection.


An element of \mathbb{P}_k is a **partition** of the set $\{1, \dots, 2k\}$, corresponding to the connected components of the associated diagram, where then enumeration of vertices located in the right column ranges from 1 to k , while the enumeration of vertices situated in the left column ranges from $k+1$ to $2k$. As a monoid, \mathbb{P}_k has an identity $1_{\mathbb{P}_k}$ given by the partition $1(k+1) \mid \dots \mid k(2k)$:



Consider two partitions p and q of \mathbb{P}_k , the composition $p \circ q$ is obtained by positioning the diagram of q immediately to the right of the diagram of p , associating the leftmost column of q 's diagram with the rightmost column of p 's diagram, and finally removing any **loops**, which are the components of the resulting diagram not connected to either the left or the right column. For example, given the two partitions $p := 13 \mid 26 \mid 45$ and $q := 12 \mid 35 \mid 46$ of \mathbb{P}_3 ,



the composition $p \circ q = 12 \mid 36 \mid 45$ becomes



where the gray loop is removed.

Remark. Considering the generators of the partition monoid, the inclusion of diagrams $\mathfrak{S}_k \subseteq \mathbb{P}_k$ holds for every $k \in \mathbb{N}$. However, it is important to note that the partition monoid \mathbb{P}_k does not constitute a group, e.g. the partition $1\,2|3\,6|4\,5$ of \mathbb{P}_3 has no inverse with respect to the composition in \mathbb{P}_3 .

The order of the partition monoid \mathbb{P}_k is the number of partition of the set $\{1, \dots, 2k\}$, denoted as the even **Bell number** B_{2k} . In general, the Bell number B_k is given by a recursive formula, with initial condition $B_0 := 1$, and

$$B_{k+1} = \sum_{i=0}^k \binom{k}{i} B_i.$$


Starting at $k = 0$, the first values of the Bell numbers are [OEIS, sequence A000110]:

$$1, 1, 2, 5, 15, 52, 203, 877, 4140, \dots$$


The **partition algebra**, denoted by $\mathbb{P}_k(\delta)$ is defined for some $\delta \in \mathbb{C}$, as the complex vector space spanned by the diagrams of \mathbb{P}_k , i.e.

$$\mathbb{P}_k(\delta) := \text{Span}_{\mathbb{C}} \{p \in \mathbb{P}_k\}.$$

The multiplication in $\mathbb{P}_k(\delta)$, given two elements p and q of \mathbb{P}_k , is denoted by $p \cdot q$ and is defined by $p \cdot q := \delta^l(p \circ q)$, where l is the number of loops removed during the composition in \mathbb{P}_k . For example, let $p := 1\,4|2\,8|3\,5|6\,7|9\,10$ and $q := 1\,2|3\,5|4\,7|6\,9|8\,10$ be two partitions of \mathbb{P}_5 ,



the composition $p \circ q = 1\,2|3\,5|4\,8|6\,7|9\,10$ becomes, in the partition algebra $\mathbb{P}_5(\delta)$,



where the number of gray loops removed is 2.

Remark. The partition algebra $\mathbb{P}_k(\delta)$ is not semisimple for all $\delta \in \mathbb{C}$. Specifically, it is semisimple if and only if δ belongs to the set $\mathbb{C} \setminus \{0, \dots, 2k-2\}$ [MS94; HR05].

1.1.3 Others diagrammatic algebras


In Section 1.1.1, the focus was on the symmetric group \mathfrak{S}_k , generated by $k-1$ permutation diagrams. Then, in Section 1.1.2, attention was turned to the partition monoid \mathbb{P}_k , generated by $4k-3$ diagrams distributed in 4 distinct collections.

The purpose of the present Section is to highlight the relationships between the 4 collections of diagrams generating the partition monoid \mathbb{P}_k , and the algebraic structures that emerge. Specifically, the choice of specific collections may yield distinct monoids.

Remark. A monoid is a unitary semigroup. To obtain the identity diagram $1_{\mathbb{P}_k}$ of the partition monoid \mathbb{P}_k , corresponding to the partition $1(k+1) \mid \dots \mid k(2k)$, the collection of transpositions is required, as none of the 3 other collections is composed of invertible elements.

Uniform block permutation algebra $\mathbb{U}_k(\delta)$


The **uniform block permutation monoid**, denoted by \mathbb{U}_k , is the submonoid of \mathbb{P}_k generated by the 2 collections of diagrams




for all $1 \leq i < k$.

The elements of the uniform block permutation monoid \mathbb{U}_k are precisely those elements from \mathbb{P}_k that satisfy the following condition: the number of vertices located on the left column equals the number of vertices located on the right column for each connected component of the diagram. This condition is a consequence of the fact that the 2 collections generating \mathbb{U}_k satisfy it and that it is preserved under multiplication.

As a consequence, each connected component contains vertices situated in both the left and right columns. For example, the partition $p := 1\,2\,4\,5 \mid 3\,6$ is in \mathbb{U}_3 ,



while the partition $q := 1\,2 \mid 3\,6 \mid 4\,5$ is not in \mathbb{U}_3 ,



The order of the uniform block permutation monoid \mathbb{U}_k is given by a recursive formula [SPS01], with initial condition $|\mathbb{U}_0| := 1$, and

$$|\mathbb{U}_{k+1}| = \sum_{i=1}^k \binom{k}{i} \binom{k+1}{i} |\mathbb{U}_i|.$$

Starting at $k = 0$, the first values of $|\mathbb{U}_k|$ are [OEIS, sequence A023998]:

$$1, 1, 3, 16, 131, 1496, \dots$$

The **uniform block permutation algebra**, denoted by $\mathbb{U}_k(\delta)$ is defined for some $\delta \in \mathbb{C}$, as the subalgebra of the partition algebra $\mathbb{P}_k(\delta)$, spanned by the element of the uniform block permutation monoid \mathbb{U}_k , i.e


$$\mathbb{U}_k(\delta) := \text{Span}_{\mathbb{C}} \{p \in \mathbb{U}_k\}.$$

The multiplication in $\mathbb{U}_k(\delta)$ does not yield any loops. Consequently, all the uniform block permutation algebras are isomorphic, for all $\delta \in \mathbb{C}$.

Remark. The uniform block permutation monoid \mathbb{U}_k is a finite **inverse monoid**, i.e. for all $x \in \mathbb{U}_k$ there exists a unique $x^* \in \mathbb{U}_k$ satisfying $x \circ x^* \circ x = x$ and $x^* \circ x \circ x^* = x^*$. As a consequence, all the uniform block permutations algebras $\mathbb{U}_k(\delta)$ are semisimple for all $\delta \in \mathbb{C}$ [Ore+21; Ste+16].


Brauer algebra $\mathbb{B}_k(\delta)$

The **Brauer monoid**, denoted by \mathbb{B}_k , is the submonoid of \mathbb{P}_k generated by the 2 collections of diagrams



for all $1 \leq i < k$.

The elements of the Brauer monoid \mathbb{B}_k are precisely all the pairings on the set $\{1, \dots, 2k\}$. Specifically, each vertex of a diagram in \mathbb{B}_k has precisely a degree of 1. Given a diagram of \mathbb{B}_k , the **vertical edges** are the edges that connect vertices within the same column, whereas the **horizontal edges** are the edges that connect vertices of both the left and right columns. For example, the partition $p := 1\,3 \mid 2\,6 \mid 4\,5$ is in \mathbb{B}_3 ,



The order of the Brauer monoid \mathbb{B}_k is given by the **odd factorial**,

$$|\mathbb{B}_k| = (2k - 1)!!.$$

Starting at $k = 0$, the first values of $|\mathbb{B}_k|$ are [OEIS, sequence A001147]:

$$1, 1, 3, 15, 105, 945, 10395, \dots$$

The **Brauer algebra**, denoted by $\mathbb{B}_k(\delta)$ is defined for some $\delta \in \mathbb{C}$, as the subalgebra of the partition algebra $\mathbb{P}_k(\delta)$, spanned by the element of the Brauer monoid \mathbb{B}_k , i.e

$$\mathbb{B}_k(\delta) := \text{Span}_{\mathbb{C}} \{p \in \mathbb{B}_k\}.$$


Remark. The Brauer algebra $\mathbb{B}_k(\delta)$ is not semisimple for all $\delta \in \mathbb{C}$. Specifically it is semisimple if and only if one of the following conditions holds [Wen88; DWH99;

[Rui05; RS06; AST17]:


- $\delta = 0$ and $k \in \{1, 3, 5\}$;
- $\delta \in \mathbb{Z} \setminus \{0\}$ and $k \leq |\delta| + 1$;
- $\delta \notin \mathbb{Z}$.

Walled Brauer algebra $\mathbb{B}_{k,l}(\delta)$

The **walled Brauer monoid**, denoted by $\mathbb{B}_{k,l}$, is the submonoid of \mathbb{B}_{k+l} generated by the 2 collections of diagrams




for all $1 \leq i < k < j < k + l$, as well as the diagram




The **wall** of the walled Brauer monoid $\mathbb{B}_{k,l}$ denotes the vertical separation between the uppermost $2k$ vertices and the lowermost $2l$ vertices. The diagram elements of $\mathbb{B}_{k,l}$ are precisely those elements from \mathbb{B}_{k+l} that satisfy the following condition: every vertical edge must cross the wall, while no horizontal edge shall cross the wall. The condition arises from the fact that the $k + l - 1$ diagrams generating $\mathbb{B}_{k,l}$ satisfy the

condition, in addition to preserving this property under multiplication. For example, the partition $p := 1\,4 \mid 2\,7 \mid 3\,9 \mid 5\,10 \mid 6\,12 \mid 8\,11$ is in $\mathbb{B}_{3,3}$,




while the partition $q := 1\,3 \mid 2\,7 \mid 4\,9 \mid 5\,10 \mid 6\,12 \mid 8\,11$ is in \mathbb{B}_6 but not in $\mathbb{B}_{3,3}$,



Remark. Given the generators of the walled Brauer monoid $\mathbb{B}_{k,l}$, the inclusion of diagrams $\mathfrak{S}_k \times \mathfrak{S}_l \subseteq \mathbb{B}_{k,l}$ holds for every $k, l \in \mathbb{N}$. The diagrams of $\mathfrak{S}_k \times \mathfrak{S}_l$ consists of those from $\mathbb{B}_{k,l}$ with no edges crossing the wall.

The operation denoted as **partial transposition** corresponds to the process that exchanges vertex i and vertex $k + l + i$, both situated on the same row. However, it should be noted that the diagram obtained after performing such an operation may not necessarily belong to $\mathbb{B}_{k,l}$ anymore. For example, the transposition of the 1-st row of the partition $1\,4 \mid 2\,7 \mid 3\,9 \mid 5\,10 \mid 6\,12 \mid 8\,11$ in $\mathbb{B}_{3,3}$ becomes,



which correspond to the partition $1\,2\,|\,3\,9\,|\,4\,7\,|\,5\,10\,|\,6\,12\,|\,8\,11$ in \mathbb{B}_6 but not in $\mathbb{B}_{3,3}$.

The partial transposition involving the l lowest vertices constitutes a one-to-one mapping from the walled Brauer monoid $\mathbb{B}_{k,l}$ to the symmetric group \mathfrak{S}_{k+l} . As a consequence,

$$|\mathbb{B}_{k,l}| = (k+l)!.$$

The **walled Brauer algebra**, denoted by $\mathbb{B}_{k,l}(\delta)$ is defined for some $\delta \in \mathbb{C}$, as the subalgebra of the Brauer algebra $\mathbb{B}_{k+l}(\delta)$, spanned by the element of the walled Brauer monoid $\mathbb{B}_{k,l}$, i.e

$$\mathbb{B}_{k,l}(\delta) := \text{Span}_{\mathbb{C}} \{p \in \mathbb{B}_{k,l}\}.$$

Remark. The walled Brauer algebra $\mathbb{B}_{k,l}(\delta)$ is not semisimple for all $\delta \in \mathbb{C}$. Specifically it is semisimple if and only if one of the following conditions hold [Cox+08; Bul20]:


- $k = 0$ or $l = 0$;
- $\delta = 0$ and $(k, l) \in \{(1, 2), (2, 1), (1, 3), (3, 1)\}$;
- $\delta \in \mathbb{Z} \setminus \{0\}$ and $k + l \leq |\delta| + 1$;
- $\delta \notin \mathbb{Z}$.

1.2 Tensor representation

The diagrammatic algebras described in Section 1.1 act on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$ by considering the mapping ψ , called **tensor representation**, from the diagrammatic monoids on $2n$ vertices to \mathcal{M}_{d^n} , and defined for each diagram p by,

$$(\psi(p))_{i_{n+1}, \dots, i_{2n}}^{i_1, \dots, i_n} := \begin{cases} 1 & \text{if } i_k = i_l, \text{ for all vertices } k \text{ and } l \text{ connected in } p \\ 0 & \text{otherwise,} \end{cases}$$

and linearly extended to the entire diagrammatic algebra. For example, let the partition $p := 1\,3\,|\,2\,6\,|\,4\,5$ in the Brauer algebra $\mathbb{B}_3(d)$,



with some vector $x_1, x_2, x_3 \in \mathbb{C}^d$ and $y_1, y_2, y_3 \in \mathbb{C}^d$, then

$$\begin{aligned} \langle y_1 \otimes y_2 \otimes y_3 | \psi(p) | x_1 \otimes x_2 \otimes x_3 \rangle &= \begin{array}{c} y_1 \bullet \\ y_2 \bullet \\ y_3 \bullet \end{array} \begin{array}{c} \curvearrowright \\ \diagup \\ \diagdown \end{array} \begin{array}{c} \bullet x_1 \\ \bullet x_2 \\ \bullet x_3 \end{array} \\ &= \langle x_1, x_3 \rangle \cdot \langle x_2, y_3 \rangle \cdot \langle y_1, y_2 \rangle. \end{aligned}$$

Remark. In the case where the diagrammatic algebras depends on a complex parameter $\delta \in \mathbb{C}$, the tensor representation acting on the tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, requires that $\delta = d$.

The tensor representations on $\mathbb{C}^2 \otimes \mathbb{C}^2$, of all the 3 diagrams spanning the Brauer algebra $\mathbb{B}_2(2)$ are

$$\psi \left(\begin{array}{cc} \bullet & \bullet \\ \curvearrowright & \curvearrowleft \\ \bullet & \bullet \end{array} \right) = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \quad \psi \left(\begin{array}{cc} \bullet & \bullet \\ \diagdown & \diagup \\ \bullet & \bullet \end{array} \right) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

and the identity tensor

$$\psi \left(\begin{array}{cc} \bullet & \bullet \\ \text{---} & \text{---} \\ \bullet & \bullet \end{array} \right) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$


For the symmetric group \mathfrak{S}_n , spanning the symmetric group algebra \mathbb{S}_n , this action corresponds to the permutation of the tensor positions, i.e. for all $\sigma \in \mathfrak{S}_n$ and all v_1, \dots, v_n in \mathbb{C}^d , the tensor representation of σ on $(\mathbb{C}^d)^{\otimes n}$ gives the action

$$\psi(\sigma) \cdot (v_1 \otimes \dots \otimes v_n) = v_{\sigma^{-1}(1)} \otimes \dots \otimes v_{\sigma^{-1}(n)}.$$


The partial transposition operation on a row, for a given diagram p , corresponds to the partial transposition of a tensor, for the matrix $\psi(p)$. For example, let the partition $12 | 34$ in the Brauer algebra $\mathbb{B}_2(2)$, then taking the partial transposition on the 1-st row gives:

$$\psi \left(\begin{array}{cc} \bullet & \bullet \\ \curvearrowright & \curvearrowleft \\ \bullet & \bullet \end{array} \right) = \psi \left(\begin{array}{cc} \bullet & \bullet \\ \diagdown & \diagup \\ \bullet & \bullet \end{array} \right) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}^T.$$

The operation referred to as **closing** a diagram consists in connecting each vertex of the diagram, to the vertex located in the same row, yielding a collection of loops. The trace of the tensor representation on $(\mathbb{C}^d)^{\otimes n}$, for a diagram p , can be obtained by $\text{Tr} [\psi(p)] = d^l$, where l is the number of loops after closing the diagram p . For example, let the partition $p := 1\ 2\ 4\ 5 \mid 3\ 6$ in the uniform block permutation algebra $\mathbb{U}_3(d)$,



the closing of p is



with 2 loops, then the trace of $\psi(p)$ becomes $\text{Tr} [\psi(p)] = d^2$.

The tensor representation of a diagrammatic algebra is in general non-faithfull. For example, let \mathbb{S}_3 be the symmetric group algebra with its tensor representation on $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ given by the map ψ , and define **sign**(σ) to be a **signature** of the permutation $\sigma \in \mathfrak{S}_3$, then

$$\sum_{\sigma \in \mathfrak{S}_3} \text{sign}(\sigma) \cdot \psi(\sigma) = 0.$$

Remark. In certain cases, diagrammatic algebras may not exhibit semisimplicity. However, the algebra \mathcal{A} defined by the complex span of $\psi(p)$, for all p in some diagrammatic monoid, always exhibits semisimplicity, as a matrix algebra:

$$\mathcal{A} \simeq \mathcal{M}_{d_1}^{\oplus n_1} \oplus \cdots \oplus \mathcal{M}_{d_k}^{\oplus n_k},$$

with some multiplicities n_i and dimensions d_i . In this basis, an element $A \in \mathcal{A}$ is written,

$$A \simeq I_{n_1} \otimes A_1 \oplus \cdots \oplus I_{n_k} \otimes A_k.$$

1.3 Schur-Weyl dualities

1.3.1 Commutant

Given a matrix algebra $\mathcal{A} \subseteq \mathcal{M}_d$, the **commutant** of \mathcal{A} , denoted \mathcal{A}' , is the set of matrices that commute with all elements of \mathcal{A} :

$$\mathcal{A}' := \{M \in \mathcal{M}_d \mid MA = AM, \text{ for all } A \in \mathcal{A}\}.$$

Theorem 1.1 ([Ser+77; FH13]). *Let \mathcal{A} be a matrix algebra, and \mathcal{B} the commutant of \mathcal{A} . Suppose \mathcal{A} decomposes as $\mathcal{A} \simeq \mathcal{M}_{d_1}^{\oplus n_1} \oplus \cdots \oplus \mathcal{M}_{d_k}^{\oplus n_k}$. Then for all $A \in \mathcal{A}$ and $B \in \mathcal{B}$,*

$$\begin{aligned} A &\simeq \bigoplus_{i=1}^k I_{n_i} \otimes A_i, \\ B &\simeq \bigoplus_{i=1}^k B_i \otimes I_{d_i}. \end{aligned}$$

Furthermore both \mathcal{A} and \mathcal{B} are commutants of each other, i.e. $\mathcal{B} = \mathcal{A}'$ and $\mathcal{A} = \mathcal{B}'$.

1.3.2 Schur-Weyl duality for \mathfrak{S}_n

Let GL_d be the **complex general linear group** of degree d , which consists of the $d \times d$ invertible complex matrices acting on \mathbb{C}^d . This action extends diagonally to an action on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, defined for $M \in \text{GL}_d$ on tensor $v_1 \otimes \cdots \otimes v_n \in (\mathbb{C}^d)^{\otimes n}$ by,

$$M^{\otimes n} \cdot (v_1 \otimes \cdots \otimes v_n) = M \cdot v_1 \otimes \cdots \otimes M \cdot v_n,$$

and extended linearly.

Let \mathcal{A} and \mathcal{B} be the matrix algebras generated, respectively, by the actions of the symmetric group \mathfrak{S}_n and the complex general linear group GL_d , on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, i.e.

$$\begin{aligned} \mathcal{A} &:= \text{Span}_{\mathbb{C}} \{\psi(\sigma) \mid \sigma \in \mathfrak{S}_n\} \\ \mathcal{B} &:= \text{Span}_{\mathbb{C}} \{M^{\otimes n} \mid M \in \text{GL}_d\}. \end{aligned}$$

Theorem ([GW09]). *Both \mathcal{A} and \mathcal{B} are commutants of each other.*

The matrix algebra \mathcal{A} , generated by the tensor representation of the symmetric group \mathfrak{S}_n , can be decomposed as the direct sum,

$$\mathcal{A} \simeq \bigoplus_{\substack{\lambda \vdash n \\ \lambda'_1 \leq d}} \mathcal{M}_{d_\lambda}^{\oplus n_\lambda},$$

indexed by the Young diagrams λ with n boxes and at most d rows.¹ Then according to [Theorem 1.1](#), for all $A \in \mathcal{A}$ and $B \in \mathcal{B}$,

$$A \simeq \bigoplus_{\substack{\lambda \vdash n \\ \lambda'_1 \leq d}} I_{n_\lambda} \otimes A_\lambda \quad \text{and} \quad B \simeq \bigoplus_{\substack{\lambda \vdash n \\ \lambda'_1 \leq d}} B_\lambda \otimes I_{d_\lambda},$$

where the A_λ act on a space denoted V_λ , and the B_λ act on a space denoted V_λ^d .

Theorem 1.2 (Schur-Weyl duality [[Sch27](#); [Wey46](#)]). *The space of n -fold tensors over \mathbb{C}^d decomposes under the action of the direct product group $\text{GL}_d \times \mathfrak{S}_n$ as follows:*

$$(\mathbb{C}^d)^{\otimes n} \simeq \bigoplus_{\substack{\lambda \vdash n \\ \lambda'_1 \leq d}} V_\lambda^d \otimes V_\lambda.$$

The diagonal action of GL_d , on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, can be restricted to the subgroup of the **unitary group** of degree d , denoted U_d , which consists of the $d \times d$ unitary matrices acting on \mathbb{C}^d .

Theorem 1.3. *Let \mathcal{C} be the matrix algebra generated by the action of the unitary group U_d , on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, i.e. $\mathcal{C} := \text{Span}_{\mathbb{C}} \{U^{\otimes n} \mid U \in U_d\}$. Then*

$$\mathcal{B} \simeq \mathcal{C}.$$

Proof. The present proof is from an unpublished note by Guillaume Aubrun [[Aub18](#)]. Due to the inclusion $U_d \subseteq \text{GL}_d$, it follows that $\mathcal{C} \subseteq \mathcal{B}$. To establish the result, it suffices to prove that for $M \in \text{GL}_d$, the n -fold $M^{\otimes n}$ can be expressed as a limit of linear combinations of n -fold $U^{\otimes n}$, for some $U \in U_d$.

Without loss of generality, assume that M can be multiplied by a real scalar

1. see [Appendix A](#)

to obtain a singular value decomposition given by,

$$M = \sum_{i=1}^d s_i |e_i \rangle \langle f_i|,$$

where e_1, \dots, e_d and f_1, \dots, f_d are two orthonormal bases of \mathbb{C}^d , and s_i are non-negative real numbers satisfying $-1 < s_i < 1$. Notice that the matrix obtained by replacing each s_i with a complex number z_i satisfying $|z_i| = 1$ is unitary. Then

$$\begin{aligned} M^{\otimes n} &= \left(\sum_{i=1}^d s_i |e_i \rangle \langle f_i| \right)^{\otimes n} \\ &= \sum_{i_1, \dots, i_n=1}^d \prod_{j=1}^n s_{i_j} |e_{i_1} \otimes \dots \otimes e_{i_n} \rangle \langle f_{i_1} \otimes \dots \otimes f_{i_n}|. \end{aligned}$$

Let γ be a counterclockwise unit circle in the complex plane. It is well-known from Cauchy's formula, that for all $s \in \mathbb{R}$ such that $-1 < s < 1$, and for all $k \in \mathbb{N}$,

$$s^k = \frac{1}{2\pi i} \int_{\gamma} z^k \frac{dz}{z-s}.$$

Then, using Fubini's theorem, for all $s \in \mathbb{R}^d$ such that $-1 < s_i < 1$, and for all $i_1, \dots, i_n \in \{1, \dots, d\}$,

$$\prod_{j=1}^n s_{i_j} = \frac{1}{(2\pi i)^d} \int_{\gamma^{\times d}} \prod_{j=1}^n z_{i_j} \frac{dz_1}{z_1 - s_1} \dots \frac{dz_d}{z_d - s_d}.$$

Finally,

$$\begin{aligned} &M^{\otimes n} \\ &= \sum_{i_1, \dots, i_n=1}^d \prod_{j=1}^n s_{i_j} |e_{i_1} \otimes \dots \otimes e_{i_n} \rangle \langle f_{i_1} \otimes \dots \otimes f_{i_n}| \\ &= \sum_{i_1, \dots, i_n=1}^d \frac{1}{(2\pi i)^d} \int_{\gamma^{\times d}} \prod_{j=1}^n z_{i_j} \frac{dz_1}{z_1 - s_1} \dots \frac{dz_d}{z_d - s_d} |e_{i_1} \otimes \dots \otimes e_{i_n} \rangle \langle f_{i_1} \otimes \dots \otimes f_{i_n}| \\ &= \frac{1}{(2\pi i)^d} \int_{\gamma^{\times d}} U_{z_1, \dots, z_d}^{\otimes n} \frac{dz_1}{z_1 - s_1} \dots \frac{dz_d}{z_d - s_d}, \end{aligned}$$

where $U_{z_1, \dots, z_d}^{\otimes n}$ is the n -fold unitary matrix defined by

$$U_{z_1, \dots, z_d} := \sum_{i=1}^d z_i |e_i\rangle\langle f_i|.$$

Corollary. *The space of n -fold tensors over \mathbb{C}^d decomposes under the action of the direct product group $U_d \times \mathfrak{S}_n$ as follows:*

$$(\mathbb{C}^d)^{\otimes n} \simeq \bigoplus_{\substack{\lambda \vdash n \\ \lambda'_1 \leq d}} V_\lambda^d \otimes V_\lambda.$$

1.3.3 Other Schur-Weyl dualities

Schur-Weyl duality for \mathbb{P}_n

The symmetric group \mathfrak{S}_d acts on the d -dimensional complex vector space \mathbb{C}^d by considering the mapping ϕ , called **permutation matrix**, from \mathfrak{S}_d to \mathcal{M}_d , and defined for each permutation σ by,

$$(\phi(p))_{i,j} := \begin{cases} 1 & \text{if } i = \sigma(j) \\ 0 & \text{otherwise,} \end{cases}$$

To be explicit, given a basis e_1, \dots, e_d of \mathbb{C}^d and vector $v = \sum_{i=1}^d v_i |e_i\rangle$ in \mathbb{C}^d , a permutation $\sigma \in \mathfrak{S}_d$ acts on v by

$$\phi(\sigma) \cdot v = \sum_{i=1}^d v_{\sigma^{-1}(i)} |e_i\rangle.$$

This action extends diagonally to an action on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, defined for $\sigma \in \mathfrak{S}_d$ on tensor $v_1 \otimes \dots \otimes v_n \in (\mathbb{C}^d)^{\otimes n}$ by,

$$\phi(\sigma)^{\otimes n} \cdot (v_1 \otimes \dots \otimes v_n) = \phi(\sigma) \cdot v_1 \otimes \dots \otimes \phi(\sigma) \cdot v_n,$$

and extended linearly.

Let \mathcal{A} and \mathcal{B} be the matrix algebras generated, respectively, by the actions of the partition monoid \mathbb{P}_n and the symmetric group \mathfrak{S}_d , on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, i.e.

$$\begin{aligned} \mathcal{A} &:= \text{Span}_{\mathbb{C}} \{ \psi(p) \mid p \in \mathbb{P}_n \} \\ \mathcal{B} &:= \text{Span}_{\mathbb{C}} \{ \phi(\sigma)^{\otimes n} \mid \sigma \in \mathfrak{S}_d \}. \end{aligned}$$

■ **Theorem** ([MR98; Mar00; HR05]). *Both \mathcal{A} and \mathcal{B} are commutants of each other.*

Schur-Weyl duality for \mathbb{U}_n

Let **diag. \mathbb{U}_d** be the subgroup of \mathbb{U}_d consisting of $d \times d$ **diagonal unitary matrices** acting on \mathbb{C}^d . That is for all U in **diag. \mathbb{U}_d** , there is $\theta \in [0, 2\pi)^d$ such that,

$$U = \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_d}).$$

As subgroups of \mathbb{U}_d , the action of **diag. \mathbb{U}_d** extends diagonally on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$.

The product of a diagonal unitary matrix $U := \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_d})$ in **diag. \mathbb{U}_d** , and a permutation matrix $\phi(\sigma)$ for some $\sigma \in \mathfrak{S}_d$, is a **monomial matrix** in $[0, 2\pi)$, i.e. a permutation matrix whose nonzero components are $e^{i\theta}$ with $\theta \in [0, 2\pi)$:

$$(U \cdot \phi(\sigma))_{i,j} = \begin{cases} \theta_i & \text{if } i = \sigma(j) \\ 0 & \text{otherwise,} \end{cases}$$

Let \mathcal{A} and \mathcal{B} be the matrix algebras generated, respectively, by the actions of the uniform block permutation monoid \mathbb{U}_n and the monomial matrices in $[0, 2\pi)$, on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, i.e.

$$\begin{aligned} \mathcal{A} &:= \text{Span}_{\mathbb{C}} \{ \psi(p) \mid p \in \mathbb{U}_n \} \\ \mathcal{B} &:= \text{Span}_{\mathbb{C}} \{ U^{\otimes n} \cdot \phi(\sigma)^{\otimes n} \mid U \in \text{diag. } \mathbb{U}_d \text{ and } \sigma \in \mathfrak{S}_d \}. \end{aligned}$$

■ **Theorem 1.4** ([Tan97]). *Both \mathcal{A} and \mathcal{B} are commutants of each other.*

Schur-Weyl duality for \mathbb{B}_n

Let **\mathbb{O}_d** denote the **ortogonal group** of degree d , which consists of the $d \times d$ orthogonal matrices acting on \mathbb{C}^d . As subgroups of \mathbb{U}_d , the action of **\mathbb{O}_d** extends diagonally on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$.

Let \mathcal{A} and \mathcal{B} be the matrix algebras generated, respectively, by the actions of the Brauer monoid \mathbb{B}_n and the orthogonal group **\mathbb{O}_d** , on the d^n -dimensional tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, i.e.

$$\begin{aligned} \mathcal{A} &:= \text{Span}_{\mathbb{C}} \{ \psi(p) \mid p \in \mathbb{B}_n \} \\ \mathcal{B} &:= \text{Span}_{\mathbb{C}} \{ O^{\otimes n} \mid O \in \mathbb{O}_d \}. \end{aligned}$$

■ **Theorem** ([Bra37]). *Both \mathcal{A} and \mathcal{B} are commutants of each other.*

Schur-Weyl duality for $\mathbb{B}_{m,n}$

Let the action of the complex general linear group GL_d on the d^n -dimensional mixed tensor product complex vector space $(\mathbb{C}^d)^{\otimes n} \otimes (\mathbb{C}^d)^{\otimes m}$, defined for $M \in \mathrm{GL}_d$ by,

$$M^{\otimes m} \otimes \left((M^{-1})^\top \right)^{\otimes n}.$$

As a subgroup of GL_d , this action is defined for $U \in \mathrm{U}_d$ by,

$$U^{\otimes m} \otimes \bar{U}^{\otimes n}.$$

Let \mathcal{A}, \mathcal{B} and \mathcal{C} be the matrix algebras generated, respectively, by the actions of the walled Brauer monoid $\mathbb{B}_{m,n}$, the complex general linear group GL_d and the unitary group U_d , on the d^n -dimensional mixed tensor product complex vector space $(\mathbb{C}^d)^{\otimes n}$, i.e.

$$\begin{aligned} \mathcal{A} &:= \mathrm{Span}_{\mathbb{C}} \{ \psi(p) \mid p \in \mathbb{B}_n \} \\ \mathcal{B} &:= \mathrm{Span}_{\mathbb{C}} \left\{ M^{\otimes m} \otimes \left((M^{-1})^\top \right)^{\otimes n} \mid M \in \mathrm{GL}_d \right\} \\ \mathcal{C} &:= \mathrm{Span}_{\mathbb{C}} \{ U^{\otimes m} \otimes \bar{U}^{\otimes n} \mid U \in \mathrm{U}_d \}. \end{aligned}$$

Theorem ([Ben+94]). *Both \mathcal{A} and \mathcal{B} (or \mathcal{C}) are commutants of each other.*

Remark. It is important to note that the various Schur-Weyl dualities presented in [Section 1.3](#) are given only in terms of the matrix algebras generated by the map ψ , rather than the diagrammatic algebras. The map ψ may not always exhibit faithfulness.

Quantum Information Theory

Chapter contents

2.1	Postulates of quantum mechanics	31
2.1.1	Pure states	31
2.1.2	Mixed states	33
2.2	Quantum entanglement	34
2.2.1	Schmidt decomposition	34
2.2.2	Pure quantum state entanglement	35
2.2.3	Mixed quantum state entanglement	36
2.2.4	Monogamy of entanglement	37
2.3	Quantum channels	38
2.3.1	Structure of quantum channels	38
2.3.2	Compatibility of quantum channels	39
2.4	Quantum fidelity	39
2.5	Graphical calculus	40

This chapter provides a comprehensive overview of the mathematical foundations of quantum mechanics in the context of quantum information theory, focusing on the postulates of quantum mechanics and their inherent probabilistic nature.

In quantum information theory, the attention is mainly directed to quantum systems with a finite number of degrees of freedom.

First, the formalism of pure quantum states is introduced, which is particularly suitable for closed quantum systems. Then the formalism of mixed quantum states is then introduced, in particular to describe open quantum systems that interact with an environment that is not intended to be described.

References for the different postulates and the mathematical foundations of quantum mechanics can be found in the textbooks [Wat18; NC02; AS17].

2.1 Postulates of quantum mechanics

Let $\mathcal{H} := \mathbb{C}^d$ be a finite d -dimensional Hilbert complex vector space. The convex set of unit trace, positive semi-definite $d \times d$ matrices, acting on \mathcal{H} is denoted,

$$\mathcal{D}_d := \{\rho \in \mathcal{M}_d \mid \text{Tr } \rho = 1 \text{ and } \rho \geq 0\}.$$

An element of \mathcal{D}_d is called a **density matrix**, to highlight the fact that its eigenvalues represent a probability distribution. The extremal points of \mathcal{D}_d are the unit rank projections $|\psi\rangle\langle\psi|$, for some $\psi \in \mathcal{H}$ with $\|\psi\| = 1$.

Definition. A **quantum system** is represented by a finite d -dimensional Hilbert complex vector space \mathcal{H} .

Let $\mathcal{H} := \mathbb{C}^d$ be a finite d -dimensional Hilbert complex vector space, the computational basis of the quantum system \mathcal{H} is denoted:

$$|0\rangle, \dots, |d-1\rangle.$$

Definition. A **composite quantum system** is represented by a tensor product of Hilbert complex vector spaces $\mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_n$.

The computational basis of the n -fold composite quantum system $\mathcal{H}^{\otimes n}$, with the finite d -dimensional Hilbert complex vector space $\mathcal{H} := \mathbb{C}^d$, is the set:

$$\{ |i_1\rangle \otimes \dots \otimes |i_n\rangle \mid i_1, \dots, i_n \in \{0, \dots, d-1\} \}.$$

2.1.1 Pure states

Definition. A **pure quantum state** on \mathcal{H} is an extremal point of \mathcal{D}_d .

A pure quantum state on $\mathcal{H} := \mathbb{C}^2$ is a unit rank projection $|\psi\rangle\langle\psi|$, for some vector $|\psi\rangle := \alpha \cdot |0\rangle + \beta \cdot |1\rangle$ with α, β satisfying,

$$|\alpha|^2 + |\beta|^2 = 1.$$

Remark. In the following the description of a pure quantum state as a unit rank projection $|\psi\rangle\langle\psi|$ or as a unit norm vector $|\psi\rangle$ is used interchangeably. Note that $|\psi\rangle\langle\psi|$ is simply the orthogonal projection onto the complex line spanned by $|\psi\rangle$.

Definition. The **evolution** of a pure quantum state ρ on \mathcal{H} is governed by a unitary matrix U on \mathcal{H} , through the conjugation mapping

$$\rho \mapsto U\rho U^*.$$

The evolution of pure quantum states is a transitive action: for all pure quantum states ρ and σ there exists a unitary matrix U such that $\rho = U\sigma U^*$.

As a unit norm vector $|\psi\rangle$, the evolution a pure quantum state, through a unitary matrix U , is given by $|\psi\rangle \mapsto U|\psi\rangle$.

Definition. The **projective measure** of a pure quantum state ρ on \mathcal{H} is described by a set of orthogonal projections $\{P_1, \dots, P_n\}$ on \mathcal{H} , which sum to the identity. The **outcome** of the measure is $i \in \{1, \dots, n\}$ with probability,

$$\text{Tr}[P_i \rho P_i^*],$$

and the resulting pure quantum state after the measure becomes,

$$\frac{P_i \rho P_i^*}{\text{Tr}[P_i \rho P_i^*]}.$$

Let $|\psi\rangle := \alpha \cdot |0\rangle + \beta \cdot |1\rangle$ be a quantum pure state on $\mathcal{H} := \mathbb{C}^2$. A projective measure in the computational basis $|0\rangle, |1\rangle$ is described by the two orthogonal projections $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$, and yields outcome 0 with probability $|\alpha|^2$ and outcome 1 with probability $|\beta|^2$.

Remark. In the context of a projective measure, due to the orthogonality property of the projections $\{P_1, \dots, P_n\}$, and the cyclic property of the trace, the probability of the outcome i given a pure quantum state $\rho := |\psi\rangle\langle\psi|$ is just $\text{Tr}[P_i \rho] = \langle\psi| P_i |\psi\rangle$, and the resulting pure quantum state after the measure

becomes,

$$\frac{P_i \rho P_i}{\langle \psi | P_i | \psi \rangle}.$$

Let $\rho := \rho_A \otimes \rho_B$ be a pure quantum state on $\mathcal{H}_A \otimes \mathcal{H}_B$, a composite quantum system. The projective measure of ρ described by the set of orthogonal projections $\{P_1, \dots, P_n\}$ on $\mathcal{H}_A \otimes \mathcal{H}_B$, yields outcome $i \in \{1, \dots, n\}$ with probability

$$\begin{aligned} \text{Tr} [P_i \rho] &= \text{Tr}_A \left[\underbrace{P_i \text{Tr}_B [I_A \otimes \rho_B]}_{M_i} \rho_A \right] \\ &= \text{Tr} [M_i \rho_A], \end{aligned}$$

for some positive semidefinite M_i on $\mathcal{H}_A \otimes \mathcal{H}_B$, which sum to the identity. Such a measure on part of a composite system is called a **generalized measure**.

2.1.2 Mixed states

Let $\rho := |\psi\rangle\langle\psi|$ be a pure quantum state on $\mathcal{H}_A \otimes \mathcal{H}_B$, a composite quantum system with $\mathcal{H}_A := \mathbb{C}^d$ and \mathcal{H}_B an unknown quantum system. The projective measure of ρ described by the set of orthogonal projections $\{P_1, \dots, P_n\}$ on \mathcal{H}_A only, yields outcome $i \in \{1, \dots, n\}$ with probability

$$\begin{aligned} \langle \psi | (P_i \otimes I_B) | \psi \rangle &= \text{Tr}_A \left[P_i \underbrace{\text{Tr}_B [\rho]}_{\sigma} \right] \\ &= \text{Tr} [P_i \sigma], \end{aligned}$$

for some $\sigma \in \mathcal{D}_d$, which is in general not a pure quantum state.

Definition. A **mixed quantum state** on \mathcal{H} is an element of \mathcal{D}_d .

Since the set of mixed quantum state \mathcal{D}_d is a convex set, and since the extremal point are the pure quantum states, a mixed quantum state is a convex combination of pure quantum states, of the form,

$$\sum_{i=1}^k p_i \cdot |\psi_i\rangle\langle\psi_i|,$$

with some positive real numbers p_i satisfying $\sum_{i=1}^k p_i = 1$, and $|\psi_i\rangle\langle\psi_i|$ some unit rank projections. From the spectral Theorem, a mixed quantum state on \mathcal{D}_d is a

convex sum of at most d terms. The most central element of \mathcal{D}_d is the mixed quantum state $\mathbf{I} := \frac{I_d}{d}$ called **maximally mixed**.

As a fundamental consequence, maximizing a convex function or minimizing a concave function over the set \mathcal{D}_d of mixed quantum states will lead to extremal values of the function on a pure quantum state.

In the case $\mathcal{H} := \mathbb{C}^2$, the mixed quantum states \mathcal{D}_2 can be written in a spherical representation, called **Bloch sphere**,

$$\mathcal{D}_2 = \left\{ \frac{1}{2}(I_2 + r_x \cdot \sigma_x + r_y \cdot \sigma_y + r_z \cdot \sigma_z) \mid r := (r_x, r_y, r_z) \in \mathbb{R}^3 \text{ and } \|r\| \leq 1 \right\},$$

where σ_x, σ_y and σ_z are the **Pauli matrices** defined by

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The pure quantum states of \mathcal{D}_2 are the elements satisfying $\|r\| = 1$.

2.2 Quantum entanglement

The **quantum entanglement** is a fundamental concept in quantum information theory that refers to the non-classical correlations that exist between two or more quantum systems.

2.2.1 Schmidt decomposition

Recall the **singular value decomposition** (SVD) for a matrix $M \in \mathcal{M}_d$ acting on \mathbb{C}^d : there exists two orthonormal bases e_1, \dots, e_d and f_1, \dots, f_d for the vector space \mathbb{C}^d , and d non-negative real numbers s_1, \dots, s_d , such that,

$$M = \sum_{i=1}^d s_i |e_i\rangle\langle f_i|.$$

Using the isomorphism $\text{Hom}(V, W) \simeq V \otimes W$ between two finite dimensional complex vector spaces V and W , the singular value decomposition becomes the **Schmidt decomposition** of vector on a bipartite tensor product $V \otimes W$.

Theorem (Schmidt decomposition [Sch07; Eve57]). *Let ψ be a vector in a bipartite tensor product of d -dimensional Hilbert complex vector spaces $\mathcal{H}_1 \otimes \mathcal{H}_2$. Then there exists two orthonormal bases e_1, \dots, e_d and f_1, \dots, f_d for \mathcal{H}_1 and \mathcal{H}_2 ,*

respectively, and d non-negative real numbers s_1, \dots, s_d called **Schmidt coefficients**, such that,

$$\psi = \sum_{i=1}^d s_i \cdot e_i \otimes f_i.$$

The number of nonzero Schmidt coefficients is called the **Schmidt number**.

Apart from the bipartite scenario, direct multipartite extension of the Schmidt decomposition does not exist in general [Per95].

2.2.2 Pure quantum state entanglement

A bipartite pure quantum state $|\psi\rangle$ on $\mathcal{H}_A \otimes \mathcal{H}_B$ is said to be **entangled** if its Schmidt number is strictly greater than 1, otherwise it is said to be **separable** and can be written,

$$|\psi\rangle = |\psi\rangle_A \otimes |\psi\rangle_B,$$

for some $|\psi\rangle_A \in \mathcal{H}_A$ and $|\psi\rangle_B \in \mathcal{H}_B$.

The Schmidt number of a bipartite pure quantum state gives a canonical quantitative **measure of entanglement**.

A bipartite pure quantum state $\omega := |\Omega\rangle\langle\Omega|$, on the 2-fold composite quantum system $\mathcal{H} \otimes \mathcal{H}$, with $\mathcal{H} \simeq \mathbb{C}^d$, is called **maximally entangled** if it has, on the computational basis, the form

$$|\Omega\rangle = \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |ii\rangle.$$

The partial transpose of an unnormalized maximally entangled pure quantum state $d \cdot \omega$ is the **flip operator**,

$$d \cdot \omega^\Gamma = \sum_{i,j=0}^{d-1} |ij\rangle\langle ij|,$$

on the computational basis. On product vector $x \otimes y$, the flip operator acts through $d \cdot \omega^\Gamma(x \otimes y) = y \otimes x$. However, since $y \otimes x - x \otimes y$ is a eigenvector for the negative eigenvalue -1 , the flip operator ω^Γ is not a quantum state. The normalized flip operator on \mathcal{D}_{d^2} is denoted $\mathbf{F} := \frac{\omega^\Gamma}{d}$.

Theorem (PPT criterion [Per96; HHH01]). *If ρ is a separable bipartite pure quantum state on $\mathcal{H}_A \otimes \mathcal{H}_B$, then the partially transposed ρ^Γ is a quantum state. The converse is true if and only if $\dim \mathcal{H}_A \times \mathcal{H}_B \leq 6$.*

In general, in the multipartite scenario $\mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_k$, a pure quantum state $|\psi\rangle$ is separable if it can be written as a product vector

$$|\psi\rangle = |\psi_1\rangle \otimes \cdots \otimes |\psi_k\rangle,$$

and entangled otherwise.

Remark. As a unit rank projection $|\psi\rangle\langle\psi|$, a separable pure quantum state on a multipartite $\mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_k$ can be written

$$|\psi\rangle\langle\psi| = |\psi_1\rangle\langle\psi_1| \otimes \cdots \otimes |\psi_k\rangle\langle\psi_k|.$$

2.2.3 Mixed quantum state entanglement

A mixed quantum state is said to be separable if it can be written as a convex combination of separable pure quantum states. Therefore the convex set of separable quantum states on $\mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_k$ is

$$\text{Conv} \{ |\psi_1\rangle\langle\psi_1| \otimes \cdots \otimes |\psi_k\rangle\langle\psi_k| : \psi_i \in \mathcal{H}_i \text{ for all } i \in \{1, \dots, k\} \},$$

with extremal points the separable pure quantum states.

An **isotropic state** is a convex combination of a maximally entangled and maximally mixed quantum states:

$$\rho = \lambda \cdot \omega + (1 - \lambda)\text{I},$$

with $0 \leq \lambda \leq 1$ and $\omega, \text{I} \in \mathcal{D}_{d^2}$. If $\frac{-1}{d^2-1} \leq \lambda \leq 0$, then ρ is still in \mathcal{D}_{d^2} , and thus still a mixed quantum state.

A **Werner state** is an affine combination of a normalized flip operator and maximally mixed quantum states:

$$\sigma = \lambda \cdot \text{F} + (1 - \lambda)\text{I},$$

with $\frac{1}{1-d} \leq \lambda \leq \frac{1}{1+d}$ and $\text{F}, \text{I} \in \mathcal{D}_{d^2}$.

Any isotropic state ρ and Werner state σ satisfy the following commutation relations:

$$[\rho, \bar{U} \otimes U] = 0 \quad \text{and} \quad [\sigma, U \otimes U] = 0,$$

for all unitary matrices U .

Theorem ([HH99]). Let $\rho := \lambda \cdot \omega + (1 - \lambda)\text{I}$ be an isotropic state on \mathcal{D}_{d^2} . Then ρ is separable if and only if $\lambda \leq \frac{1}{d+1}$.

Theorem ([Wer89]). *Let $\sigma := \lambda \cdot F + (1 - \lambda)I$ be a Werner state, on \mathcal{D}_{d^2} . Then σ is separable if and only if $\lambda \geq \frac{1}{1-d^2}$.*

In general, deciding whether a given quantum state is separable is known to be NP-hard [Gur03].

2.2.4 Monogamy of entanglement

A bipartite quantum state ρ on $\mathcal{H}_A \otimes \mathcal{H}_B$ is said to be k -extendible, with respect to \mathcal{H}_B if there exists a quantum state σ on $\mathcal{H}_A \otimes \mathcal{H}_B^{\otimes k}$ such that for all $i \in \{1, \dots, k\}$:

$$\rho = \text{Tr}_{\{B_1, \dots, B_k\} \setminus \{B_i\}} [\sigma].$$

Theorem (Entanglement hierarchy [DPS04]). *A bipartite quantum state ρ on $\mathcal{H}_A \otimes \mathcal{H}_B$ is separable if and only if it is k -extendible for all $k \in \mathbb{N}$.*

Let $\rho_{A,B,C}$ be a tripartite quantum state on $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ with $\mathcal{H}_A \simeq \mathcal{H}_B \simeq \mathcal{H}_C$, and such that the **reduced quantum state** on $\mathcal{H}_A \otimes \mathcal{H}_B$,

$$\rho_{A,B} := \text{Tr}_C[\rho_{A,B,C}],$$

is maximally entangled, i.e. $\rho_{A,B} = \omega$. From the spectral Theorem, the quantum state $\rho_{A,B,C}$ is a convex combination,

$$\rho_{A,B,C} = \sum_{i=1}^k p_i \cdot |\psi_i\rangle\langle\psi_i|,$$

for some orthonormal pure quantum states $|\psi_i\rangle$ on $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$. Then the reduced quantum state $\rho_{A,B}$ becomes

$$\rho_{A,B} = \sum_{i=1}^k p_i \cdot \text{Tr}_C \left[|\psi_i\rangle\langle\psi_i| \right].$$

But since $\rho_{A,B}$ is a pure quantum state, i.e. $\rho_{A,B} = |\Omega\rangle\langle\Omega|$ is an extremal point of the convex set of density matrices, every $\text{Tr}_C \left[|\psi_i\rangle\langle\psi_i| \right]$ in the sum must necessarily be equal to $\rho_{A,B}$. This implies that $\rho_{A,B,C} = \rho_{A,B} \otimes \rho_C$, for some reduced quantum state ρ_C on \mathcal{H}_C . Thus none of the reduced quantum states $\rho_{A,C}$ and $\rho_{B,C}$ can be maximally entangled. This phenomenon is known as **monogamy of entanglement**.

2.3 Quantum channels

Let $\Phi : \mathcal{M}_d \rightarrow \mathcal{M}_{d'}$ be a map such that $\Phi(\mathcal{D}_d) \subseteq \mathcal{D}_{d'}$, i.e. mapping d -dimensional quantum states to d' -dimensional quantum states. Under linearity assumption this is equivalent to:

- Φ **positive**: $X \geq 0 \implies \Phi(X) \geq 0$.
- Φ **trace preserving**: $\text{Tr } X = \text{Tr } \Phi(X)$.

The transpose of a matrix is a linear map that satisfies both positivity and trace preserving properties. But when partially applied to a composite quantum system, this can lead to non-quantum states, e.g. the partial transpose of a maximally entangled pure quantum state is the flip operator $F = \omega^\Gamma$.

A linear map $\Phi : \mathcal{M}_d \rightarrow \mathcal{M}_{d'}$ is called **completely positive** if all the partial applications of Φ on any positive semidefinite matrix results in another positive semidefinite matrix, i.e. $\forall D \in \mathbb{N}, \mathcal{M}_d \otimes \mathcal{M}_D \ni X \geq 0 \implies (\Phi \otimes \text{id}_D)(X) \geq 0$.

Definition. The most general transformations of quantum states, called **quantum channels**, are the Completely Positive Trace Preserving (CPTP) linear maps.

2.3.1 Structure of quantum channels

The **Choi matrix** of a linear map $\Phi : \mathcal{M}_d \rightarrow \mathcal{M}_{d'}$ is the matrix C_Φ in $\mathcal{M}_{d \times d'}$, defined by,

$$\begin{aligned} C_\Phi &:= (\text{id}_d \otimes \Phi) \left(\sum_{i,j=1}^d |ii\rangle\langle jj| \right) \\ &= (\text{id}_d \otimes \Phi)(d \cdot \omega). \end{aligned}$$

It is possible to retrieve the original linear map Φ from its associated Choi matrix C_Φ using the formula,

$$\Phi(X) = \text{Tr}_d [C_\Phi (X^\top \otimes I_{d'})].$$

Theorem 2.1 ([Wat18]). *Let $\Phi : \mathcal{M}_d \rightarrow \mathcal{M}_{d'}$ be a linear map. The following are equivalent:*

- the map Φ is CPTP;
- the Choi matrix C_Φ is positive semidefinite and $\text{Tr}_{d'} [C_\Phi] = I_d$;

— there exist $A_1, \dots, A_k \in \mathcal{M}_{d \times d'}$ such that,

$$\Phi(X) = \sum_{i=1}^k A_i X A_i^* \quad \text{and} \quad \sum_{i=1}^k A_i^* A_i = I_d;$$

— there exist $D \in \mathbb{N}$ and an isometry $V : \mathbb{C}^d \rightarrow \mathbb{C}^{d'} \otimes \mathbb{C}^D$ such that,

$$\Phi(X) = \text{Tr}_D [V X V^*].$$

2.3.2 Compatibility of quantum channels

Let $\Phi : \mathcal{M}_d \rightarrow \mathcal{M}_{d'}^{\otimes n}$ be a quantum channel from 1 to n quantum states, the i -th **marginal** of Φ , denoted Φ_i is defined by,

$$\Phi_i(X) := \text{Tr}_{[n] \setminus \{i\}} [\Phi(X)].$$

A marginal of a quantum channel is also a quantum channel.

Let $\Phi_i : \mathcal{M}_d \rightarrow \mathcal{M}_{d_i}$ be a family of k quantum channels. The **quantum channel compatibility problem** consists in determining whether there exists a global quantum channel $\Psi : \mathcal{M}_d \rightarrow \mathcal{M}_{d_1} \otimes \dots \otimes \mathcal{M}_{d_k}$ **compatible** with all the Φ_i , that is,

$$\Psi_i = \Phi_i,$$

for all marginals Ψ_i .

■ *Remark.* Quantum channels can be incompatible with themselves.

2.4 Quantum fidelity

The **quantum fidelity** is a measure of the closeness between two quantum states ρ and σ , defined as the function F on $\mathcal{D}_d \times \mathcal{D}_d$ by,

$$F(\rho, \sigma) = \text{Tr} \left[\sqrt{\sqrt{\sigma} \rho \sqrt{\sigma}} \right]^2.$$

■ **Proposition 2.2** ([Wat18]). *The quantum fidelity F between two quantum states has the following properties:*

- $F(\rho, \sigma) = F(\sigma, \rho)$;
- $F(\rho, \sigma) \in [0, 1]$;
- $F(\rho, \sigma) = 1 \iff \rho = \sigma$;


- $F(\rho, |\psi\rangle\langle\psi|) = \langle\psi|\rho|\psi\rangle = \text{Tr}[\rho|\psi\rangle\langle\psi|]$;
- $F(U\rho U^*, U\sigma U^*) = F(\rho, \sigma)$, for all unitary matrices U ;
- $F(\Phi(\rho), \Phi(\sigma)) \geq F(\rho, \sigma)$, for all quantum channels Φ ;
- F is jointly concave, i.e. for all $\lambda \in [0, 1]$,

$$F(\lambda \cdot \rho_1 + (1 - \lambda)\rho_2, \lambda \cdot \sigma_1 + (1 - \lambda)\sigma_2) \geq \lambda \cdot F(\rho_1, \sigma_1) + (1 - \lambda) \cdot F(\rho_2, \sigma_2).$$

2.5 Graphical calculus

The present Section introduces a graphical calculus for tensors, which is built upon the graphical notation developed by Penrose [Pen71]. Recently, analogous calculi have been formulated within the tensor network states framework and the framework of categorical quantum information theory, which are elaborated in [WBC15; BC17; CK17]. The graphical calculus introduced here is consistent with the tensor representation of a diagram algebra, introduced in Section 1.2.

In this graphical notation, tensors are represented by **boxes** and **wires**,



More specifically, the wires are labeled by **indices**, such that a box represents the value of the tensor at the given indices,

$$T_{ij} = \text{---}^i \boxed{T} \text{---}^j$$

In particular, a vector is a box with only 1 wire pointing to the left, and labeled by the index of the coordinate. A dual vector has its wire pointing to the right,

$$v_i = \text{---}^i \boxed{v} \qquad \bar{v}_i = \boxed{v^*} \text{---}^i$$

Note that these left and right directions are just a convention corresponding to the usual **right-to-left** composition in linear algebra.

The tensor diagrams can be combined in two ways. The **tensor product** combines two diagrams vertically

$$\begin{array}{c} \text{---} \boxed{A \otimes B} \text{---} \end{array} = \begin{array}{c} \text{---} \boxed{A} \text{---} \\ \text{---} \boxed{B} \text{---} \end{array}$$

or horizontally, with the isomorphism $\text{Hom}(V, W) \simeq V^* \otimes W$ between two finite dimensional complex vector spaces V and W ,

$$\text{---} \boxed{uv^*} \text{---} = \text{---} \boxed{u} \quad \boxed{v^*} \text{---}$$

The operation of joining together two diagrams by one of their wire labeled with the same index corresponds to multiplying the values of the two tensors at the given indices,

$$u_i \cdot v_i = \boxed{u} \overset{i}{\text{---}} \boxed{v}$$

The **tensor contraction** combines two diagrams using the previous operation by taking the sum over all common indices

$$\begin{array}{c} \diagup \boxed{A} \diagdown \\ \text{---} \boxed{A} \text{---} \end{array} \overset{\Sigma_i}{\text{---}} \begin{array}{c} \diagup \boxed{B} \diagdown \\ \text{---} \boxed{B} \text{---} \end{array}$$

In particular, this leads to the scalar product,

$$\langle u, v \rangle = \boxed{u} \text{---} \boxed{v}$$

the matrix product,

$$A \cdot B = \text{---} \boxed{A} \text{---} \boxed{B} \text{---}$$

and the matrix trace,

$$\text{Tr } T = \text{---} \boxed{T} \text{---}$$

Scalars multiply diagrams and are depicted next to them. The following three special tensors of \mathbb{C}^d , important in quantum information theory, have wire-only diagrams,

$$I_d = \text{---} \quad |\Omega\rangle = \frac{1}{\sqrt{d}} \cdot \text{---} \quad d = \bigcirc$$

Finally, the **matrix transpose** can be graphically depicted by swapping the *input* and the *output* wires of a box depicting a matrix. The partial transpose permutes just the wires of the corresponding spaces,

$$\text{---} \boxed{T^\top} \text{---} = \text{---} \boxed{T} \text{---}$$

Quantum Cloning

Chapter contents

3.1	Quantum cloning problem	45
3.1.1	Universal quantum cloning problem	50
3.1.2	Equatorial quantum cloning problem	51
3.1.3	Average vs. worst fidelity	53
3.2	Upper bound	54
3.3	Universal $1 \rightarrow 2$ quantum cloning problem	57
3.3.1	Restricted quantum cloning channels	60
3.3.2	General quantum cloning channels	63
3.4	Universal $1 \rightarrow N$ quantum cloning problem	65
3.4.1	Partially transposed permutations	65
3.4.2	Optimal symmetric quantum cloning channels	70
3.4.3	Optimal asymmetric quantum cloning channels	70
3.4.4	Optimal quantum cloning channel necessary condition	74
3.4.5	The \mathcal{Q} -norm	74
3.4.6	Achievable fidelity region	79

The present Section discusses the results of the papers “A geometrical description of the universal $1 \rightarrow 2$ asymmetric quantum cloning region” [NPR21], and “The asymmetric quantum cloning region” [NPR22], of which I am a co-author.

The problem of quantum cloning has received considerable attention over the last thirty years. This area of research began with the early investigations of universal quantum cloners [BH96] and has since expanded to include various cloning scenarios: symmetric [Wer98; KW99], asymmetric [Cer98; FFC05], qubits [GM97], qudits [Cer00], universal [Bru+98; IAG06; Ibl+05], equatorial [Bru+00; DM03; DD04b], economical [DD04a; DFC05], probabilistic [DG98; DG98], continuous quantum systems [Cer+04; Cer+05].

Of particular interest are two sets of papers dealing with the most general case of asymmetric universal $1 \rightarrow N$ quantum cloning problem. The first set of papers, by Kay and collaborators, deals with the optimisation of this problem [KRK12; Kay14; Kay16]. The second set of papers, by Ćwikliński, Horodecki, Mozrzyński, and Studziński, uses techniques from group representation theory [ĆHS12; SHM13; MHS14; Stu+14; MSH18].

In particular, Hashagen’s article on the asymmetric $1 \rightarrow 2$ quantum cloning problem [Has17], should be read in conjunction with the Section 3.3. Indeed, the quantum cloning problem is studied there using similar techniques: twirling of the quantum channel, Choi matrix as an element the algebra of operator permutations, decomposition this algebra into matrix algebras of size 2×2 and 1×1 . The main contribution of [NPR21] is the complete spectral analysis of the Choi matrix, which allows a block diagonalization of the matrix and a characterisation of the figures of merit as ellipses.

3.1 Quantum cloning problem

The problem known as the **quantum cloning problem** aims to identify a specific quantum channel, called the **quantum cloning channel**, denoted by $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$, which maps an *input* pure quantum state to an *output* mixed quantum state on a N -fold composite quantum system, such that the *output* marginals of Φ are as close as possible to the *input*. To achieve this, a **direction vector** denoted a satisfying $a \in [0, 1]^N$ and $\sum_{i=1}^N a_i = 1$, together with a subset Γ of the pure quantum states, are introduced. The quantum cloning problem is defined as the optimisation

problem given by,

$$\sup_{\Phi \in \text{CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F(\Phi_i(\rho), \rho) \right],$$

where the expected value is taken with respect to the uniform measure on Γ .

Remark. The quantum cloning problem belongs to a more general class of quantum marginal problems, which includes the quantum state marginal problem [Haa+21], the quantum channel marginal problem [HLA22], and the quantum measurement marginal problem [LN21].

The **no-cloning theorem** states that a perfect quantum cloning channel, i.e. a quantum channel with marginals the identity quantum channels, cannot exist in general. Indeed, a linearity argument shows that, if $|0\rangle, \dots, |d-1\rangle$ is the computational basis of \mathbb{C}^d , and $U \in \text{U}_{d^2}$ is a unitary matrix such that on this basis,

$$U |i\rangle \otimes |v\rangle = |i\rangle \otimes |i\rangle,$$

Then U is a perfect quantum cloning unitary for this basis, given an auxiliary pure quantum state $|v\rangle$. But if $|\psi\rangle := \frac{|i\rangle + |j\rangle}{\sqrt{2}}$ for distinct $i, j \in \{0, \dots, d-1\}$, then,

$$\begin{aligned} U |\psi\rangle \otimes |v\rangle &= \frac{|i\rangle \otimes |i\rangle + |j\rangle \otimes |j\rangle}{\sqrt{2}} \\ &\neq |\psi\rangle \otimes |\psi\rangle. \end{aligned}$$

Many quantum cryptographic schemes rely on the existence of the no-cloning theorem [BB84; BL20].

Theorem 3.1 (No-cloning theorem [WZ82; Die82]). *For any subset Γ of the pure quantum states, there is no quantum cloning channel $\Psi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ such that for all pure quantum states $\rho \in \Gamma$ and all marginals Ψ_i ,*

$$\Psi_i(\rho) = \rho,$$

unless Γ is a set of mutually orthogonal pure quantum states.

Remark. A perfect quantum cloning channel $\Psi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ has marginals Φ_i of the form $\Phi_i = \text{id}_d$ the identity quantum channels, and Choi matrices $C_{\Phi_i} = d \cdot \omega$ an unnormalized maximally entangled quantum state. The no-cloning Theorem 3.1 is a reformulation of the monogamy of the entanglement, and the

fact that there is no quantum channel compatible with the identity quantum channels.

A generalisation of the quantum cloning problem can be considered given a quantum cloning channel $\Phi : (\mathcal{M}_d)^{\otimes M} \rightarrow (\mathcal{M}_d)^{\otimes N}$, which maps an *input* pure quantum state as an identical product state on a M -fold composite quantum system, to an *output* mixed quantum state on a N -fold composite quantum system,

$$\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F\left(\Phi_i(\rho^{\otimes M}), \rho\right) \right].$$

Even with M identical copies as *input*, the no-cloning theorem hold.

Theorem 3.2 ($M \rightarrow N$ no-cloning theorem [Wer98]). *For any subset Γ of the pure quantum states, there is no quantum cloning channel $\Phi : (\mathcal{M}_d)^{\otimes M} \rightarrow (\mathcal{M}_d)^{\otimes N}$, with $M < N$, such that for all pure quantum states $\rho \in \Gamma$ and all marginals Φ_i ,*

$$\Phi_i(\rho^{\otimes M}) = \rho,$$

unless Γ is a set of mutually orthogonal pure quantum states.

It is important to note that the quantum cloning problem, the $1 \rightarrow N$ no-cloning Theorem 3.1 and $M \rightarrow N$ no-cloning Theorem 3.2 are states on pure states only. When mixed states are considered, the problem becomes the **quantum broadcasting problem**, and the equivalent no-broadcasting theorems do not hold in the same generality. In particular for mixed enough states, it is possible to broadcast.

Theorem (Superbroadcasting [DMP05]). *For $0 < M < N$ large enough, there exists a quantum channel $\Phi : (\mathcal{M}_2)^{\otimes M} \rightarrow (\mathcal{M}_2)^{\otimes N}$ such that for all 2-dimensional mixed states $\rho \in \mathcal{D}_2$, where $\rho = \frac{1}{2}(I_2 + r_x \cdot \sigma_x + r_y \cdot \sigma_y + r_z \cdot \sigma_z)$ and with $\|r\|$ small enough, the mixed state,*

$$\sigma := \Phi_i(\rho^{\otimes M}),$$

commutes with ρ , for all marginals Φ_i . I.e. they are collinear in the spherical representation:

$$\mathcal{D}_2 = \left\{ \frac{1}{2}(I_2 + r_x \cdot \sigma_x + r_y \cdot \sigma_y + r_z \cdot \sigma_z) \mid r := (r_x, r_y, r_z) \in \mathbb{R}^3 \text{ and } \|r\| \leq 1 \right\}.$$

Moreover, if $\sigma = \frac{1}{2}(I_2 + r'_x \cdot \sigma_x + r'_y \cdot \sigma_y + r'_z \cdot \sigma_z)$, then $\|r'\| \geq \|r\|$.

From [Proposition 2.2](#) and the joint concavity quantum fidelity, given two quantum cloning channels Φ and Ψ from \mathcal{M}_d to $(\mathcal{M}_d)^{\otimes N}$, then for all subset Γ of the pure quantum states and for all $1 \leq i \leq N$,

$$\mathbb{E}_{\rho \in \Gamma} \left[F \left(\frac{(\Phi + \Psi)_i(\rho)}{2}, \rho \right) \right] \geq \frac{\mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)] + \mathbb{E}_{\rho \in \Gamma} [F(\Psi_i(\rho), \rho)]}{2}.$$

Hence, in order to address the quantum cloning optimization problem, the approach would be to identify the largest uniform sum of quantum cloning channels.

Let Γ be a subset of the pure quantum states, and G be a compact subgroup of the unitary group U_d acting on \mathbb{C}^d , such that for all $\rho \in \Gamma$ and for all $M \in G$,

$$M\rho M^* \in \Gamma.$$

The **twirling** of a quantum channel $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$, with respect to Γ and G , denoted $\tilde{\Phi}$, is defined for all $\rho \in \Gamma$ by

$$\tilde{\Phi}(\rho) := \int_G (M^*)^{\otimes N} \left(\Phi(M\rho M^*) \right) M^{\otimes N} dM,$$

where the integral is taken with respect to the normalized Haar measure on the group G . Then for all $1 \leq i \leq N$,

$$\mathbb{E}_{\rho \in \Gamma} [F(\tilde{\Phi}_i(\rho), \rho)] \geq \mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)].$$

The approach would be to consider the largest group G . In particular the partially transposed Choi matrix $C_{\tilde{\Phi}}^\Gamma$ is in the commutant of G .

Proposition 3.3. *Let Γ be a subset of the pure quantum states, let G be a compact subgroup of U_d , and $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ a quantum cloning channel. If $\tilde{\Phi}$ is the twirling of Φ with respect to Γ and G , then for all $M \in G$,*

$$[C_{\tilde{\Phi}}^\Gamma, M^{\otimes(N+1)}] = 0.$$

Proof. Since G is a subgroup of U_d , for any M in G , the following two equalities hold,

$$(\bar{M} \otimes M) |\Omega\rangle = |\Omega\rangle \quad \text{and} \quad \langle \Omega | (M^\top \otimes M^*) = \langle \Omega |.$$

Then for any $M \in G$,

$$\begin{aligned} C_{\tilde{\Phi}}^{\Gamma} &= \left(\left(\text{id}_d \otimes \tilde{\Phi} \right) (d \cdot |\Omega \rangle \langle \Omega|) \right)^{\Gamma} \\ &= \left(\left(\text{id}_d \otimes \tilde{\Phi} \right) \left(d(\bar{M} \otimes M) |\Omega \rangle \langle \Omega| (M^{\top} \otimes M^*) \right) \right)^{\Gamma}. \end{aligned}$$

From the definition of the twirling $\tilde{\Phi}$, then for all $M \in G$ and for all $\rho \in \Gamma$

$$\tilde{\Phi}(M \rho M^*) = M^{\otimes N} (\tilde{\Phi}(\rho)) (M^*)^{\otimes N}.$$

Then the following commutation relation on the partially transposed Choi matrix $C_{\tilde{\Phi}}^{\Gamma}$ holds for any $M \in G$,

$$\begin{aligned} C_{\tilde{\Phi}}^{\Gamma} &= \left((\bar{M} \otimes M^{\otimes N}) \left(\left(\text{id}_d \otimes \tilde{\Phi} \right) (d \cdot |\Omega \rangle \langle \Omega|) \right) (M^{\top} \otimes (M^*)^{\otimes N}) \right)^{\Gamma} \\ &= (M \otimes M^{\otimes N}) \left(\left(\text{id}_d \otimes \tilde{\Phi} \right) (d \cdot |\Omega \rangle \langle \Omega|) \right)^{\Gamma} (M^* \otimes (M^*)^{\otimes N}) \\ &= (M \otimes M^{\otimes N}) C_{\tilde{\Phi}}^{\Gamma} (M^* \otimes (M^*)^{\otimes N}). \end{aligned}$$

That is $[C_{\tilde{\Phi}}^{\Gamma}, M^{\otimes(N+1)}] = 0$.

Corollary 3.4. *Let Γ be a subset of the pure quantum states, let G be a compact subgroup of U_d , and $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ a quantum cloning channel. If $\tilde{\Phi}$ is the twirling of Φ with respect to Γ and G , then for all $M \in G$ and all $i \in \{1, \dots, N\}$,*

$$[C_{\tilde{\Phi}_i}^{\Gamma}, M^{\otimes 2}] = 0.$$

Remark. The importance of twirling a quantum channel is twofold: first, it allows to improve the performance of the quantum channel with respect to the optimization problem, and second, it allows to simplify the optimization problem through the induced symmetries. The study of optimization problems under symmetries has been the subject of substantial work [FST22; GO22].

3.1.1 Universal quantum cloning problem

When the subset Γ of the pure quantum states is the full set of pure quantum states, it can choose the subgroup G of U_d to be the full unitary group. The quantum cloning problem becomes the **universal quantum cloning problem**. Then from [Proposition 3.3](#), the partially transposed Choi matrix $C_{\tilde{\Phi}}^\Gamma$ of a twirled quantum cloning channel $\tilde{\Phi}$ commutes with all the unitary matrices $U^{\otimes(N+1)}$, i.e. it is in the commutant of the algebra

$$\text{Span}_{\mathbb{C}} \{U^{\otimes(N+1)} \mid U \in U_d\}.$$

From [Theorem 1.3.2](#) and [Theorem 1.3](#), the partially transposed Choi matrix $C_{\tilde{\Phi}}^\Gamma$ is in the algebra

$$\text{Span}_{\mathbb{C}} \{\psi(\sigma) \mid \sigma \in \mathfrak{S}_{N+1}\}.$$

That is, it exists a family of complex numbers c_σ indexed by the permutations of \mathfrak{S}_{N+1} such that

$$C_{\tilde{\Phi}} = \sum_{\sigma \in \mathfrak{S}_{N+1}} c_\sigma \cdot \psi(\sigma)^\Gamma.$$

Therefore, the Choi matrix $C_{\tilde{\Phi}}$ is a sum of partially transposed tensor representation of the symmetric group \mathfrak{S}_{N+1} . Hence, a sum of partially transposed tensor representation of the symmetric group \mathfrak{S}_{N+1} is a Choi matrix of a quantum channel if both the positivity and the trace conditions of [Theorem 2.1](#) hold. These conditions depend on the $(N+1)!$ coefficients, and in particular it does not depend on the dimension of the quantum system. Recall from the Stirling's formula that the asymptotic growth of the factorial function is,

$$n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

From [Proposition 3.3](#), the partially transposed Choi matrix $C_{\tilde{\Phi}_i}$ of each marginal $\tilde{\Phi}_i$ has the form

$$\begin{aligned} C_{\tilde{\Phi}_i} &= \alpha_i \cdot \psi((1)(2))^\Gamma + \beta_i \cdot \psi((1\ 2))^\Gamma \\ &= \alpha_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \text{---} & \text{---} \\ \bullet & \bullet \end{array} \right)^\Gamma + \beta_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \diagdown & \diagup \\ \bullet & \bullet \end{array} \right)^\Gamma \\ &= \alpha_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \text{---} & \text{---} \\ \bullet & \bullet \end{array} \right) + \beta_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \text{---} & \text{---} \\ \bullet & \bullet \end{array} \right), \end{aligned}$$

for some complex numbers α_i and β_i . For all $\rho \in \Gamma$, the marginals $\tilde{\Phi}_i$ on ρ are

$$\begin{aligned}\tilde{\Phi}_i(\rho) &= \text{Tr}_{\{0\}} [C_{\tilde{\Phi}_i}(\rho^\top \otimes I_d)] \\ &= \alpha_i \cdot \text{Tr}_{\{0\}} [\psi((1)(2))^\Gamma(\rho \otimes I_d)] + \beta_i \cdot \text{Tr}_{\{0\}} [\psi((1\ 2))^\Gamma(\rho \otimes I_d)] \\ &= \alpha_i \cdot I_d + \beta_i \cdot \rho,\end{aligned}$$

and the fidelity $F(\tilde{\Phi}_i(\rho), \rho)$ becomes,

$$\begin{aligned}F(\tilde{\Phi}_i(\rho), \rho) &= \text{Tr} [(\tilde{\Phi}_i(\rho))\rho] \\ &= \text{Tr} [C_{\tilde{\Phi}_i}((\rho^\top \cdot \rho) \otimes I_d)] \\ &= \text{Tr} [C_{\tilde{\Phi}_i}(\rho \otimes I_d)] \\ &= \alpha_i \cdot \text{Tr} [\psi((1)(2))^\Gamma(\rho \otimes I_d)] + \beta_i \cdot \text{Tr} [\psi((1\ 2))^\Gamma(\rho \otimes I_d)] \\ &= d \cdot \alpha_i + \beta_i.\end{aligned}$$

3.1.2 Equatorial quantum cloning problem

Recall that in the case $\mathcal{H} := \mathbb{C}^2$, the pure quantum states can be written,

$$\frac{1}{2}(I_2 + r_x \cdot \sigma_x + r_y \cdot \sigma_y + r_z \cdot \sigma_z),$$

with $r := (r_x, r_y, r_z) \in \mathbb{R}^3$ and $\|r\| = 1$, and thus are isomorphic to the unit sphere in \mathbb{R}^3 . The **equatorial pure quantum states** are the pure quantum states located on the $x - y$ equator of this sphere, i.e. $r_z = 0$. The equatorial pure quantum states are of the form,

$$\frac{e^{i\theta_0} |0\rangle + e^{i\theta_1} |1\rangle}{\sqrt{2}},$$

for some $\theta \in [0, 2\pi)^2$.

Remark. The 4 states used in the BB84 protocol [BB84] are all in the $x - z$ equator, and the two equators $x - y$ and $x - z$ are connected by a change of basis.

When the subset Γ of the pure quantum states is the set of states of the form,

$$\frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} e^{i\theta_k} |k\rangle,$$

for some $\theta \in [0, 2\pi)^d$, the subgroup G of U_d can be chosen to be the group of monomial matrices in $[0, 2\pi)$. The quantum cloning problem becomes the **equatorial quantum cloning problem**. Then from [Proposition 3.3](#), the partially transposed Choi matrix $C_{\tilde{\Phi}}^\Gamma$ of a twirled quantum cloning channel $\tilde{\Phi}$ commutes with all the diagonal unitary matrices $U^{\otimes(N+1)}$ and all the tensor representation of permutations $\psi(\sigma)^{\otimes(N+1)}$, i.e. it is in the commutant of the algebra

$$\text{Span}_{\mathbb{C}} \{ U^{\otimes n} \cdot \phi(\sigma)^{\otimes n} \mid U \in \text{diag. } U_d \text{ and } \sigma \in \mathfrak{S}_d \}.$$

From [Theorem 1.4](#), the partially transposed Choi matrix $C_{\tilde{\Phi}}^\Gamma$ is in the algebra

$$\text{Span}_{\mathbb{C}} \{ \psi(p) \mid p \in \mathbb{U}_n \}.$$

That is, it exists a family of complex numbers c_p indexed by the uniform block permutations of \mathbb{U}_{N+1} such that

$$C_{\tilde{\Phi}} = \sum_{p \in \mathbb{U}_{N+1}} c_p \cdot \psi(p)^\Gamma.$$

From [Proposition 3.3](#), the partially transposed Choi matrix $C_{\tilde{\Phi}_i}$ of each marginal $\tilde{\Phi}_i$ has the form

$$\begin{aligned} C_{\tilde{\Phi}_i} &= \alpha_i \cdot \psi(13 | 24)^\Gamma + \beta_i \cdot \psi(14 | 23)^\Gamma + \gamma_i \cdot \psi(1234)^\Gamma \\ &= \alpha_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \right)^\Gamma + \beta_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \right)^\Gamma + \gamma_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \begin{array}{c} \text{H} \\ \text{H} \end{array} \right)^\Gamma \\ &= \alpha_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \right) + \beta_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \begin{array}{c} \text{C} \quad \text{C} \end{array} \right) + \gamma_i \cdot \psi \left(\begin{array}{cc} \bullet & \bullet \\ \bullet & \bullet \end{array} \begin{array}{c} \text{H} \\ \text{H} \end{array} \right), \end{aligned}$$

for some complex numbers α_i, β_i and γ_i . For all $\rho \in \Gamma$, the marginals $\tilde{\Phi}_i$ on ρ are

$$\begin{aligned} \tilde{\Phi}_i(\rho) &= \text{Tr}_{\{0\}} [C_{\tilde{\Phi}_i}(\rho^\Gamma \otimes I_d)] \\ &= \text{Tr}_{\{0\}} \left[\left(\alpha_i \cdot \psi(13 | 24)^\Gamma + \beta_i \cdot \psi(14 | 23)^\Gamma + \gamma_i \cdot \psi(1234)^\Gamma \right) (\rho \otimes I_d) \right] \\ &= \alpha_i \cdot I_d + \beta_i \cdot \rho + \gamma_i \cdot \text{diag}(\rho_{11}, \dots, \rho_{dd}) \\ &= \alpha_i \cdot I_d + \beta_i \cdot \rho + \gamma_i \cdot I_d, \end{aligned}$$

and the fidelity $F(\tilde{\Phi}_i(\rho), \rho)$ becomes,

$$\begin{aligned}
F(\tilde{\Phi}_i(\rho), \rho) &= \text{Tr} \left[(\tilde{\Phi}_i(\rho)) \rho \right] \\
&= \text{Tr} \left[C_{\tilde{\Phi}_i} \left((\rho^\top \cdot \rho) \otimes I_d \right) \right] \\
&= \text{Tr} \left[C_{\tilde{\Phi}_i} (\rho \otimes I_d) \right] \\
&= \text{Tr} \left[\left(\alpha_i \cdot \psi(13 | 24)^\Gamma + \beta_i \cdot \psi(14 | 23)^\Gamma + \gamma_i \cdot \psi(1234)^\Gamma \right) (\rho \otimes I_d) \right] \\
&= d \cdot \alpha_i + \beta_i + \gamma_i.
\end{aligned}$$

3.1.3 Average vs. worst fidelity

In [Section 3.1.1](#) and [Section 3.1.2](#), the marginals $\tilde{\Phi}_i$ and $\tilde{\Psi}_i$ of a twirled quantum cloning channel $\tilde{\Phi}$ for the universal quantum cloning problem and a twirled quantum cloning channel $\tilde{\Psi}$ for the equatorial quantum cloning problem, were shown to be, on all $\rho \in \Gamma$, some linear combinations of the trace, identity maps. Using the trace preserving property of quantum channels, the marginals becomes on all $\rho \in \Gamma$, some affine combinations

$$\tilde{\Phi}_i(\rho) = p \cdot \rho + (1-p) \frac{I_d}{d} \quad \text{and} \quad \tilde{\Psi}_i(\rho) = q \cdot \rho + (1-q) \frac{I_d}{d},$$

for some $p, q \in \mathbb{R}$, and with fidelities $F(\tilde{\Phi}_i(\rho), \rho)$ and $F(\tilde{\Psi}_i(\rho), \rho)$,

$$F(\tilde{\Phi}_i(\rho), \rho) = p + \frac{1-p}{d} \quad \text{and} \quad F(\tilde{\Psi}_i(\rho), \rho) = q + \frac{1-q}{d},$$

In particular, none of the fidelities depend on the quantum state. Hence both optimization problems,

$$\begin{aligned}
(\text{average}) \quad & \sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F(\Phi_i(\rho), \rho) \right] \\
(\text{worst}) \quad & \sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \inf_{\rho \in \Gamma} \left[F(\Phi_i(\rho), \rho) \right],
\end{aligned}$$

are equal on universal and equatorial Γ 's. However, the expectation value will be more convenient to manipulate, especially for to determine an upper bound.

3.2 Upper bound

This section is dedicated to the determination of an upper bound for both the universal and equatorial quantum cloning problems, given a direction vector $a \in [0, 1]^N$. For all quantum cloning channels $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$, and for all subset Γ of the pure quantum states,

$$\begin{aligned} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)] &= \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[\text{Tr} \left[(\Phi_i(\rho)) \rho \right] \right] \\ &= \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[\text{Tr} \left[(\Phi(\rho)) (\rho_{(i)} \otimes I_d^{\otimes(N-1)}) \right] \right] \\ &= \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[\text{Tr} \left[C_\Phi(\rho_{(0)}^\top \otimes \rho_{(i)} \otimes I_d^{\otimes(N-1)}) \right] \right] \\ &= \sum_{i=1}^N a_i \cdot \text{Tr} \left[C_\Phi \left(\mathbb{E}_{\rho \in \Gamma} [\rho_{(0)}^\top \otimes \rho_{(i)}] \otimes I_d^{\otimes(N-1)} \right) \right]. \end{aligned}$$

Theorem 3.5. *For any direction vector $a \in [0, 1]^N$ the universal quantum cloning problem is upper bounded by*

$$\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)] \leq \frac{\lambda_{\max}(R_a)}{d+1},$$

where $\lambda_{\max}(R_a)$ is the largest eigenvalue of the matrix

$$R_a := \sum_{i=1}^N a_i \cdot (d^2 \cdot I_{(0,i)} + d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)},$$

with quantum states $I_{(0,i)}$ and $\omega_{(0,i)}$, respectively maximally mixed and maximally entangled, between the 0-th and i -th quantum systems.

Proof. Let $\vee^N \mathbb{C}^d$ be the **symmetric subspace** of $(\mathbb{C}^d)^{\otimes N}$ defined by

$$\vee^N \mathbb{C}^d := \text{Span}_{\mathbb{C}} \{ v^{\otimes N} \mid v \in \mathbb{C}^d \}.$$

It is well known [Har13] that $\vee^N \mathbb{C}^d$ is an irreducible representation vector space for the representation $U \mapsto U^{\otimes N}$ of the unitary group U_d . Since the pure quan-

tum states Γ are generated by the unitary matrices, i.e. $\Gamma \simeq \{U\rho U^* \mid U \in \mathcal{U}_d\}$ for any pure quantum state ρ , then

$$\begin{aligned} \mathbb{E}_{\rho \in \Gamma} [\rho_{(0)}^\top \otimes \rho_{(i)}] &= \left(\mathbb{E}_{\rho \in \Gamma} [\rho_{(0)} \otimes \rho_{(i)}] \right)^\Gamma \\ &= \left(\int_{\Gamma} \rho \otimes \rho \, d\rho \right)^\Gamma \\ &= \left(\int_{\mathcal{U}_d} U |0\rangle\langle 0| U^* \otimes U |0\rangle\langle 0| U^* \, dU \right)^\Gamma. \end{aligned}$$

Note that the integral, before taking the partial transpose, commutes with all the unitary matrices $U \otimes U$ and lives in $\text{End}(\vee^2 \mathbb{C}^d)$, by Schur's Lemma it must be a multiple of the identity in $\vee^2 \mathbb{C}^d$. The identity of the bipartite symmetric subspace is

$$\frac{\psi((1)(2)) + \psi((1 \, 2))}{2}.$$

The unit trace condition, together with the partial transpose, give

$$\begin{aligned} \mathbb{E}_{\rho \in \Gamma} [\rho \otimes \rho] &= \frac{\psi((1)(2))^\Gamma + \psi((1 \, 2))^\Gamma}{d(d+1)} \\ &= \frac{d^2 \cdot \mathbf{I}_{(0,1)} + d \cdot \omega_{(0,1)}}{d(d+1)}. \end{aligned}$$

Finally the universal quantum cloning problem becomes

$$\begin{aligned} &\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)] \\ &= \sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \text{Tr} \left[C_\Phi \left(\mathbb{E}_{\rho \in \Gamma} [\rho_{(0)}^\top \otimes \rho_{(i)}] \otimes I_d^{\otimes (N-1)} \right) \right] \\ &= \sup_{\Phi \text{ CPTP}} \frac{\text{Tr} [C_\Phi R_a]}{d(d+1)}. \end{aligned}$$

Then from the inequality $\text{Tr}[CR] \leq \text{Tr}[C] \cdot \lambda_{\max}(R)$ that holds for any positive semidefinite matrix C and symmetric matrix R , and the equality $\text{Tr} [C_\Phi] = d$

for any Choi matrix C_Φ of a quantum channel $\Phi : \mathcal{M}_d \rightarrow \mathcal{M}_{d'}$,

$$\begin{aligned} \sup_{\Phi \text{ CPTP}} \frac{\text{Tr}[C_\Phi R_a]}{d(d+1)} &\leq \sup_{\Phi \text{ CPTP}} \frac{\text{Tr}[C_\Phi]}{d(d+1)} \lambda_{\max}(R_a) \\ &= \frac{\lambda_{\max}(R_a)}{d+1}. \end{aligned}$$

Remark. The upper bound in [Theorem 3.5](#) is a special case of the result of Jaromír Fiurášek on the extremal equation for optimal completely-positive maps [[Fiu01](#)].

The spectrum of the matrix R_a has been considered in a recent series of papers for the port-based teleportation protocol [[Stu+17](#); [Moz+18](#); [Led22](#)]. In particular, in [[Chr+21](#), Lemma 3.6], all the eigenvalues of the operator R_a , up to shift factor, are given in the special case of $a = \frac{1}{N}(1, \dots, 1)$.

Theorem 3.6. *For any direction vector $a \in [0, 1]^N$ the equatorial quantum cloning problem is upper bounded by*

$$\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)] \leq \frac{\lambda_{\max}(R_a)}{d},$$

where $\lambda_{\max}(R_a)$ is the largest eigenvalue of the matrix

$$R_a := \sum_{i=1}^N a_i \cdot (d^2 \cdot I_{(0,i)} + d \cdot \omega_{(0,i)} - d \cdot X_{(0,i)}) \otimes I^{\otimes(N-1)},$$

with quantum states $I_{(0,i)}$ and $\omega_{(0,i)}$, respectively maximally mixed and maximally entangled, between between the 0-th and i -th quantum systems, and quantum state $X_{(0,i)}$ defined by

$$X := \frac{1}{d} \sum_{i=0}^{d-1} |ii\rangle\langle ii|,$$

between between the 0-th and i -th quantum systems.

Proof. Let $|+\rangle$ be the pure quantum state defined by $|+\rangle := \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |i\rangle$, then for any pure quantum state $|\psi\rangle \in \Gamma$, i.e., of the form $\frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} e^{i\theta_k} |k\rangle$, there

exists a diagonal unitary matrix U in $\text{diag. } U_d$ such that $|\psi\rangle = U|+\rangle$. Then

$$\begin{aligned} \mathbb{E}_{\rho \in \Gamma} [\rho_{(0)}^\top \otimes \rho_{(i)}] &= \int_{\Gamma} \rho^\top \otimes \rho \, d\rho \\ &= \int_{\text{diag. } U_d} \bar{U} |+\rangle\langle+| U^\top \otimes U |+\rangle\langle+| U^* \, dU. \end{aligned}$$

Using integration over random diagonal unitary matrices [NS21],

$$\int_{\text{diag. } U_d} \bar{U} |+\rangle\langle+| U^\top \otimes U |+\rangle\langle+| U^* \, dU = \frac{d^2 \cdot I_{(0,1)} + d \cdot \omega_{(0,1)} - d \cdot X_{(0,1)}}{d^2}.$$

Which leads to the upper bound,

$$\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} [F(\Phi_i(\rho), \rho)] \leq \frac{\lambda_{\max}(R_a)}{d}.$$

Remark. For both the upper bounds in Theorem 3.5 and Theorem 3.6, the identity terms in the matrices R_a only lead to a shift in the largest eigenvalues $\lambda_{\max}(R_a)$.

3.3 Universal $1 \rightarrow 2$ quantum cloning problem

This section is devoted to the universal $1 \rightarrow 2$ quantum cloning problem, in the general case of quantum systems of arbitrary dimension [NPR21]. The problem is studied from the following perspective: given some target pair of fidelities (f_1, f_2) , does there exist a quantum cloning channel Φ from \mathcal{M}_d to $(\mathcal{M}_d)^{\otimes 2}$, such that $F(\Phi_i(\rho), \rho) = f_i$ for all pure quantum state ρ . As seen in Section 3.1.3, by twirling the quantum cloning channel Φ , the marginals becomes for all pure quantum states ρ ,

$$\tilde{\Phi}_1(\rho) = p_1 \cdot \rho + (1 - p_1) \frac{I_d}{d} \quad \text{and} \quad \tilde{\Phi}_2(\rho) = p_2 \cdot \rho + (1 - p_2) \frac{I_d}{d},$$

for some $p_1, p_2 \in \mathbb{R}$. Therefore, the following transformations rules for $\tilde{\Phi}$ hold:

$$f_1 = p_1 + \frac{(1 - p_1)}{d} \quad \quad \quad f_2 = p_2 + \frac{(1 - p_2)}{d}$$

$$p_1 = \frac{df_1 - 1}{d - 1} \qquad p_2 = \frac{df_2 - 1}{d - 1}$$

The main result is expressed as the following: the **achievable fidelity region**, defined as

$$\left\{ (f_1, f_2) \mid \exists \Phi : \mathcal{M}_d \xrightarrow{\text{CPTP}} (\mathcal{M}_d)^{\otimes 2} \text{ such that } \mathbb{E}_{\rho \in \mathcal{U}_d} [F(\Phi_i(\rho), \rho)] = f_i \right\},$$

is a **union of ellipses**, with the optimal one coming from a restricted class of quantum cloning channel (see [Figure 1](#)).

Theorem ([NPR21]). *The achievable fidelity region for the universal $1 \rightarrow 2$ quantum cloning problem is the union of a family of ellipses indexed by $\lambda \in (0, d]$, given by*

$$\frac{x^2}{a_\lambda^2} + \frac{(y - c_\lambda)^2}{b_\lambda^2} \leq 1,$$

with $a_\lambda := \frac{\lambda}{\sqrt{d^2-1}}$, $b_\lambda := \frac{\lambda}{d^2-1}$ and $c_\lambda := \frac{\lambda d - 2}{d^2-1}$. The parameters x and y can be expressed as,

$$\begin{cases} x &= \frac{d(f_1 - f_2)}{d-1} \\ y &= \frac{d(f_1 + f_2) - 2}{d-1} \end{cases} \quad \text{or} \quad \begin{cases} x &= p_1 - p_2 \\ y &= p_1 + p_2. \end{cases}$$

The optimal quantum cloning channels correspond to $\lambda = d$.

In [Section 3.1.1](#), the Choi matrix $C_{\tilde{\Phi}}$ of a twirled quantum cloning channel $\tilde{\Phi}$ from \mathcal{M}_d to $(\mathcal{M}_d)^{\otimes 2}$, was shown to be a linear combination of the 6 partially transposed tensor representations of the symmetric group \mathfrak{S}_3 :

$$C_{\tilde{\Phi}} = c_1 \cdot \psi \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right) + c_2 \cdot \psi \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right) + c_3 \cdot \psi \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right) + c_4 \cdot \psi \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right) + c_5 \cdot \psi \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right) + c_6 \cdot \psi \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right),$$

for complex numbers $c_1, \dots, c_6 \in \mathbb{C}$. Such linear combination is the Choi matrix of a quantum channel if both the positivity and the trace conditions of [Theorem 2.1](#) are satisfied.

By taking the corresponding partial traces, the two marginals of $\tilde{\Phi}$ becomes on all pure quantum states ρ ,

$$\begin{aligned} \tilde{\Phi}_1(\rho) &= \underbrace{(dc_2 + c_5 + c_6)}_{p_1} \rho + \underbrace{(d^2c_1 + dc_3 + dc_4)}_{1-p_1} \frac{I_d}{d} \\ \tilde{\Phi}_2(\rho) &= \underbrace{(dc_3 + c_5 + c_6)}_{p_2} \rho + \underbrace{(d^2c_1 + dc_2 + dc_4)}_{1-p_2} \frac{I_d}{d}. \end{aligned}$$




Figure 1 – The achievable fidelity region \square of the universal $1 \rightarrow 2$ quantum cloning problem is the union of a continuous family of ellipses \square .

The partially transposed tensor representations $\psi(\text{diagram 1})$ and $\psi(\text{diagram 2})$ corresponding to the two cycles $(1\ 2\ 3)$ and $(3\ 2\ 1)$ of \mathfrak{S}_3 contribute, with coefficients c_5 and c_6 , to both the p_1 and p_2 of $\tilde{\Phi}_1$ and $\tilde{\Phi}_2$. However, due to the no-cloning [Theorem 3.1](#), a Choi matrix composed only of the two cycles $(1\ 2\ 3)$ and $(3\ 2\ 1)$ of \mathfrak{S}_3 cannot be the Choi matrix of a quantum channel, since no linear combination,

$$\alpha \cdot \psi(\text{diagram 1}) + \beta \cdot \psi(\text{diagram 2})$$

is positive semidefinite. Hence, contributions from the other 4 partially transposed tensor representations are needed to ensure the positivity of the Choi matrix $C_{\tilde{\Phi}}$. The trace condition implies that

$$d^2 c_1 + d(c_2 + c_3 + c_4) + c_5 + c_6 = 1.$$

In the following, the tensor representation function ψ is dropped, and instead the partially transposed permutations are depicted using the graphical calculus from [Section 2.5](#). The focus now shifts to the problem of characterizing the positivity of the Choi matrix $C_{\tilde{T}}$ using the coefficients c_i . To achieve this, let the vectors u_i and

v_i of be defined for all $i \in \{0, \dots, d-1\}$ by,

$$u_i := \sqrt{d} \cdot |\Omega\rangle_{(0,1)} \otimes |i\rangle \quad \text{and} \quad v_i := \sqrt{d} \cdot |\Omega\rangle_{(0,2)} \otimes |i\rangle,$$

depicted as

$$u_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i \quad \text{and} \quad v_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i.$$

Then, the action of the partially transposed permutations on these vectors is given by,

$$\begin{array}{ccc} \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = d \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i \\ \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = d \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i, \end{array} \quad (3.1)$$

and,

$$\begin{array}{ccc} \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = d \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i \\ \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i & \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i = d \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}_i. \end{array} \quad (3.2)$$

The vectors $u_i + v_i$ and $u_i - v_i$ are eigenvectors of $\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}$ and $\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}$. Since they are all d -rank matrices, their complete (nonzero) spectrums is known:

$$\begin{aligned} \text{Spec} \left(\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) &= \begin{cases} (d+1) & \times d \\ (d-1) & \times d \end{cases} \\ \text{Spec} \left(\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) &= \begin{cases} +(d+1) & \times d \\ -(d-1) & \times d. \end{cases} \end{aligned}$$

The other two partially transposed permutations $\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}$ and $\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}$, are unitary matrices and thus have full rank.

In the next two Sections the positivity of the Choid matrix $C_{\mathfrak{F}}$ is characterize in terms of the coefficients c_i , first by restricting to the first 4, and then considering the general case.

3.3.1 Restricted quantum cloning channels

In this Section, the universal $1 \rightarrow 2$ quantum cloning problem will be solved when the Choi matrix $C_{\mathfrak{F}}$ is a linear combination of only 4 partially transposed permutations of \mathfrak{S}_3 , that is,

$$C_{\mathfrak{F}} = c_1 \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + c_2 \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + c_3 \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} + c_4 \cdot \begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array}.$$

Since $C_{\tilde{\tau}}$ must be positive semidefinite and in particular Hermitian, the following must hold: $c_1, c_2 \in \mathbb{R}$ and $c_3 = \bar{c}_4$. That is;

$$C_{\tilde{\Phi}} = \alpha \cdot \begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} + \beta \cdot \begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} + \gamma \cdot \begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} + \bar{\gamma} \cdot \begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix},$$

such that the trace condition becomes $d(\alpha + \beta) + 2\Re(\gamma) = 1$. Using Eq. (3.1) and Eq. (3.2), The 4 partially transposed permutations can be block diagonalized in the basis of the $2d$ vectors u_i and v_i , i.e.,

$$\left(\begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} \right)_i = \begin{pmatrix} u_i & v_i \\ d & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} u_i \\ v_i \end{pmatrix}$$

and,

$$\left(\begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} \right)_i = \begin{pmatrix} 0 & 0 \\ 1 & d \end{pmatrix} \quad \left(\begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} \right)_i = \begin{pmatrix} 0 & 0 \\ d & 1 \end{pmatrix} \quad \left(\begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} \right)_i = \begin{pmatrix} 1 & d \\ 0 & 0 \end{pmatrix}$$

such that each partially transposed permutation operators is a direct sum of d such blocks and $d^3 - 2d$ zero blocks, e.g.,

$$\begin{smallmatrix} \text{---} & \text{---} \\ \text{---} & \text{---} \end{smallmatrix} = \begin{pmatrix} d & 1 \\ 0 & 0 \end{pmatrix}^{\oplus d}.$$

Then the block diagonal decomposition of $C_{\tilde{\Phi}}$ becomes

$$(C_{\tilde{\Phi}})_i = \begin{pmatrix} d\alpha + \bar{\gamma} & \alpha + d\bar{\gamma} \\ \beta + d\gamma & d\beta + \gamma \end{pmatrix}.$$

It is well known that a 2×2 hermitian matrix M is positive semidefinite if and only if $\det(M) \geq 0$ and $\text{Tr}[M] \geq 0$. Then $C_{\tilde{\Phi}}$ is positive semi-definite if and only if each of its 2×2 blocks, in its block diagonal decomposition, are positive semidefinite. That is

$$\text{Tr} \left[(C_{\tilde{\Phi}})_i \right] = d(\alpha + \beta) + 2\Re(\gamma) \geq 0$$

and

$$\det \left((C_{\tilde{\Phi}})_i \right) = \alpha\beta - |\gamma|^2 \geq 0.$$

The first condition is always true since $d(\alpha + \beta) + 2\Re(\gamma) = 1$. Finally, the Choi matrix $C_{\tilde{\Phi}}$ is the Choi matrix of a quantum channel, and thus a quantum cloning channel, when both

$$d(\alpha + \beta) + 2\Re(\gamma) = 1 \quad \text{and} \quad \alpha\beta \geq |\gamma|^2.$$

The two marginals of $\tilde{\Phi}$ becomes on all pure quantum states ρ ,

$$\begin{aligned}\tilde{\Phi}_1(\rho) &= \underbrace{(d\alpha + 2\Re(\gamma))}_{p_1} \rho + \underbrace{d\beta}_{1-p_1} \frac{I_d}{d} \\ \tilde{\Phi}_2(\rho) &= \underbrace{(d\beta + 2\Re(\gamma))}_{p_2} \rho + \underbrace{d\alpha}_{1-p_2} \frac{I_d}{d}.\end{aligned}$$

Theorem. *The achievable fidelity region for the restricted universal $1 \rightarrow 2$ quantum cloning problem is the ellipse given by:*

$$\frac{x^2}{a^2} + \frac{(y-c)^2}{b^2} \leq 1,$$

with $a := \frac{d}{\sqrt{d^2-1}}$, $b := \frac{d}{d^2-1}$ and $c := \frac{d^2-2}{d^2-1}$. The parameters x and y can be expressed as,

$$\begin{cases} x &= \frac{d(f_1-f_2)}{d-1} \\ y &= \frac{d(f_1+f_2)-2}{d-1} \end{cases} \quad \text{or} \quad \begin{cases} x &= p_1 - p_2 \\ y &= p_1 + p_2. \end{cases}$$

Equivalently, the achievable fidelity region for the restricted universal $1 \rightarrow 2$ quantum cloning problem is the set,

$$\left\{ (p_1, p_2) \in \left[\frac{-1}{d^2-1}, 1 \right] \mid \frac{(1-p_1)(1-p_2)}{d^2} \geq \left(\frac{p_1+p_2-1}{2} \right)^2 \right\}.$$

Proof. A pair $(p_1, p_2) \in \left[\frac{-1}{d^2-1}, 1 \right]$ is in the achievable fidelity region for the restricted universal $1 \rightarrow 2$ quantum cloning problem if and only if there exists coefficients $\alpha, \beta \in \mathbb{R}$ and $\gamma \in \mathbb{C}$ satisfying both,

$$d(\alpha + \beta) + 2\Re(\gamma) = 1 \quad \text{and} \quad \alpha\beta \geq |\gamma|^2,$$

with,

$$\begin{aligned} p_1 &= d\alpha + 2\Re(\gamma) & p_2 &= d\beta + 2\Re(\gamma) \\ &= 1 - d\beta & &= 1 - d\alpha. \end{aligned}$$

Such a complex number γ exists if and only if,

$$\left(\frac{1 - d(\alpha + \beta)}{2} \right)^2 \leq \alpha\beta.$$

Rewriting the this inequality in terms of p_1 and p_2 yields,

$$\frac{(1-p_1)(1-p_2)}{d^2} \geq \left(\frac{p_1 + p_2 - 1}{2} \right)^2.$$




Figure 2 – The achievable fidelity region for the restricted universal $1 \rightarrow 2$ quantum cloning problem is an ellipse.

3.3.2 General quantum cloning channels

In this Section, the general case of the universal $1 \rightarrow 2$ quantum cloning problem will be solved when the Choi matrix $C_{\tilde{\Phi}}$ is a linear combination of all the 6 partially transposed permutations of \mathfrak{S}_3 , that is,

$$C_{\tilde{\Phi}} = \alpha \cdot \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array} + \beta \cdot \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array} + \gamma \cdot \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array} + \bar{\gamma} \cdot \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array} + \varepsilon_1 \cdot \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array} + \varepsilon_2 \cdot \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array}.$$

The hermitian condition on $C_{\tilde{\Phi}}$ imposes that $\alpha, \beta, \varepsilon_1, \varepsilon_2 \in \mathbb{R}$, and the trace condition reads $d(\alpha + \beta) + 2\Re(\gamma) + d^2\varepsilon_1 + d\varepsilon_2 = 1$. As in the previous Section, the action of $C_{\tilde{\Phi}}$ decomposes on with respect to the the $2d$ vectors u_i and v_i .

Let \mathbf{V} be the complex span of the $2d$ vectors u_i and v_i :

$$\mathbf{V} := \text{Span}_{\mathbb{C}} \left\{ \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array}, \begin{array}{c} \text{---} \text{---} \\ \text{---} \end{array} \right\},$$

then $\mathbf{V} \subset \mathbb{C}^d \otimes \vee^2 \mathbb{C}^d$, and is invariant by the two partially transposed permutations $\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}$ and $\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}$ that have both full rank. On \mathbf{V}^\perp , the spectrum of $\varepsilon_1 \cdot \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} + \varepsilon_2 \cdot \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}$ is

$$\text{Spec}|_{\mathbf{V}^\perp} \left(\varepsilon_1 \cdot \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} + \varepsilon_2 \cdot \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix} \right) = \begin{cases} \varepsilon_1 + \varepsilon_2 & \times d \frac{d(d+1)}{2} - 2d \\ \varepsilon_1 - \varepsilon_2 & \times d \frac{d(d-1)}{2}. \end{cases}$$

Then the complete block diagonal decomposition of $C_{\tilde{\Phi}}$ is made of 2×2 and 1×1 blocks. The 1×1 blocks $(\varepsilon_1 + \varepsilon_2)$ and $(\varepsilon_1 - \varepsilon_2)$ are positive when $\varepsilon_1 \geq |\varepsilon_2|$. On \mathbf{V} the block diagonalization of $C_{\tilde{\Phi}}$ becomes,

$$(C_{\tilde{\Phi}})_i = \begin{pmatrix} d\alpha + \bar{\gamma} + \varepsilon_1 & \alpha + d\bar{\gamma} + \varepsilon_2 \\ \beta + d\gamma + \varepsilon_2 & d\beta + \gamma + \varepsilon_1 \end{pmatrix}.$$

After the change of basis

$$\begin{aligned} \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}_i &\mapsto \frac{1}{\sqrt{2(d+1)}} \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}_i + \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}_i \right) \\ \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}_i &\mapsto \frac{1}{\sqrt{2(d-1)}} \left(\begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}_i - \begin{smallmatrix} \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{smallmatrix}_i \right), \end{aligned}$$

the block diagonal decomposition becomes the Hermitian,

$$(C_{\tilde{\Phi}})_i = \begin{pmatrix} (d+1) \frac{\alpha+\beta+2\cdot\Re(\gamma)}{2} + \varepsilon_1 + \varepsilon_2 & \frac{\sqrt{d^2-1}}{2}(\alpha - \beta) \\ \frac{\sqrt{d^2-1}}{2}(\alpha - \beta) & (d+1) \frac{\alpha+\beta-2\cdot\Re(\gamma)}{2} + \varepsilon_1 - \varepsilon_2 \end{pmatrix}.$$

The two marginals of $\tilde{\Phi}$ becomes on all pure quantum states ρ ,

$$\begin{aligned} \tilde{\Phi}_1(\rho) &= \underbrace{(d\alpha + 2\Re(\gamma))}_{p_1} \rho + \underbrace{(d\beta + d^2\varepsilon_1 + d\varepsilon_2)}_{1-p_1} \frac{I_d}{d} \\ \tilde{\Phi}_2(\rho) &= \underbrace{(d\beta + 2\Re(\gamma))}_{p_2} \rho + \underbrace{(d\alpha + d^2\varepsilon_1 + d\varepsilon_2)}_{1-p_2} \frac{I_d}{d}. \end{aligned}$$

Theorem. *The achievable fidelity region for the universal $1 \rightarrow 2$ quantum cloning problem is the union of a family of ellipses indexed by $\lambda \in (0, d]$, given by*

$$\frac{x^2}{a_\lambda^2} + \frac{(y - c_\lambda)^2}{b_\lambda^2} \leq 1,$$

with $a_\lambda := \frac{\lambda}{\sqrt{d^2-1}}$, $b_\lambda := \frac{\lambda}{d^2-1}$ and $c_\lambda := \frac{\lambda d-2}{d^2-1}$. The parameters x and y can be expressed as,

$$\begin{cases} x &= \frac{d(f_1-f_2)}{d-1} \\ y &= \frac{d(f_1+f_2)-2}{d-1} \end{cases} \quad \text{or} \quad \begin{cases} x &= p_1 - p_2 \\ y &= p_1 + p_2. \end{cases}$$

Proof. Setting $x := p_1 - p_2$ and $y := p_1 + p_2$, together with the relation $d(\alpha + \beta) + 2\Re(\gamma) + d^2\varepsilon_1 + d\varepsilon_2 = 1$ yields,

$$(C_{\tilde{\Phi}})_i = \begin{pmatrix} \frac{(d^2-1)y+d(d-1)((d^2-2)\varepsilon_1+d\varepsilon_2-1)+2}{2d} & \frac{\sqrt{d^2-1}}{2d}x \\ \frac{\sqrt{d^2-1}}{2d}x & -\frac{(d^2-1)y+d(d+1)((d^2-2)\varepsilon_1+d\varepsilon_2-1)+2}{2d} \end{pmatrix}.$$

Let $\lambda := -d((d^2-2)\varepsilon_1 + d\varepsilon_2 - 1)$, then the block diagonal decomposition of $C_{\tilde{\Phi}}$ reduces to

$$(C_{\tilde{T}})_i = \frac{1}{2d} \begin{pmatrix} (d^2-1)y - (d-1)\lambda + 2 & \sqrt{d^2-1}x \\ \sqrt{d^2-1}x & -((d^2-1)y - (d+1)\lambda + 2) \end{pmatrix}.$$

In this way, the positivity of each of the 2×2 blocks becomes

$$0 \leq \lambda \leq d \quad \text{and} \quad \frac{x^2}{a_\lambda^2} + \frac{(y - c_\lambda)^2}{b_\lambda^2} \leq 1,$$

with $a_\lambda := \frac{\lambda}{\sqrt{d^2-1}}$, $b_\lambda := \frac{\lambda}{d^2-1}$ and $c_\lambda := \frac{\lambda d-2}{d^2-1}$.

3.4 Universal $1 \rightarrow N$ quantum cloning problem

This section covers the arbitrary universal $1 \rightarrow N$ quantum cloning problem, in the general case of quantum systems of any dimension [NPR22].

3.4.1 Partially transposed permutations

This section is devoted to the study of the partially transposed tensor representations $\psi(\sigma)^\Gamma$ of the symmetric group \mathfrak{S}_{N+1} , that appear in the Choi matrix of the

twirled quantum cloning channels. In this section, the symmetric group \mathfrak{S}_{N+1} is the group of permutations of the set $\{0, 1, \dots, N\}$, starting from 0.

Remark. In the Choi matrix of a twirled quantum cloning channel, the *input* tensor corresponds to the set $\{0\}$ of its partially transposed tensor representations, and the *output* tensors correspond to the set $\{1, \dots, N\}$.

Let $\sigma \in \mathfrak{S}_{N+1}$ such that 0 is a fixed point of σ , i.e. $\sigma(0) = 0$. Assume that $\psi(\sigma)^\Gamma$ appears in the Choi matrix of a twirled quantum cloning channel, then σ does not contribute to the performance of the quantum cloning channel. Indeed on all pure quantal states ρ ,

$$\text{Tr}_{\{0\}} \left[\psi(\sigma)^\Gamma (\rho^\Gamma \otimes I_d^{\otimes N}) \right] = \text{Tr} [\rho^\Gamma] \cdot \psi(\hat{\sigma}) = \psi(\hat{\sigma}),$$

where $\hat{\sigma}$ is the permutation of the symmetric group \mathfrak{S}_N on $\{1, \dots, N\}$, obtained from σ by dropping $\{0\}$. Each marginals in thus a scalar multiple of the identity.

Let $\Sigma_{a,b}$ be the subset of \mathfrak{S}_{N+1} be defined for all $1 \leq a, b \leq N$ by,

$$\Sigma_{a,b} := \{ \sigma \in \mathfrak{S}_{N+1} \mid \sigma(0) = a \text{ and } \sigma^{-1}(0) = b \}.$$

This gives a partition of

$$\{ \sigma \in \mathfrak{S}_{N+1} \mid \sigma(0) \neq 0 \} = \bigcup_{1 \leq a, b \leq N} \Sigma_{a,b},$$

where each set $\Sigma_{a,b}$ contains $(N-1)!$ permutations.

Note that for all $1 \leq a, b \leq N$ and for all $\sigma \in \Sigma_{a,b}$, there exists a unique $\hat{\sigma} \in \mathfrak{S}_{N-1}$ such that the partially transposed tensor representation $\psi(\sigma)^\Gamma$ decomposes into,

$$\psi(\sigma)^\Gamma = \psi((1 \ a)) \cdot (d \cdot \omega_{(0,1)} \otimes \psi(\hat{\sigma})) \cdot \psi((1 \ b)), \quad (3.3)$$

where $(d \cdot \omega_{(0,1)} \otimes \psi(\hat{\sigma}))$ is the partially transposed tensor representation of a permutation in $\Sigma_{1,1}$.

Lemma 3.7. *Let distinct $1 \leq a, b, c \leq N$, then*

$$\begin{aligned} \text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma)^\Gamma \right] &= \frac{1}{d} \sum_{\sigma \in \mathfrak{S}_{N-1}} \text{Tr} [\psi(\sigma)] \cdot I_d \\ \text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{c,c}} \psi(\sigma)^\Gamma \right] &= \sum_{\sigma \in \mathfrak{S}_{N-1}} \text{Tr} [\psi(\sigma)] \cdot I_d. \end{aligned}$$

Proof. For the second equation, using the decomposition of Eq. (3.3),

$$\begin{aligned}
& \text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{c,c}} \psi(\sigma)^\Gamma \right] \\
&= \text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\hat{\sigma} \in \mathfrak{S}_{N-1}} \psi((1\ c)) \cdot (d \cdot \omega_{(0,1)} \otimes \psi(\hat{\sigma})) \cdot \psi((1\ c)) \right] \\
&= \text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\hat{\sigma} \in \mathfrak{S}_{N-1}} \sum_{i,j=0}^{d-1} \psi((1\ c)) \cdot (|ii\rangle\langle jj| \otimes \psi(\hat{\sigma})) \cdot \psi((1\ c)) \right] \\
&= \text{Tr} \left[\sum_{\hat{\sigma} \in \mathfrak{S}_{N-1}} \sum_{i,j=0}^{d-1} \psi((0\ (c-1))) \cdot (|ii\rangle\langle jj| \otimes \psi(\hat{\sigma})) \cdot \psi((0\ (c-1))) \right] \cdot |i\rangle\langle j| \\
&= \sum_{\sigma \in \mathfrak{S}_{N-1}} \text{Tr} [\psi(\sigma)] \cdot I_d,
\end{aligned}$$

where $\psi((0\ (c-1)))$ is the tensor representations of the permutation $(0\ (c-1))$ on $\{0, \dots, (N-1)\}$. For the first equation, since the partial transpose is a linear operator,

$$\text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma)^\Gamma \right] = \left(\text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma) \right] \right)^\Gamma.$$

For any $\sigma \in \mathfrak{S}_{N+1}$, the partial trace of the tensor representation $\psi(\sigma)$ is a multiple of the identity, i.e.,

$$\text{Tr}_{[N+1] \setminus \{0\}} [\psi(\sigma)] = c \cdot I_d,$$

with $c = \frac{1}{d} \text{Tr} [\psi(\sigma)]$. Then

$$\left(\text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma) \right] \right)^\Gamma = \frac{1}{d} \text{Tr} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma) \right] \cdot I_d.$$

For a permutation $\sigma \in \mathfrak{S}_{N+1}$, let $\#\sigma$ denotes the number of disjoint cycles of σ . Then

$$\begin{aligned}
\text{Tr} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma) \right] &= \sum_{\sigma \in \Sigma_{a,b}} d^{\#\sigma} \\
&= \sum_{\sigma \in \Sigma_{a,a}} d^{\#[\sigma \circ (a\ b)]}.
\end{aligned}$$

Let distinct $a, b \in \{0, \dots, N\}$ and a permutation $\sigma \in \mathfrak{S}_{N+1}$ with its decomposition into disjoint cycles $\sigma = c_1 \circ \dots \circ c_k$. If there exists $i \in \{1, \dots, k\}$ such that both $a, b \in c_i$, then the permutation $c_i \circ (a \ b)$ can be decomposed into two disjoint cycles. Otherwise, if there exist distinct $i, j \in \{1, \dots, k\}$ such that $a \in c_i$ and $b \in c_j$, then the permutation $c_i \circ c_j \circ (a \ b)$ can be decomposed into only one disjoint cycle. Finally

$$\#[\sigma \circ (a \ b)] = \begin{cases} \#\sigma + 1 & \text{if } \exists i \in \{1, \dots, k\} \text{ s.t. } a, b \in c_i \\ \#\sigma - 1 & \text{otherwise} \end{cases}.$$

But when $\sigma \in \Sigma_{a,a}$ and since $b \neq a$, in the decomposition of σ into disjoint cycles, a is in the cycle $(0 \ a)$, and $\#[\sigma \circ (a \ b)] = \#\sigma - 1$. That is

$$\text{Tr} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma) \right] = \frac{1}{d} \text{Tr} \left[\sum_{\sigma \in \Sigma_{a,a}} \psi(\sigma) \right].$$

Using the first equation,

$$\text{Tr}_{[N+1] \setminus \{0\}} \left[\sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma)^\Gamma \right] = \frac{1}{d} \sum_{\sigma \in \mathfrak{S}_{N-1}} \text{Tr} [\psi(\sigma)] \cdot I_d.$$

The next lemma establishes the relationship between a Choi matrix of a partially transposed tensor representation $\psi(\sigma)^\Gamma$, and its corresponding quantum channel.

Lemma 3.8. *Let some $1 \leq a, b \leq N$ and $\sigma \in \Sigma_{a,b}$, then there exist $\mu, \nu \in \mathfrak{S}_N$ satisfying $\mu(0) = a - 1$ and $\nu^{-1}(0) = b - 1$, such that $\psi(\sigma)^\Gamma$ is the Choi matrix of the linear map,*

$$X \mapsto \psi(\mu) \cdot (X \otimes I_d^{\otimes(N-1)}) \cdot \psi(\nu).$$

Proof. Using the decomposition of [Eq. \(3.3\)](#),

$$\psi(\sigma)^\Gamma = \psi((1 \ a)) \cdot (d \cdot \omega_{(0,1)} \otimes \psi(\hat{\sigma})) \cdot \psi((1 \ b)),$$

for some unique permutation $\hat{\sigma} \in \mathfrak{S}_{N-1}$. Then,

$$\begin{aligned} & \text{Tr}_{\{0\}} \left[\psi(\sigma)^\Gamma (X^\top \otimes I_d^{\otimes N}) \right] \\ &= \text{Tr}_{\{0\}} \left[\psi((1 \ a)) \cdot (d \cdot \omega_{(0,1)} \otimes \psi(\hat{\sigma})) \cdot \psi((1 \ b)) (X^\top \otimes I_d^{\otimes N}) \right] \end{aligned}$$

$$\begin{aligned}
&= \text{Tr}_{\{0\}} \left[\sum_{i,j=0}^{d-1} \psi((1 \ a)) \cdot (|ii\rangle\langle jj| \otimes \psi(\hat{\sigma})) \cdot \psi((1 \ b)) (X^\top \otimes I_d^{\otimes N}) \right] \\
&= \sum_{i,j=0}^{d-1} \langle i| X |j\rangle \cdot \psi((0 \ (a-1))) \cdot (|ii\rangle\langle jj| \otimes \psi(\hat{\sigma})) \cdot \psi((0 \ (b-1))) \\
&= \psi((0 \ (a-1))) \cdot (X \otimes \psi(\hat{\sigma})) \cdot \psi((0 \ (b-1))).
\end{aligned}$$

Thus, by setting $\psi(\mu) := \psi((0 \ (a-1)))$ and $\psi(\nu) := (I_d \otimes \psi(\hat{\sigma})) \cdot \psi((0 \ (b-1)))$, the result holds.

Recall that the upper bound of the universal quantum cloning problem for a direction vector a , from [Section 3.2](#), is given as the largest eigenvalue of the the matrix R_a defined by,

$$R_a := \sum_{i=1}^N a_i \cdot (d^2 \cdot I_{(0,i)} + d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)}.$$

The eigenvectors of R_a are the same as those of the matrix S_a defined by,

$$S_a := \sum_{i=1}^N a_i \cdot (d \cdot \omega_{(0,i)}) \otimes I^{\otimes(N-1)}.$$

Lemma 3.9. *[NPR22] The normalized largest eigenvectors of S_a are of the form*

$$\chi = \sum_{i=1}^N b_i \cdot (\sqrt{d} \cdot |\Omega\rangle_{(0,i)}) \otimes |v\rangle,$$

for some vector $|v\rangle$ in the symmetric subspace $\vee^{(N-1)}\mathbb{C}^d$, and some positive real numbers b_i satisfy the equation,

$$(d-1) \sum_{i=1}^N b_i^2 + \left(\sum_{i=1}^N b_i \right)^2 = 1.$$

The largest eigenvalue becomes $\lambda_{\max} = \sum_{i=1}^N a_i \left((d-1)b_i + \sum_{j=1}^N b_j \right)^2$.

Remark. Note that the positive real numbers b_i depend on the direction vector a .

3.4.2 Optimal symmetric quantum cloning channels

The **symmetric** universal $1 \rightarrow N$ quantum cloning problem is a special case of the quantum cloning problem, where all the marginals Φ_i of the quantum cloning channel are asked to be equal on all pure quantum states ρ , i.e.,

$$\Phi_i(\rho) = p \cdot \rho + (1 - p) \frac{I_d}{d},$$

where p does not depend on the choice of the marginal Φ_i .

Theorem ([KW99]). *The optimal quantum cloning channel Φ_{opt} from \mathcal{M}_d to $(\mathcal{M}_d)^{\otimes N}$, for the symmetric universal $1 \rightarrow N$ quantum cloning problem, is defined on all pure quantum states ρ by*

$$\Phi_{opt}(\rho) := \frac{d}{\text{Tr}[P_{\mathfrak{S}_N}^+]} P_{\mathfrak{S}_N}^+(\rho \otimes I_d^{\otimes(N-1)}) P_{\mathfrak{S}_N}^+,$$

where $P_{\mathfrak{S}_N}^+$ is the orthogonal projector onto the symmetric subspace $\vee^N \mathbb{C}^d$, defined by

$$P_{\mathfrak{S}_N}^+ := \frac{1}{N!} \sum_{\sigma \in \mathfrak{S}_N} \psi(\sigma).$$

Then each marginal $(\Phi_{opt})_i$ is equal, on all pure quantum states ρ , to,

$$(\Phi_{opt})_i(\rho) = \underbrace{\frac{d+N}{N(d+1)}}_{p_{opt}} \rho + \underbrace{\left(1 - \frac{d+N}{N(d+1)}\right)}_{1-p_{opt}} \frac{I_d}{d}.$$

3.4.3 Optimal asymmetric quantum cloning channels

The **asymmetric** universal $1 \rightarrow N$ quantum cloning problem is the general case of the quantum cloning problem, where the marginals Φ_i of the quantum cloning channel can be arbitrary.

In Section 3.3, the main technical difficulty was the positivity condition of the Choi matrix of the quantum cloning channels. Let $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ be a linear map defined by,

$$\Phi(X) := A(X \otimes I_d^{\otimes(N-1)})A^*,$$

for some matrix $A \in \mathcal{M}_{d^N}$. Let $\Psi : \mathcal{M}_{d^N} \rightarrow \mathcal{M}_{d^N}$ be a linear map defined by $\Psi(X) := AXA^*$, then,

$$\Psi(x \otimes I_d^{\otimes(N-1)}) = \Phi(X).$$




Figure 3 – The optimal fidelity for the symmetric universal $1 \rightarrow N$ quantum cloning problem.

Through this conjugacy form Ψ and in particular Φ are both completely positive.

Theorem 3.10. *Given a direction vector a , let $\chi = \sum_{i=1}^N b_i \cdot (\sqrt{d} \cdot |\Omega\rangle_{(0,i)}) \otimes |v\rangle$ be normalized largest eigenvectors of S_a . The optimal quantum cloning channel Φ_{opt}^a from \mathcal{M}_d to $(\mathcal{M}_d)^{\otimes N}$, for the asymmetric universal $1 \rightarrow N$ quantum cloning problem in the direction a , is defined on all pure quantum states ρ by*

$$\Phi_{\text{opt}}^a(\rho) := \frac{dN(N+d-1)}{\text{Tr}[P_{\mathfrak{S}_N}^+]} P_{\mathfrak{S}_N}^a(\rho \otimes I_d^{\otimes(N-1)})(P_{\mathfrak{S}_N}^a)^*,$$

where,

$$P_{\mathfrak{S}_N}^a := \frac{1}{N!} \sum_{\sigma \in \mathfrak{S}_N} b_{\sigma(0)+1} \cdot \psi(\sigma).$$

Proof. The complete positivity is given by the conjugacy form of the quantum channel Φ_{opt}^a .

For the trace preserving property, the Choi matrix $C_{\Phi_{\text{opt}}^a}$ of Φ_{opt}^a will be first

determined. For all pure quantum states ρ ,

$$\begin{aligned}\Phi_{\text{opt}}^a(\rho) &= \frac{dN(N+d-1)}{\text{Tr}[P_{\mathfrak{S}_N}^+]} P_{\mathfrak{S}_N}^a(\rho \otimes I_d^{\otimes(N-1)})(P_{\mathfrak{S}_N}^a)^* \\ &= \frac{dN(N+d-1)}{(N!)^2 \cdot \text{Tr}[P_{\mathfrak{S}_N}^+]} \sum_{\sigma, \tau \in \mathfrak{S}_N} (b_{\sigma(0)+1} \cdot b_{\tau^{-1}(0)+1}) \cdot \psi(\sigma)(\rho \otimes I_d^{\otimes(N-1)})\psi(\tau).\end{aligned}$$

As a consequence of [Lemma 3.8](#), for any $1 \leq a, b \leq N$, the matrix

$$(N-1)! \sum_{\sigma \in \Sigma_{a,b}} \psi(\sigma)^\Gamma,$$

is the Choi matrix of the linear map

$$X \mapsto \sum_{\substack{\mu, \nu \in \Sigma_{a,b} \\ \mu(0)=a-1 \\ \nu^{-1}(0)=b-1}} \psi(\mu) \cdot (X \otimes I_d^{\otimes(N-1)}) \cdot \psi(\nu).$$

This implies that the Choi matrix $C_{\Phi_{\text{opt}}^a}$ of Φ_{opt}^a is,

$$C_{\Phi_{\text{opt}}^a} = \frac{d(N+d-1)}{N! \cdot \text{Tr}[P_{\mathfrak{S}_N}^+]} \sum_{\substack{1 \leq a, b \leq N \\ \sigma \in \Sigma_{a,b}}} b_a b_b \cdot \psi(\sigma)^\Gamma.$$

Finally, using [Lemma 3.7](#), the equation $(d-1) \sum_{i=1}^N b_i^2 + (\sum_{i=1}^N b_i)^2 = 1$ satisfied by the positive real numbers b_i , and the relation $\text{Tr}[P_{\mathfrak{S}_{N-1}}^+] = \frac{N}{N+d-1} \text{Tr}[P_{\mathfrak{S}_N}^+]$, the trace condition on the Choi matrix $C_{\Phi_{\text{opt}}^a}$ holds, i.e.,

$$\text{Tr}_{[N+1] \setminus \{0\}} [C_{\Phi_{\text{opt}}^a}] = I_d.$$

Hence Φ_{opt}^a is trace preserving, and thus a quantum channel.

It remains to prove that the optimal quantum cloning channel Φ_{opt}^a saturates the upper bound,

$$\sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F((\Phi_{\text{opt}}^a)_i(\rho), \rho) \right] \leq \frac{\lambda_{\max}(R_a)}{d+1}.$$

For all marginals $(\Phi_{\text{opt}}^a)_i$, the average fidelity $\mathbb{E}_{\rho \in \Gamma} [F((\Phi_{\text{opt}}^a)_i(\rho), \rho)]$ becomes,

$$\begin{aligned} \mathbb{E}_{\rho \in \Gamma} [F((\Phi_{\text{opt}}^a)_i(\rho), \rho)] &= \mathbb{E}_{\rho \in \Gamma} \left[\text{Tr} \left[C_{\Phi_{\text{opt}}^a} (\rho_{(0)}^\top \otimes \rho_{(i)} \otimes I_d^{\otimes(N-1)}) \right] \right] \\ &= \frac{1}{d(d+1)} \text{Tr} \left[C_{\Phi_{\text{opt}}^a} \left((d^2 \cdot \mathbf{I}_{(0,i)} + d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} \right) \right] \\ &= \frac{\text{Tr} [C_{\Phi_{\text{opt}}^a}] + \text{Tr} [C_{\Phi_{\text{opt}}^a} (d \cdot \omega_{(0,i)} \otimes I_d^{\otimes(N-1)})]}{d(d+1)} \\ &= \frac{d + \text{Tr} [C_{\Phi_{\text{opt}}^a} (d \cdot \omega_{(0,i)} \otimes I_d^{\otimes(N-1)})]}{d(d+1)} \end{aligned}$$

Using the form of the Choi matrix $C_{\Phi_{\text{opt}}^a}$,

$$\begin{aligned} C_{\Phi_{\text{opt}}^a} (d \cdot \omega_{(0,i)} \otimes I_d^{\otimes(N-1)}) &= \frac{d(N+d-1)}{N! \cdot \text{Tr} [P_{\mathfrak{S}_N}^+]} \sum_{\substack{1 \leq a, b \leq N \\ \sigma \in \Sigma_{a,b}}} b_a b_b \cdot \left(\psi(\sigma)^\Gamma \cdot \psi((0 \ i))^\Gamma \right) \\ &= \frac{d(N+d-1)}{N! \cdot \text{Tr} [P_{\mathfrak{S}_N}^+]} \sum_{\substack{1 \leq a \leq N \\ \sigma \in \Sigma_{a,i}}} b_a \left((d-1)b_i + \sum_{1 \leq b \leq N} b_b \right) \psi(\sigma)^\Gamma. \end{aligned}$$

Such that taking the trace yields, using [Lemma 3.7](#),

$$\text{Tr} [C_{\Phi_{\text{opt}}^a} (d \cdot \omega_{(0,i)} \otimes I_d^{\otimes(N-1)})] = d \left((d-1)b_i + \sum_{j=1}^N b_j \right)^2,$$

and finally the average fidelity becomes,

$$\mathbb{E}_{\rho \in \Gamma} [F((\Phi_{\text{opt}}^a)_i(\rho), \rho)] = \frac{1 + ((d-1)b_i + \sum_{j=1}^N b_j)^2}{d+1}.$$

But since

$$\lambda_{\max}(S_a) = \langle \chi | S_a | \chi \rangle = \sum_{i=1}^N a_i \left((d-1)b_i + \sum_{j=1}^N b_j \right)^2,$$

then the equality $\sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} [F((\Phi_{\text{opt}}^a)_i(\rho), \rho)] = \frac{\lambda_{\max}(R_a)}{d+1}$ holds.

3.4.4 Optimal quantum cloning channel necessary condition

Let a be a direction vector and Φ_{opt}^a be the optimal quantum cloning channel for the universal $1 \rightarrow N$ quantum cloning problem in the direction a . For all marginals $(\Phi_{\text{opt}}^a)_i$, the average fidelity $f_i := \mathbb{E}_{\rho \in \Gamma} \left[F((\Phi_{\text{opt}}^a)_i(\rho), \rho) \right]$ is

$$f_i = \frac{1 + ((d-1)b_i + \sum_{j=1}^N b_j)^2}{d+1}.$$

Summing over all i yields,

$$\sum_{i=1}^N \sqrt{(d+1)f_i - 1} = (N+d-1) \sum_{i=1}^N b_i.$$

Using the equation $(d-1) \sum_{i=1}^N b_i^2 + (\sum_{i=1}^N b_i)^2 = 1$ satisfied by the positive real numbers b_i ,

$$\begin{aligned} \sum_{i=1}^N ((d+1)f_i - 1) &= \sum_{i=1}^N \left((d-1)b_i + \sum_{j=1}^N b_j \right)^2 \\ &= (d-1) + (N+d-1) \left(\sum_{i=1}^N b_i \right)^2 \\ &= (d-1) + \frac{\left(\sum_{i=1}^N \sqrt{(d+1)f_i - 1} \right)^2}{N+d-1}. \end{aligned}$$

Finally from the relation $f_i = p_i + \frac{1-p_i}{d}$, the necessary condition for the p_i 's of the optimal quantum cloning channels becomes,

$$N + (d^2 - 1) \sum_{i=1}^N p_i = d(d-1) + \frac{\left(\sum_{i=1}^N \sqrt{(d^2 - 1)p_i + 1} \right)^2}{N+d-1}. \quad (3.4)$$

3.4.5 The \mathcal{Q} -norm

Let $x \in \mathbb{R}^N$, and define its \mathcal{Q} -norm by

$$\|x\|_{\mathcal{Q}} := \frac{d\lambda_{\max}(S_x) - \|x\|_1}{d^2 - 1},$$

Necessary condition optimal quantum cloning channel




Figure 4 – The necessary condition [Eq. \(3.4\)](#) of the optimal quantum cloning channels for the universal $1 \rightarrow 2$ quantum cloning problem.

where $\|x\|_1 = \sum_{i=1}^N |x_i|$ is the ℓ_1 norm of the vector x and the matrix $S_x \in \mathcal{M}_{d^{N+1}}$ is given by,

$$S_x := \sum_{i=1}^N |x_i| \cdot (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)}.$$

Note that the \mathcal{Q} -norm depends implicitly on a parameter d , and for non-negative $x \in \mathbb{R}^N$, the matrix S_x coincides with that of [Lemma 3.9](#).

Remark. Note that on its own $\lambda_{\max}(S_x)$ defines a norm, but in general, except for trivial examples, subtracting two norms don't provide a new norm.

Proposition 3.11. *For all $N \geq 1$, the quantity $\|\cdot\|_{\mathcal{Q}}$ is a norm on \mathbb{R}^N .*

Proof. The absolute homogeneity of $\|\cdot\|_{\mathcal{Q}}$ comes from the fact that both $\lambda_{\max}(S_x)$ and $\|x\|_1$ are also absolutely homogeneous.

Let $x \in \mathbb{R}^N$ such that $\|x\|_{\mathcal{Q}} = 0$, then $d \cdot \lambda_{\max}(S_x) = \|x\|_1 = \text{Tr}[S_x]$. In particular all eigenvalues of S_x are equal to $\lambda_{\max}(S_x)$, and hence

$$S_x = \lambda_{\max}(S_x) \cdot I_d^{\otimes(N+1)},$$

with $\lambda_{\max}(S_x) \geq 0$. Let distinct $i, j = \{0, \dots, d-1\}$ and $|\psi\rangle$ defined by,

$$|\psi\rangle = |\underbrace{ij \cdots j}_{N \text{ times}}\rangle.$$

Then

$$\begin{aligned} \langle \psi | \sum_{i=1}^N |x_i| \cdot (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} | \psi \rangle &= 0 \\ \langle \psi | \lambda_{\max}(S_x) \cdot I_d^{\otimes(N+1)} | \psi \rangle &= \lambda_{\max}(S_x). \end{aligned}$$

Finally $\lambda_{\max}(S_x) = 0$ implies $x = 0$, and $\|\cdot\|_{\mathcal{Q}}$ is positive definite.

For the subadditivity of $\|\cdot\|_{\mathcal{Q}}$, it is sufficient to prove that for any $x, y \in \mathbb{R}^N$ such that $\|x\|_{\mathcal{Q}} \leq 1$ and $\|y\|_{\mathcal{Q}} \leq 1$, then $\|\frac{x+y}{2}\|_{\mathcal{Q}} \leq 1$. It is possible to assert, without loss of generality, that by multiplying the vectors by a scalar, the vector $x + y$ is in \mathbb{R}_+^N . Let $x' \in \mathbb{R}_+^N$ be defined by

$$x' := \begin{cases} [x, \frac{x+y}{2}] \cap \partial \mathbb{R}_+^N & \text{if } [x, \frac{x+y}{2}] \cap \partial \mathbb{R}_+^N \neq \emptyset \\ x & \text{otherwise} \end{cases}$$

and similarly for $y' \in \mathbb{R}_+^N$, where $\partial \mathbb{R}_+^N$ denote the boundary of \mathbb{R}_+^N . Then $\frac{x+y}{2} \in [x', y']$, and there exists $\lambda \in [0, 1]$ such that $\frac{x+y}{2} = \lambda \cdot x' + (1 - \lambda) \cdot y'$. Then using,

$$\begin{aligned} &\lambda_{\max} \left(\sum_{i=1}^N (x'_i + y'_i) \cdot (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} \right) \\ &\leq \lambda_{\max} \left(\sum_{i=1}^N x'_i \cdot (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} \right) + \lambda_{\max} \left(\sum_{i=1}^N y'_i \cdot (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} \right), \end{aligned}$$

and the absolute homogeneity of $\|\cdot\|_{\mathcal{Q}}$,

$$\left\| \frac{x+y}{2} \right\|_{\mathcal{Q}} \leq \lambda \cdot \|x'\|_{\mathcal{Q}} + (1 - \lambda) \cdot \|y'\|_{\mathcal{Q}} \leq \max(\|x'\|_{\mathcal{Q}}, \|y'\|_{\mathcal{Q}}).$$

Now there exists $t_x \in [0, 1]^N$ and $t_y \in [0, 1]^N$ such that element-wise product equations $x' = t_x \cdot x$ and $y' = t_y \cdot y$ hold. It remains to prove that, for all $t \in [0, 1]^N$ and $x \in \mathbb{R}_+^N$, the inequality $\|t \cdot x\|_{\mathcal{Q}} \leq \|x\|_{\mathcal{Q}}$ hold. With this both $\|x'\|_{\mathcal{Q}} \leq 1$ and $\|y'\|_{\mathcal{Q}} \leq 1$, such that $\|\frac{x+y}{2}\|_{\mathcal{Q}} \leq 1$.

Let $x, y \in \mathbb{R}_+^N$ such that $x_i \leq y_i$ for all $i \in \{1, \dots, N\}$, then $\|x\|_{\mathcal{Q}} \leq \|y\|_{\mathcal{Q}}$. Indeed, let $\chi = \sum_{i=1}^N b_i \cdot (\sqrt{d} \cdot |\Omega\rangle_{(0,i)} \otimes |v\rangle)$ be normalized largest eigenvectors of S_x , from [Lemma 3.9](#), then using that $|v\rangle \in \vee^{(N-1)} \mathbb{C}^d$ and then all the b_i are positive real numbers,

$$\begin{aligned} & \langle \chi | (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} | \chi \rangle \\ &= \sum_{k,j=1}^N (d^2 b_k b_l) \cdot (\langle \Omega |_{(0,i)} \otimes \langle v |) (\omega_{(0,i)} \otimes I_d^{\otimes(N-1)}) (|\Omega\rangle_{(0,i)} \otimes |v\rangle) \\ &= \left((d-1)b_i + \sum_{j=1}^N b_j \right)^2 \\ &\geq \left(\sum_{j=1}^N b_j \right)^2. \end{aligned}$$

Now, using the equation $(d-1) \sum_{i=1}^N b_i^2 + \left(\sum_{i=1}^N b_i \right)^2 = 1$ satisfied by the positive real numbers b_i ,

$$\begin{aligned} 1 &= \langle \chi | \chi \rangle \\ &= (d-1) \sum_{i=1}^N b_i^2 + \left(\sum_{i=1}^N b_i \right)^2 \\ &\leq d \left(\sum_{i=1}^N b_i \right)^2. \end{aligned}$$

this implies that $\langle \chi | (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} | \chi \rangle \geq \frac{1}{d}$. This way since $x_i \leq y_i$ for all $i \in \{1, \dots, N\}$, then $\lambda_{\max}(S_y) \geq \langle \chi | S_y | \chi \rangle$ and,

$$\begin{aligned} & \lambda_{\max}(S_y) - \lambda_{\max}(S_x) \langle \chi | S_y - S_x | \chi \rangle \\ &= \sum_{i=1}^N (y_i - x_i) \cdot \langle \chi | (d \cdot \omega_{(0,i)}) \otimes I_d^{\otimes(N-1)} | \chi \rangle \\ &\geq \sum_{i=1}^N (y_i - x_i) \frac{1}{d}. \end{aligned}$$

Then $\|y\|_{\mathcal{Q}} = \lambda_{\max}(S_y) - \frac{1}{d}\|y\|_1 \geq \lambda_{\max}(S_x) - \frac{1}{d}\|x\|_1 = \|x\|_{\mathcal{Q}}$, which conclude the proof since $t \cdot y = x$ for some $t \in [0, 1]^N$.

With the help of the \mathcal{Q} -norm, the upper bound from [Theorem 3.5](#) can be reformulated as: for any direction vector $a \in [0, 1]^N$ the universal quantum cloning problem is upper bounded by

$$\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F(\Phi_i(\rho), \rho) \right] \leq \frac{1}{d}\|a\|_1 + \left(1 - \frac{1}{d}\right)\|a\|_{\mathcal{Q}}.$$

Let the **dual** \mathcal{Q} -norm, be defined on $y \in \mathbb{R}^N$ by,

$$\|y\|_{\mathcal{Q}}^* := \sup_{\substack{x \in \mathbb{R}^N \\ x \neq 0}} \frac{\langle y, x \rangle}{\|x\|_{\mathcal{Q}}}.$$

\mathcal{Q} -norm & dual \mathcal{Q} -norm 1-dimensional unit spheres [$d = 2$]




Figure 5 – The 1-dimensional unit spheres of the \mathcal{Q} -norm ■ and the dual \mathcal{Q} -norm ■.

3.4.6 Achievable fidelity region

In this Section, the characterization of the achievable fidelity region for the universal $1 \rightarrow N$ quantum cloning problem will be given geometrically, as the non-negative part of the unit ball of the dual \mathcal{Q} -norm.

Let $\mathcal{R}_{N,d}$ be the achievable fidelity region for the universal $1 \rightarrow N$ quantum cloning problem, that is,

$$\mathcal{R}_{N,d} := \left\{ p \in [0, 1]^N \mid \exists \Phi : \mathcal{M}_d \xrightarrow{\text{CPTP}} (\mathcal{M}_d)^{\otimes N} \text{ s.t. } \Phi_i(\rho) = p_i \cdot \rho + (1 - p_i) \frac{I_d}{d} \right\}.$$

Theorem 3.12. $\mathcal{R}_{N,d}$ is the non-negative part of the unit ball of the dual \mathcal{Q} -norm, i.e. $p \in [0, 1]^N$ is in $\mathcal{R}_{N,d}$ if and only if $\|p\|_{\mathcal{Q}}^* \leq 1$.

Proof. The following more general result will be first proved. Let X be a finite dimensional real vector space and let X^* denote its dual space. Let $\|\cdot\|_1$ be a norm on X , let $\|\cdot\|_2$ be a norm on X^* , and define the dual norms as follows,

$$\|y\|_1^* := \sup_{\substack{x \in X \\ x \neq 0}} \frac{\langle y, x \rangle}{\|x\|_1}, \quad \forall y \in X^*,$$

$$\|y\|_2^* := \sup_{\substack{x \in X^* \\ x \neq 0}} \frac{\langle y, x \rangle}{\|x\|_2}, \quad \forall y \in X.$$

Provided that both conditions,

$$\forall x \in X, \forall y \in X^*, \quad \langle x, y \rangle \leq \|x\|_1 \|y\|_2 \quad (3.5)$$

$$\forall y \in X^*, \exists x \in X, \quad \langle x, y \rangle = \|x\|_1 \|y\|_2, \quad (3.6)$$

are satisfied, then $\forall x \in X, \|x\|_1 = \|x\|_2^*$. In particular the unit balls of $\|\cdot\|_1$ and $\|\cdot\|_2^*$ are equal. Indeed, from Eq. (3.5), then $\forall x \in X$,

$$\|x\|_1 \geq \sup_{y \in X^*} \frac{\langle x, y \rangle}{\|y\|_2} = \|x\|_2^*,$$

and from Eq. (3.6), then $\forall y \in X^*, \exists x \in X$,

$$\|x\|_1 = \frac{\langle x, y \rangle}{\|y\|_2} \leq \sup_{y \in X^*} \frac{\langle x, y \rangle}{\|y\|_2} \leq \|x\|_2^*,$$

which conclude the proof of this first result.

Now to prove the theorem, define the norm $\|\cdot\|_{\mathcal{R}}$ on $x \in \mathbb{R}^N$ by,

$$\|x\|_{\mathcal{R}} := \max \left\{ t > 0 \mid t \cdot |x| \in \mathcal{R}_{N,d} \right\},$$

where $t \cdot |x|$ denote the vector of \mathbb{R}^N with coefficients $t \cdot |x_i|$. The norms $\|\cdot\|_{\mathcal{R}}$ and $\|\cdot\|_{\mathcal{Q}}$ are as in the settings above. If both conditions [Eq. \(3.5\)](#) and [Eq. \(3.6\)](#), i.e.

$$\begin{aligned} \forall p \in \mathbb{R}^N, \forall a \in \mathbb{R}^N, \quad \langle p, a \rangle &\leq \|p\|_{\mathcal{R}} \|a\|_{\mathcal{Q}} \\ \forall a \in \mathbb{R}^N, \exists p \in \mathbb{R}^N, \quad \langle p, a \rangle &= \|p\|_{\mathcal{R}} \|a\|_{\mathcal{Q}}, \end{aligned}$$

are satisfied, then the theorem holds.

Let $p \in \mathcal{R}_{N,d}$ and a quantum cloning channel $\Phi_p : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ such that for all marginals $(\Phi_p)_i$ and all pure quantum states ρ ,

$$(\Phi_p)_i(\rho) = p_i \cdot \rho + (1 - p_i) \frac{I_d}{d}.$$

Then for all direction vectors $a \in [0, 1]^N$,

$$\begin{aligned} \frac{1}{d} \|a\|_1 + \left(1 - \frac{1}{d}\right) \|a\|_{\mathcal{Q}} &\geq \sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F(\Phi_i(\rho), \rho) \right] \\ &\geq \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F((\Phi_p)_i(\rho), \rho) \right] \\ &= \frac{1}{d} \|a\|_1 + \left(1 - \frac{1}{d}\right) \langle p, a \rangle. \end{aligned}$$

This gives $\langle p, a \rangle \leq \|a\|_{\mathcal{Q}}$ for all direction vectors $a \in [0, 1]^N$. For arbitrary $a \in \mathbb{R}^N$, note that

$$\langle p, a \rangle \leq \langle p, |a| \rangle \leq \| |a| \|_{\mathcal{Q}} = \|a\|_{\mathcal{Q}},$$

showing that the condition [Eq. \(3.5\)](#) is satisfied, since $\|p\|_{\mathcal{R}} = 1$.

Let some direction vector $a \in [0, 1]^N$, then from [Theorem 3.10](#), there a quantum cloning channel Φ_{opt}^a such that,

$$\sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F((\Phi_{\text{opt}}^a)_i(\rho), \rho) \right] = \frac{1}{d} \|a\|_1 + \left(1 - \frac{1}{d}\right) \|a\|_{\mathcal{Q}},$$

with some $p \in \mathcal{R}_{N,d}$, such that,

$$\sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F((\Phi_{\text{opt}}^a)_i(\rho), \rho) \right] = \frac{1}{d} \|a\|_1 + \left(1 - \frac{1}{d}\right) \langle p, a \rangle.$$

This shows that condition [Eq. \(3.6\)](#) is also satisfied.

From [Theorem 3.12](#), the achievable fidelity region $\mathcal{R}_{N,d}$ is a convex set delimited by a family of hyperplanes (see [Figure 6](#)):

$$\langle p, a \rangle = \|a\|_{\mathcal{Q}}.$$

$\mathcal{R}_{2,2}$ is delimited by a family of hyperplanes




Figure 6 – $\mathcal{R}_{2,2}$ is a convex set delimited by a family of hyperplanes $\langle p, a \rangle = \|a\|_{\mathcal{Q}}$.

Note that within the formulation of the quantum cloning problem, i.e.

$$\sup_{\Phi \text{ CPTP}} \sum_{i=1}^N a_i \cdot \mathbb{E}_{\rho \in \Gamma} \left[F(\Phi_i(\rho), \rho) \right],$$

the optimal quantum cloning channels are twirled quantum channels, and thus their marginals are of the form $\rho \mapsto p_i \cdot \rho + (1 - p_i) \frac{I_d}{d}$, but the p_i 's are not asked to be

collinear with the a_i 's. Instead they have to maximize,

$$\sum_{i=1}^N a_i \cdot \left(p_i + \frac{1 - p_i}{d} \right).$$

The optimal quantum cloning channels from [Theorem 3.10](#), can indeed give p_i 's in a different direction than the a_i 's, especially if the direction of the a_i does not intersect an extreme point of $\mathcal{R}_{N,d}$. As a consequence, the optimal quantum cloning channels from [Theorem 3.10](#) do not fill the boundary of $\mathcal{R}_{N,d}$, since some points in this boundary are not optimal with respect to the optimisation problem.

For example, from [Theorem 3.4.2](#),

$$\max \{ p \in [0, 1] \mid (p, p, 0) \in \mathcal{R}_{3,2} \} = \frac{2}{3},$$

that is the p_{opt} for the symmetric universal $1 \rightarrow 2$ quantum cloning problem is $2/3$. However $(2/3, 2/3, 0) \in \mathcal{R}_{3,2}$ is not optimal for the asymmetric universal $1 \rightarrow 3$ quantum cloning problem, given the direction vector $a := (1/2, 1/2, 0)$. Indeed, the the optimal quantum cloning channel in this direction, from [Theorem 3.10](#), gives $(2/3, 2/3, 1/9) \in \mathcal{R}_{3,2}$ (see [Figure 7](#)).




Figure 7 – $\mathcal{R}_{3,2}$ is a convex set delimited by a family of hyperplanes, but some points may intersect more than 1 hyperplanes. Here is a view of the slice (p, p, q) of $\mathcal{R}_{3,2}$, with a flat region.

In conclusion, as the previous example shows, given a $p \in [0, 1]^N$, in order to decide whether p is in the achievable fidelity region $\mathcal{R}_{N,d}$, it does not suffice to check the necessary condition [Eq. \(3.4\)](#).

Extendibility

Chapter contents

4.1	Extendibility of quantum states on a graph	87
4.2	Quantum cloning: star-graph extendibility of isotropic states	88
4.3	K_N -Extendibility of isotropic states	89
4.3.1	Lower bound	89
4.3.2	Symmetries	91
4.3.3	Dual SDP	92
4.3.4	Optimal value	101

The present Section discusses the results of the papers “Monogamy of highly symmetric states” [All+23], of which I am a co-author. At the time of writing this manuscript, that paper was in preparation. Only the section on the extendibility of isotropic states is presented below.

In recent years, the problem of entanglement in many-body systems has received particular attention in quantum physics and quantum information theory. This complex but fundamental quantum phenomenon has been the subject of much work because of its essential role in the description and understanding of quantum many-body systems [WVI03; BV07; Ami+08; Tab+22].

4.1 Extendibility of quantum states on a graph

Let G be a graph with N vertices. A quantum state on the graph G is a quantum state ρ on a N -fold composite quantum system, each associated to a vertex of G , i.e.

$$\rho \in (\mathbb{C}^d)^{\otimes N}.$$

Let $e := (u, v)$ be an edge of the graph G , the reduced quantum state ρ_e is the reduced quantum state on quantum systems u and v , i.e.

$$\rho_e := \text{Tr}_{G \setminus \{u, v\}} [\rho].$$

The **complete graph** on N vertices, denoted K_N is the graph G where for all distinct vertices u and v in G , the pair (u, v) is an edge of G , i.e. every pair of distinct vertices is connected by a unique edge (see Figure 8a). The complete graph K_N has $\frac{N(N-1)}{2}$ edges.

The **star graph** on N vertices, denoted S_N is the graph G with a distinct **central** vertex $v \in G$ such that the pair (u, v) is an edge of G for all other vertex $u \in G$, i.e. it is the 1-depth tree of order N (see Figure 8b). The star graph S_N has $n - 1$ edges.

From Section 2.2.4, a bipartite quantum state ρ is k -extendible if there exists a quantum state σ on the star graph S_{k+1} such that for all edges e , the reduced quantum states on e is,

$$\sigma_e = \rho.$$



Figure 8 – Complete graph on 5 vertices K_5 , and star graph on 6 vertices S_6 .

4.2 Quantum cloning: star-graph extendibility of isotropic states

The quantum cloning problem can be seen as an extendibility problem, when considering the Choi matrix of the quantum cloning channel. Indeed, a perfect $1 \rightarrow N$ quantum cloning channel $\Phi : \mathcal{M}_d \rightarrow (\mathcal{M}_d)^{\otimes N}$ would have a Choi matrix C_Φ living on a star graph S_{N+1} such that for all edges e , the reduced Choi matrix on e is,

$$(C_\Phi)_e = d \cdot \omega.$$

Because of the no-cloning [Theorem 3.1](#), or alternatively the monogamy of the entanglement from [Section 2.2.4](#), the normalized reduced Choi matrix on e is required to be instead an isotropic state,

$$\frac{1}{d}(C_\Phi)_e = \lambda_e \cdot \omega + (1 - \lambda_e)I,$$

where I is the maximally mixed state on $\mathbb{C}^d \otimes \mathbb{C}^d$

The symmetric $1 \rightarrow N$ quantum cloning problem can then be solved using a **semi-definite programming** (SDP) optimization problem:

$$\begin{aligned} \max_{C_\Phi, p} \quad & p \\ \text{s.t.} \quad & (C_\Phi)_e = d(p \cdot \omega + (1 - p)I), & \text{for all edges } e \\ & \text{Tr}_{[N+1] \setminus \{0\}} [C_\Phi] = I_d, \quad C_\Phi \geq 0, \quad p \in [0, 1]. \end{aligned}$$

Note that in the previous optimization problem, the conditions $\text{Tr}_{[N+1]\setminus\{0\}} [C_\Phi] = I_d$ and $p \in [0, 1]$ are superfluous, and the optimization problem reduces to,

$$\begin{aligned} \max_{C_\Phi, p} \quad & p \\ \text{s.t.} \quad & (C_\Phi)_e = d(p \cdot \omega + (1 - p)I), \quad \text{for all edges } e \\ & C_\Phi \geq 0. \end{aligned}$$

From [Theorem 3.4.2](#), this optimization problem has optimal solution:

$$p_{\text{opt}} := \frac{d + N}{N(d + 1)}.$$

4.3 K_N -Extendibility of isotropic states

The optimization problem of [Section 4.2](#), can be extended to the following quantum state marginal problem. Given a complete graph K_N , what is the largest $p \in [0, 1]$, such that there exists a quantum state ρ on K_N , with all reduced quantum state $\rho_e = p \cdot \omega + (1 - p)I$, for all edges e ? This problem depends on the number of quantum systems N and their dimension d , and can be stated as the primal semi-definite problem:

$$\begin{aligned} p(N, d) := \max_{\rho, p} \quad & p \\ \text{s.t.} \quad & \rho_e = p \cdot \omega + (1 - p)I, \quad \text{for all edges } e \\ & \rho \geq 0. \end{aligned} \tag{4.1}$$

4.3.1 Lower bound

A **perfect matching** on a graph is a set of edges, such that every vertex is contained in exactly one of those edges.

Proposition 4.1. *There are $(2N - 1)!!$ perfect matchings on K_{2N} , and if e is an edge on K_{2N} , then there are $(2N - 3)!!$ perfect matchings on K_{2N} containing e .*

Proof. Let a_N be the number of perfect matching on K_{2N} ; clearly $a_1 = 1$. Now assume $N > 1$ and let v be a vertex in K_{2N} . This vertex can be matched with $2N - 1$ other vertices, let u be such other vertex matched with v . Remove u and v from K_{2N} , the resulting graph $K_{2N} \setminus \{u, v\}$ is the complete graph $K_{2(N-1)}$.

Thus, by induction on N , the number of perfect matchings on K_{2N} satisfies the recursive relation:

$$a_N = (2N - 1)a_{N-1} \implies a_N = (2N - 1)!!.$$

Assume $e = (u, v)$, thus the number of perfect matchings containing e is the number of perfect matchings on $K_{2N} \setminus \{u, v\}$, that is $(2N - 3)!!$.

■ *Remark.* There is no perfect matching on K_N for odd N (see Figure 9).



Figure 9 – Perfect matching ■ for complete graphs K_N , with even N . For K_N with odd N , some vertices are not matched ■.

A lower bound on the optimisation problem Eq. (4.1) would be the following. For even N , let $E_1, \dots, E_{(N-1)!!}$ be all the perfect matchings on K_N , and for each perfect matching E_k , define the quantum state $\rho^{(k)}$ on K_N by,

$$\rho^{(k)} := \bigotimes_{e \in E_k} \omega_e.$$

For odd N , let v be vertex of K_N , and let $E_{v,1}, \dots, E_{v,(N-2)!!}$ be all the perfect matchings on $K_N \setminus \{v\}$, define the quantum state $\rho^{(v,k)}$ on K_N by,

$$\rho^{(v,k)} := \bigotimes_{e \in E_{v,k}} \omega_e \otimes \mathbb{I}_v.$$

That is a quantum state maximally entangled on the perfect matching edges, and maximally mixed on the remaining vertex in the odd case. Let ρ be the quantum

state defined on K_N , as a uniform combination of the previously constructed states:

$$\begin{aligned} (N \text{ even}) \quad \rho &:= \frac{1}{N-1} \sum_{1 \leq k \leq (N-1)!!} \rho^{(k)} \\ (N \text{ odd}) \quad \rho &:= \frac{1}{N} \sum_{\substack{v \in K_N \\ 1 \leq k \leq (N-2)!!}} \rho^{(v,k)}, \end{aligned}$$

where the corresponding normalization factors can be found using [Proposition 4.1](#). Then for all edges e in K_N , the reduced quantum state ρ_e is

$$\rho_e = \begin{cases} \frac{1}{N-1}\omega + \frac{N-2}{N-1}\mathbf{I} & N \text{ even} \\ \frac{1}{N}\omega + \frac{N-1}{N}\mathbf{I} & N \text{ odd.} \end{cases}$$

The lower bound becomes

$$p(N, d) \geq \begin{cases} \frac{1}{N-1} & N \text{ even} \\ \frac{1}{N} & N \text{ odd.} \end{cases}$$

In particular, the lower bound is independent of the dimension d .

4.3.2 Symmetries

Since it is a complete graph, any permutation of the vertices of K_N is a graph isomorphism. Let ρ be a quantum state on K_N such that for all edges e , the reduced quantum state ρ_e is,

$$\rho_e = p \cdot \omega + (1-p)\mathbf{I}.$$

Then ρ is invariant by vertex permutation, that is for all permutations $\sigma \in \mathfrak{S}_N$, the equality

$$\psi(\sigma) \cdot \rho \cdot \psi(\sigma^{-1}) = \rho,$$

holds.

Recall that a maximally entangled state ω is invariant by $\bar{U} \otimes U$, for all unitary matrix U , i.e. $(\bar{U} \otimes U) \cdot \omega = \omega$. On the star graph S_N with central vertex v , a quantum state ρ with reduced quantum states $\rho_e = p \cdot \omega + (1-p)\mathbf{I}$, for all edges e , would be invariant under conjugation by \bar{U} on v and $U^{\otimes(N-1)}$ on $S_N \setminus \{v\}$, i.e.

$$\left(\bar{U}_v \otimes (U^{\otimes(N-1)})_{S_N \setminus \{v\}} \right) \cdot \rho \cdot \left(U_v^\top \otimes ((U^*)^{\otimes(N-1)})_{S_N \setminus \{v\}} \right) = \rho.$$

But on the complete graph K_N , no distinct vertex can be chosen, instead, using the invariant on a maximally entangled state by $O \otimes O$, for all orthogonal matrix $O \in O_d$, i.e. $(O \otimes O) \cdot \omega = \omega$, a quantum state ρ with reduced quantum states $\rho_e = p \cdot \omega + (1 - p)\mathbf{I}$, for all edges e , is invariant under conjugation by $O^{\otimes N}$, that is,

$$O^{\otimes N} \cdot \rho \cdot (O^\top)^{\otimes N} = \rho.$$

Hence, with respect to the optimisation problem [Eq. \(4.1\)](#), the two commutation relations,

$$[\rho, \sigma] = 0, \quad \forall \sigma \in \mathfrak{S}_N \quad (4.2)$$

$$[\rho, O^{\otimes N}] = 0, \quad \forall O \in O_d. \quad (4.3)$$

hold for all N and d .

4.3.3 Dual SDP

The Lagrangian [\[BBV04\]](#) associated with the semi-definite problem [Eq. \(4.1\)](#) is defined as,

$$L(p, \rho, h_e, Z) := p + \sum_{e \text{ edge}} \langle h_e, \rho_e - p \cdot \omega + (1 - p)\mathbf{I} \rangle + \langle Z, \rho \rangle.$$

The **min-max principle** states that,

$$\max_{\rho, p} \min_{h_e, Z} L(p, \rho, h_e, Z) \leq \min_{h_e, Z} \max_{\rho, p} L(p, \rho, h_e, Z).$$

Then, the **dual** semi-definite problem of [Eq. \(4.1\)](#) is,

$$\begin{aligned} p^*(N, d) := \max_{h_e, Z} & \quad - \frac{\text{Tr} \left[\sum_{e \text{ edge}} h_e \otimes I_d^{\otimes (N-2)} \right]}{d^N} \\ \text{s.t.} & \quad \sum_{e \text{ edge}} \text{Tr} [h_e \cdot (\omega - \mathbf{I})] = 1 \\ & \quad Z + \sum_{e \text{ edge}} h_e \otimes I_d^{\otimes (N-2)} = 0, \quad Z \geq 0, \end{aligned}$$

where for each edge e of K_N , the matrix $h_e \in \mathcal{M}_{d^2}$ is Hermitian, as well as the matrix $Z \in \mathcal{M}_{d^N}$. Recall that for any Hermitian matrix M the smallest $\lambda \in \mathbb{R}$ such

that $\lambda \cdot I \geq M$ is equal to the largest eigenvalue of M . Then the dual optimization problem can be simplified to,

$$\begin{aligned} p^*(N, d) = \min_{h_e} \quad & \lambda_{\max} \left(\sum_{e \text{ edge}} h_e \otimes I_d^{\otimes (N-2)} \right) \\ \text{s.t.} \quad & \sum_{e \text{ edge}} \text{Tr} [h_e \cdot \omega] = 1 \\ & \text{Tr} [h_e] = 0, \quad \text{for all edges } e. \end{aligned}$$

From the commutation relation [Eq. \(4.3\)](#) applied to the dual optimization problem, define for all edge e , the twirled Hermitian matrix \tilde{h}_e :

$$\tilde{h}_e := \int_O (O \otimes O)(h_e)(O^\top \otimes O^\top) dO,$$

where the integral is taken with respect to the normalized Haar measure on the orthogonal group O_d . The constraints of the dual optimization problem are also satisfied with \tilde{h}_e , and since by convexity of λ_{\max} ,

$$\lambda_{\max} \left(\sum_{e \text{ edge}} h_e \otimes I_d^{\otimes (N-2)} \right) \geq \lambda_{\max} \left(\sum_{e \text{ edge}} \tilde{h}_e \otimes I_d^{\otimes (N-2)} \right),$$

hence the dual optimization problem can be restricted to twirled \tilde{h}_e .

The twirled \tilde{h}_e commutes with all the orthogonal matrices $O \otimes O$, i.e. they are in the commutant of the algebra

$$\text{Span}_{\mathbb{C}} \{O \otimes O \mid O \in O_d\}.$$

From [Theorem 1.3.3](#), the twirled \tilde{h}_e is in the complex algebra spanned by the tensor representation of the Brauer monoid:

$$\text{Span}_{\mathbb{C}} \{\psi(p) \mid p \in \mathbb{B}_2\}.$$

Hence, the twirled \tilde{h}_e are a linear combination of the unnormalized maximally mixed, maximally entangled, and flip state:

$$\tilde{h}_e = \alpha_e(d^2 \cdot \mathbf{I}_e) + \beta_e(d \cdot \omega_e) + \gamma_e(d \cdot F_e),$$

subject to the constraint $\alpha_e = -\frac{\beta_e + \gamma_e}{d}$. The dual optimization problem simplifies to,

$$\begin{aligned} p^*(N, d) = \min_{\beta_e, \gamma_e} \quad & \lambda_{\max} \left(\sum_{e \text{ edge}} \left(-\frac{\beta_e + \gamma_e}{d} (d^2 \cdot \mathbf{I}_e) + \beta_e (d \cdot \omega_e) + \gamma_e (d \cdot \mathbf{F}_e) \right) \otimes I_d^{\otimes(N-2)} \right) \\ \text{s.t.} \quad & (d^2 - 1) \left(\sum_{e \text{ edge}} \beta_e \right) + (d - 1) \left(\sum_{e \text{ edge}} \gamma_e \right) = d. \end{aligned}$$

From the commutation relation [Eq. \(4.2\)](#), applied to the dual optimization problem, all the β_e and γ_e must be equal. Writing $\beta := \beta_e$ and $\gamma := \gamma_e$ for all edges e , the dual optimization problem simplifies further to

$$\begin{aligned} p^*(N, d) = \min_{\beta, \gamma} \quad & \lambda_{\max} \left(\sum_{e \text{ edge}} \left(-\frac{\beta + \gamma}{d} (d^2 \cdot \mathbf{I}_e) + \beta (d \cdot \omega_e) + \gamma (d \cdot \mathbf{F}_e) \right) \otimes I_d^{\otimes(N-2)} \right) \\ \text{s.t.} \quad & \frac{N(N-1)}{2} (\beta(d^2 - 1) + \gamma(d - 1)) = d. \end{aligned}$$

Using the constraint $\frac{N(N-1)}{2} (\beta(d^2 - 1) + \gamma(d - 1)) = d$ to eliminate γ , let the function f be defined by $f(x) := \lambda_{\max}(H(x))$ with

$$H(x) := \sum_{e \text{ edge}} \left(\left(\frac{2}{N(N-1)(1-d)} - x \right) ((d^2 \cdot \mathbf{I}_e) - d(d \cdot \mathbf{F}_e)) + x((d \cdot \mathbf{F}_e) - (d \cdot \omega_e)) \right) \otimes I_d^{\otimes(N-2)}.$$

Then the dual optimization problem becomes finally the minimization,

$$p^*(N, d) = \min_x f(x).$$

Recall from [Theorem 1.2](#), that under the action of the tensor representation of the symmetric group algebra \mathbb{S}_N , the space of N -fold tensors over \mathbb{C}^d decomposes as

$$(\mathbb{C}^d)^{\otimes N} \simeq \bigoplus_{\lambda \in \text{Irr}(\mathfrak{S}_N)} V_{\lambda}^{\oplus m_{\lambda}},$$

where $\text{Irr}(\mathfrak{S}_N) := \{\lambda \vdash N \mid \lambda'_1 \leq d\}$. Similarly, as a consequence of [Theorem 1.3.3](#), under the action of the tensor representation of the Brauer algebra $\mathbb{B}_N(d)$, the space of N -fold tensors over \mathbb{C}^d decomposes as

$$(\mathbb{C}^d)^{\otimes N} \simeq \bigoplus_{\lambda \in \text{Irr}(\mathbb{B}_N(d))} W_{\lambda}^{\oplus n_{\lambda}},$$

where $\text{Irr}(\mathbb{B}_N(d)) := \{\lambda \vdash N - 2k \mid k \in \{0, \dots, \lfloor \frac{N}{2} \rfloor\} \text{ and } \lambda'_1 + \lambda'_2 \leq d\}$ [Wen88]. Both decompositions are indexed by some Young diagram λ .

Given a irreducible representation W_λ of the Brauer algebra $\mathbb{B}_N(d)$, the restriction of W_λ to the symmetric group algebra \mathbb{S}_N , as a subalgebra decompose into

$$\text{Res}_{\mathbb{S}_N}^{\mathbb{B}_N(d)}(W_\lambda) \simeq \bigoplus_{\mu \in \text{Irr}(\mathbb{S}_N)} V_\lambda^{\oplus m_{\lambda, \mu}}.$$

The multiplicities have no known concise formula. Even the explicit characterization of the set Ω ,

$$\Omega := \{(\lambda, \mu) \in \text{Irr}(\mathbb{B}_N(d)) \times \text{Irr}(\mathbb{S}_N) \mid m_{\lambda, \mu} \neq 0\},$$

is still unknown. However Okada [Oka16] characterizes it using an algorithm, and found the subset $\Gamma \subseteq \Omega$,

$$\Gamma := \{(\lambda, \mu) \in \text{Irr}(\mathbb{B}_N(d)) \times \text{Irr}(\mathbb{S}_N) \mid \lambda = (1^m), r(\mu) = m \text{ for some } m \in \{1, \dots, d\}\},$$

where $r(\mu)$ is the number of rows with odd size in the Young diagram μ . Moreover if $((1^m), \mu) \in \Omega$ then $((1^m), \mu) \in \Gamma$.

The **content** of a Young diagram λ , denoted $c(\lambda)$, is the the sum of the entries of the boxes of the Young tableau T , where the entry of the box in row i and column j is given by $j - i$. For example if $\lambda := (3, 3, 1)$, then

$$\lambda = \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \qquad T = \begin{array}{|c|c|c|} \hline 0 & 1 & 2 \\ \hline -1 & 0 & 1 \\ \hline -2 & & \\ \hline \end{array}$$

and the content of λ is $c(\lambda) = 1$.

The next lemmas describe how particular central elements of the symmetric group \mathbb{S}_N and the Brauer monoid \mathbb{B}_N , act on their irreducible representations.

Lemma 4.2 ([DLS19]). *Let V_λ be an irreducible representation of \mathbb{S}_N , with $\lambda \in \text{Irr}(\mathbb{S}_N)$, and define J an element of the symmetric group algebra \mathbb{S}_N , by*

$$J := \sum_{1 \leq i, j \leq N} (i \ j).$$

Then the restriction of J on V_λ is the multiple of the identity,

$$J|_{V_\lambda} = c(\lambda) \cdot I.$$

Lemma 4.3 ([DLS19]). Let W_λ be an irreducible representation of $\mathbb{B}_N(d)$, with $\lambda \in \text{Irr}(\mathbb{B}_N(d))$, and define J an element of the Brauer algebra $\mathbb{B}_N(d)$, by

$$J := \sum_{1 \leq i, j \leq N} (i \ j) - (i \ j)^\Gamma,$$

where $(i \ j)^\Gamma$ denotes the partial transposition of the diagram $(i \ j)$. Then the restriction of J on W_λ is the multiple of the identity,

$$J|_{W_\lambda} = (c(\lambda) - k(d-1)) \cdot I.$$

Using Lemma 4.2, Lemma 4.3, the decomposition of the space of N -fold tensors over \mathbb{C}^d under the action of the tensor representation of the Brauer algebra $\mathbb{B}_N(d)$, and decomposing of the restriction of the irreducible representations of the Brauer algebra $\mathbb{B}_N(d)$ to the symmetric group algebra \mathbb{S}_N , the function f becomes,

$$f(x) = \max_{(\lambda, \mu) \in \Omega} \underbrace{\frac{1}{d-1} \left(\frac{2d \cdot c(\mu)}{N(N-1)} - 1 \right) + x \left(c(\lambda) + d \cdot c(\mu) - k(d-1) - \frac{N(N-1)}{2} \right)}_{f_{\lambda, \mu}(x)},$$

where $f_{\lambda, \mu}(x)$ is an affine function. The dual optimization problem is then the minimum value of the max over a set of affine functions, i.e.,

$$p^*(N, d) = \min_x \max_{(\lambda, \mu) \in \Omega} f_{\lambda, \mu}(x). \quad (4.4)$$

Case $d > N$ or either d or N is even

An upper bound for the dual optimization problem Eq. (4.4), can be found by setting $x = \frac{2}{N(N-1)(1-d)}$. Then Eq. (4.4) becomes, since $x < 0$,

$$p^*(N, d) \leq \min_{\lambda \in \text{Irr}(\mathbb{B}_N(d))} \frac{2}{N(N-1)(1-d)} (c(\lambda) - k(d-1)).$$

Lemma 4.4. If $d > N$, or either d or N is even, then,

$$p^*(N, d) \leq \begin{cases} \frac{1}{N} & \text{if } N \text{ is odd} \\ \frac{1}{N-1} & \text{if } N \text{ is even.} \end{cases}$$

Proof. It is enough to prove that

$$\min_{\lambda \in \text{Irr}(\mathbb{B}_N(d))} c(\lambda) - k(d-1) \leq \begin{cases} \frac{(1-d)(N-1)}{2} & \text{if } N \text{ is odd} \\ \frac{(1-d)N}{2} & \text{if } N \text{ is even,} \end{cases}$$

If $d > N$, the minimization can be restricted to only single-column partitions $\lambda := (1^{(N-2k)})$, for all $k \in \{0, \dots, \lfloor \frac{N}{2} \rfloor\}$, which is always possible when $d > N$. Let $|\lambda| := N - 2k$, then

$$\begin{aligned} \min_{\lambda \in \text{Irr}(\mathbb{B}_N(d))} c(\lambda) - k(d-1) &\leq \min_{k \in \{0, \dots, \lfloor \frac{N}{2} \rfloor\}} c(1^{(N-2k)}) - k(d-1) \\ &= \min_{k \in \{0, \dots, \lfloor \frac{N}{2} \rfloor\}} -\frac{|\lambda|(|\lambda| - 1)}{2} - (d-1)\frac{N - |\lambda|}{2} \\ &= \begin{cases} \frac{(1-d)(N-1)}{2} & \text{if } N \text{ is odd} \\ \frac{(1-d)N}{2} & \text{if } N \text{ is even.} \end{cases} \end{aligned}$$

Otherwise, if d is even, let $k^* = \lceil \frac{N}{2} \rceil - \frac{d}{2}$. Then the single-column partition $\lambda := (1^{(N-2(k+k^*))})$ satisfies $\lambda'_1 + \lambda'_2 \leq d$ for all $k \in \{0, \dots, \lfloor \frac{N}{2} \rfloor - k^*\}$, and,

$$\begin{aligned} \min_{\lambda \in \text{Irr}(\mathbb{B}_N(d))} c(\lambda) - k(d-1) &\leq \min_{k \in \{0, \dots, \lfloor \frac{N}{2} \rfloor - k^*\}} c(1^{(N-2k)}) - (k+k^*)(d-1) \\ &= \begin{cases} \frac{(1-d)(N-1)}{2} & \text{if } N \text{ is odd} \\ \frac{(1-d)N}{2} & \text{if } N \text{ is even.} \end{cases} \end{aligned}$$

The same result holds if N is even.

Theorem 4.5. *If $d > N$, or either d or N is even, then,*

$$p^*(N, d) = \begin{cases} \frac{1}{N} & \text{if } N \text{ is odd} \\ \frac{1}{N-1} & \text{if } N \text{ is even.} \end{cases}$$

Proof. Using [Section 4.3.1](#) and [Lemma 4.4](#), the dual optimization problem is lower and upper bounded by,

$$\begin{cases} \frac{1}{N} & \text{if } N \text{ is odd} \\ \frac{1}{N-1} & \text{if } N \text{ is even.} \end{cases}$$

Case $N \geq d$ and both d and N are odd

Let $\tilde{x} := \frac{2}{N(N-1)(1-d)}$, then the affine functions $f_{\lambda,\mu}$ of [Eq. \(4.4\)](#) evaluated at the negative coordinate \tilde{x} become,

$$\begin{aligned} f_{\lambda,\mu}(\tilde{x}) &= \frac{1}{d-1} \left(\frac{2d \cdot c(\mu)}{N(N-1)} - 1 \right) + \tilde{x} \left(c(\lambda) + d \cdot c(\mu) - k(d-1) - \frac{N(N-1)}{2} \right) \\ &= \frac{1}{n-1} + \tilde{x} \cdot h(\lambda), \end{aligned}$$

where $h(\lambda)$ is defined by $h(\lambda) := \frac{1}{2} \sum_{i=0}^{\lambda_1} \lambda'_i (d - \lambda'_i + 2(i-1))$. At this coordinate the affine functions $f_{\lambda,\mu}$ do not depend on μ anymore. Let g be the function defined by,

$$g(\lambda) := f_{\lambda,\mu}(\tilde{x}).$$

The offsets of the affine functions $f_{\lambda,\mu}$ do not depend on λ either. Let a be the function defined by,

$$a(\mu) := \frac{1}{d-1} \left(\frac{2d \cdot c(\mu)}{N(N-1)} - 1 \right).$$

If $N \geq d$, $k := \lfloor \frac{N-d}{2}/d \rfloor$ and $m := \frac{N-d}{2} \bmod d$, let the two partitions λ_1, λ_2 in $\text{Irr}(\mathbb{B}_N(d))$ and the three partitions μ_1, μ_2, μ_3 in $\text{Irr}(\mathfrak{S}_N)$ be defined by¹,

$$\begin{aligned} \lambda_1 &:= (1^d) & \mu_1 &:= (N) \\ \lambda_2 &:= (1) & \mu_2 &:= (N-d+1, 1^{d-1}) \\ & & \mu'_3 &:= (d^{2k+1}, m^2). \end{aligned} \tag{4.5}$$

Lemma 4.6. *If $N \geq d$ and both d and N are odd, then λ_1, μ_2 from [Eq. \(4.5\)](#) satisfy*

$$g(\lambda_1) - a(\mu_2) = \frac{2d+2-N}{N-1}.$$

1. In the definition above, μ_3 is given using the column notation μ'_3 . Using the row notation it becomes $\mu_3 = ((2k+3)^m, (2k+1)^{d-m})$: m rows of size $(2k+3)$ and $d-m$ rows of size $(2k+1)$.

In particular $g(\lambda_1) < a(\mu_2)$ if $N \geq 2d + 3$, and $g(\lambda_1) > a(\mu_2)$ if $N \leq 2d + 1$.

Proof. The content of μ_2 is $\frac{(N-d+1)(N-d)}{2} - \frac{d(d-1)}{2}$. Also $h(\lambda_1) = 0$, so $g(\lambda_1) = \frac{1}{N-1}$. Then

$$\begin{aligned} g(\lambda_1) - a(\mu_2) &= \frac{1}{N-1} - \frac{1}{d-1} \left(\frac{2d \cdot c(\mu_2)}{n(n-1)} - 1 \right) \\ &= -d \frac{(N-d+1)(N-d) - d(d-1)}{N(N-1)(d-1)} + \frac{1}{d-1} + \frac{1}{N-1} \\ &= \frac{2d+2-N}{N-1}. \end{aligned}$$

Lemma 4.7. If $N \geq d$ and both d and N are odd, then the partitions from Eq. (4.5) satisfy $(\lambda_i, \mu_j) \in \Gamma$, and the relations,

$$g(\lambda) < g(\lambda_2) < g(\lambda_1) \quad \text{and} \quad a(\mu) < a(\mu_1),$$

for all $\lambda \neq \lambda_1, \lambda_2$ in $\text{Irr}(\mathbb{B}_N(d))$ and all $\mu \neq \mu_1$ in $\text{Irr}(\mathfrak{S}_N)$. Moreover for all $(\lambda_1, \mu) \in \Omega$,

$$a(\mu) \leq a(\mu_2).$$

Proof. By definition of Γ , all (λ_i, μ_j) are in Γ .

Let λ in $\text{Irr}(\mathbb{B}_N(d))$ then

$$g(\lambda) = \frac{1}{N-1} + \tilde{x} \cdot h(\lambda).$$

But since $\lambda'_1 \leq d$ holds for all λ in $\text{Irr}(\mathbb{B}_N(d))$, then $h(\lambda) \geq 0$. In particular, $h(\lambda_2) = \frac{d-1}{2}$, and $h(\lambda) = 0$ iff $\lambda = \lambda_1$. Assume there exists λ in $\text{Irr}(\mathbb{B}_N(d))$ such that

$$g(\lambda_2) \leq g(\lambda) \leq g(\lambda_1),$$

then necessarily the first term of $h(\lambda)$ is either $h(\lambda_1)$ or $h(\lambda_2)$, since it is minimized for the columns (1) and (1^d) . But since all the terms in $h(\lambda)$ are positive, then either $g(\lambda) = g(\lambda_1)$ or $g(\lambda) = g(\lambda_2)$.

Because μ_1 is the N -box Young diagram that maximizes the content function, then

$$a(\mu) < a(\mu_1),$$

for all $\mu \neq \mu_1$ in $\text{Irr}(\mathfrak{S}_N)$.

Assume there exists μ such that $(\lambda_1, \mu) \in \Omega$ and,

$$a(\mu_2) \leq a(\mu) \leq a(\mu_1).$$

Since $((1^d), \mu) \in \Omega$ implies $((1^d), \mu) \in \Gamma$, then $(\lambda_1, \mu) \in \Gamma$, and by definition $r(\mu) = d$. Thus necessarily $c(\mu_2) \leq c(\mu)$, which implies that the first row of μ_2 is of size at most $n - d + 1$. But the content of a Young diagram is a non-increasing function of the first row's size, i.e. given two Young diagrams λ and μ with the same number of boxes, if $c(\lambda) \leq c(\mu)$ then $\lambda_1 \leq \mu_1$. Then μ_2 and μ share the same first row, and $\mu_2 = \mu$.

Theorem 4.8. *If $N \geq d$ and both d and N are odd, then,*

$$p^*(N, d) = \min \left\{ \frac{2d+1}{2dN+1}, \frac{1}{N-1} \right\}.$$

Proof. Let λ_1, λ_2 and μ_1, μ_2, μ_3 be the partitions from [Eq. \(4.5\)](#). Two cases are considered in the proof, $p^*(N, d) = g(\lambda_1)$ when $N \geq 2d + 3$, and $p^*(N, d)$ lies at intersection of the affine functions f_{λ_2, μ_1} and f_{λ_1, μ_2} , when $N \leq 2d + 1$ (see [Figure 10](#)).

Since

$$\begin{aligned} c(\mu_3) &= \sum_{i=1}^{2k+1} \left(-\frac{d(d-1)}{2} + (i-1)d \right) + \sum_{i=2k+2}^{2k+3} \left(-\frac{m(m-1)}{2} + (i-1)m \right) \\ &= \frac{d(2k+1)(2k+1-d)}{2} + m(4k+4-m), \end{aligned}$$

and $N - d = 2kd + 2m$ with $m \in \{0, \dots, d-1\}$, then

$$\begin{aligned} g(\lambda_1) - a(\mu_3) &= \frac{1}{N-1} - \frac{1}{d-1} \left(\frac{2d \cdot c(\mu_3)}{N(N-1)} - 1 \right) \\ &= \frac{N(d+N-2) - 2d \cdot c(\mu_3)}{N(N-1)(d-1)} \\ &= \frac{(d+2)(2m^2 - 2dm - N + dN)}{N(N-1)(d-1)} \\ &\geq \frac{(d+2)(-(d-1)(d+1)/2 + d(d-1))}{N(N-1)(d-1)} \end{aligned}$$

$$= \frac{(d+2)(d-1)}{2N(N-1)} > 0,$$

where in the first inequality $N \geq d$ was used, and that the minimum of $2m^2 - 2dm$ on the domain $m \in \{0, \dots, d-1\}$ is achieved for $m = \frac{d-1}{2}$. Geometrically this means that the point $(0, a(\mu_3))$ is always lower than $(\tilde{x}, g(\lambda_1))$.

Suppose that $N \geq 2d + 3$, then the relation

$$g(\lambda_1) < a(\mu_2),$$

holds by [Lemma 4.6](#). Therefore $p^*(N, d) \geq g(\lambda_1)$ since the optimal point should be above the intersection of the affine functions f_{λ_1, μ_2} and f_{λ_1, μ_3} that is, above $g(\lambda_1)$. But since $g(\lambda) \leq g(\lambda_1)$ for all λ in $\text{Irr}(\mathbb{B}_N(d))$ by [Lemma 4.7](#), it must be $p^*(N, d) = g(\lambda_1)$.

Suppose that $N \leq 2d + 1$, then the relation

$$a(\mu) \leq a(\mu_2),$$

holds for all $(\lambda_1, \mu) \in \Omega$, by [Lemma 4.7](#). Then $p^*(N, d)$ lies above the affine function f_{λ_1, μ_2} . But $g(\lambda) \leq g(\lambda_1)$ for all λ in $\text{Irr}(\mathbb{B}_N(d))$, by [Lemma 4.7](#), then $p^*(N, d)$ must lie on the affine function f_{λ_1, μ_2} , at the intersection with another affine function $f_{\lambda, \mu}$ with $g(\lambda) \leq a(\mu)$. Among all such affine functions there are no functions with $\lambda = \lambda_1$ due to [Lemma 4.7](#). Because $g(\lambda) \leq g(\lambda_2)$ for all $\lambda \neq \lambda_1$ in $\text{Irr}(\mathbb{B}_N(d))$ and $a(\mu) \leq a(\mu_1)$ for all $\mu \in \text{Irr}(\mathfrak{S}_N)$, by [Lemma 4.7](#), it must be that this function is f_{λ_2, μ_1} . Therefore $p^*(N, d)$ lies at intersection of the affine functions f_{λ_2, μ_1} and f_{λ_1, μ_2} . In order to find the intersection of f_{λ_2, μ_1} and f_{λ_1, μ_2} , the equation $f_{\lambda_2, \mu_1}(x^*) = f_{\lambda_1, \mu_2}(x^*)$ must be solved, which gives $x^* = \frac{4d}{(d-1)(N-1)(2dN+1)}$ and $p^*(N, d) = \frac{2d+1}{2dN+1}$.

In conclusion, when $N \geq 2d + 3$ then $p^*(N, d) = \frac{1}{N-1}$, and when $N \leq 2d + 1$ then $p^*(N, d) = \frac{2d+1}{2dN+1}$, which is equivalent to the statement of the theorem.

4.3.4 Optimal value

Using [Theorem 4.5](#) and [Theorem 4.8](#), the dual optimization problem [Eq. \(4.4\)](#) has solution,

$$p^*(N, d) = \begin{cases} \frac{1}{N+N \bmod 2-1} & \text{if } d > N \text{ or either } d \text{ or } N \text{ is even} \\ \min \left\{ \frac{2d+1}{2dN+1}, \frac{1}{N-1} \right\} & \text{if } N \geq d \text{ and both } d \text{ and } N \text{ are odd.} \end{cases}$$

Hence $p^*(N, d)$ is decreasing with respect to N , but is not monotonic with respect to d .

Slater's condition for strong duality of SDP holds true in this case, from [Section 4.3.1](#), hence both optimal values of the optimization problem [Eq. \(4.1\)](#) and the dual optimization problem [Eq. \(4.4\)](#) are equal, i.e. $p(N, d) = p^*(N, d)$. The first values of $p(N, d)$ are summarized in the following table (in \square the values of $p(N, d)$ for which the isotropic states ρ_e are separable):

$d \backslash N$	2	3	4	5	6	7	8	9
2	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{7}$	$\frac{1}{7}$	$\frac{1}{9}$
3	1	$\frac{7}{19}$	$\frac{1}{3}$	$\frac{7}{31}$	$\frac{1}{5}$	$\frac{7}{43}$	$\frac{1}{7}$	$\frac{1}{8}$
4	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{7}$	$\frac{1}{7}$	$\frac{1}{9}$
5	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{11}{51}$	$\frac{1}{5}$	$\frac{11}{71}$	$\frac{1}{7}$	$\frac{11}{91}$
6	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{7}$	$\frac{1}{7}$	$\frac{1}{9}$
7	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{5}{33}$	$\frac{1}{7}$	$\frac{15}{127}$
8	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{7}$	$\frac{1}{7}$	$\frac{1}{9}$
9	1	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{7}$	$\frac{1}{7}$	$\frac{19}{163}$

In general, it is not known which quantum state ρ gives the optimal value $p(N, d)$, but using the commutation relation [Eq. \(4.3\)](#), it must be in the algebra,

$$\text{Span}_{\mathbb{C}} \{ \psi(\sigma) \mid \sigma \in \mathbb{B}_N \}.$$

For example when $N = 3$ and $d = 3$ the optimal quantum state ρ for the optimisation problem [Eq. \(4.1\)](#) is:

$$\rho = \frac{1}{57} \left[\begin{array}{c} \begin{array}{c} \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} \\ \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} \end{array} \\ + 2 \left(\begin{array}{c} \text{⌞⌞⌞} + \text{⌞⌞⌞} + \text{⌞⌞⌞} - \text{⌞⌞⌞} - \text{⌞⌞⌞} - \text{⌞⌞⌞} \end{array} \right) \end{array} \right].$$




Figure 10 – The optimal value of the dual optimization problem lies at the intersection of the set of affine functions $f_{\lambda, \mu}(x)$. When $N \geq 2d + 3$ (Figure 10a, with $N = 9$ and $d = 3$), then $p^*(N, d) = g(\lambda_1)$. When $N \leq 2d + 1$ (Figure 10b, with $N = 5$ and $d = 3$), then $p^*(N, d)$ lies at the intersection of f_{λ_2, μ_1} and f_{λ_1, μ_2} . In ■ the max over all affine functions, in ■ the affine functions from the set Γ , and in ■ the affine functions from the set $\Omega \setminus \Gamma$.

Appendix A

Elements of representation theory

Appendix contents	
A.1	Finite groups 107
A.1.1	Representations of finite groups 107
A.1.2	Reducibility 108
A.1.3	Character theory 112
A.1.4	Representations of \mathfrak{S}_n 120
A.1.5	Restricted and induced representations 134
A.2	Matrix groups 140
A.2.1	Representations of connected compact matrix groups 140
A.2.2	Representations of U_d 147

This Appendix provides a comprehensive overview of the fundamental principles of representation theory that are used throughout the present thesis. Its aim is to focus primarily on the investigation of representations of finite groups, with a specific emphasis on the symmetric group, and on those of infinite matrix groups, with a specific emphasis on the unitary group.

The sections devoted to finite groups provide a complete set of proofs, while more detailed treatments of the subject can be found in the books [FH13; Ser+77; Sag13; Jam06]. Regarding matrix groups, only the principal results are presented, with books [GW09; HH13; BT13; Bak03; Bum+04; Ada82; Sim96] serving as references for the corresponding proofs. A thorough investigation of the representation theory with a particular emphasis on its application in the domain of physics, may be found in the books [Lan12; Boe+70; Zhe73; Ste95; Tun85].

An alternative approach to address the representations of the symmetric group, which has been proposed by A. Okounkov and A. Vershik, can be found in [VO05; CST10].

A.1 Finite groups

A.1.1 Representations of finite groups

Let G be a finite group, a **representation** of G is a pair (ρ, V) , where V is a complex vector space with dimension d and $\rho : G \rightarrow \text{GL}(V)$ is a group homomorphism, i.e. $\rho(g \cdot h) = \rho(g)\rho(h)$ holds for all g and h in G . Specifically, ρ satisfies the condition $\rho(1_G) = I$, where I is the identity matrix on V . As a corollary of this property, it follows that $\rho(g^{-1}) = \rho(g)^{-1}$ for every $g \in G$.

Consider two representations (ρ, V) and (σ, W) of G , an **intertwining map** between these representations is a linear map $\phi : V \rightarrow W$ that satisfies $\phi \circ \rho(g) = \sigma(g) \circ \phi$ for all $g \in G$. If such a map is an isomorphism, then the representations are said to be **equivalent**, and $\rho(g) = \phi^{-1} \circ \sigma(g) \circ \phi$ for all $g \in G$. The set of all intertwining maps between (ρ, V) and (σ, W) is denoted $\text{Hom}_G(V, W)$, and forms a complex vector space.

If (ρ_V, V) is a representation of G , then the **dual representation** (ρ_{V^*}, V^*) is defined for all $g \in G$ by

$$\rho_{V^*}(g) := \rho_V(g^{-1})^T,$$

where \cdot^T denotes the transpose.

The natural bilinear pairing $\langle \cdot, \cdot \rangle$ between complex vector spaces V and V^* is defined as $\langle f, x \rangle := f(x)$ for all $x \in V$ and $f \in V^*$. Then, the dual representation

(ρ_{V^*}, V^*) preserves this natural bilinear pairing, specifically, $\langle \rho_{V^*}(g)f, \rho_V(g)x \rangle = \langle f, x \rangle$ holds for any $g \in G$ and for all $x \in V$ and $f \in V^*$.

Example. Consider a finite group G of order n , and let (ρ, V) be a representation of G . Let g be an arbitrary element of G , and let $A \in \text{GL}(V)$ be the image of g under ρ , i.e. $\rho(g) = A$. It follows that

$$A^n = I.$$

The characteristic polynomial $X^n - 1$ of A factors into n distinct eigenvalues, which are the n -th roots of unity. As a result, it can be deduced that A is diagonalizable. Furthermore, if G is an abelian group, all of the matrices $\rho(g)$ are simultaneously diagonalizable. In [Section A.1.2](#), the concept of block diagonalization will be explored in the context of non-abelian groups.

Given a finite group G and a representation (ρ, V) of G , the complex vector space V is said to carry the representation ρ of G . For clarity, it is appropriate to succinctly refer to the complex vector space V as a representation of G when no ambiguity surrounding the mapping ρ exists. For any $g \in G$ and $v \in V$, the action of g on v can be denoted by $g \cdot v$, as a compact alternative to the expression $\rho(g)(v)$.

Remark. The representation of a finite group G is a mapping of the group onto a set of matrices that operate on a complex vector space V . This mapping preserves the underlying structure of the group, such that the group law of G is equivalent to matrix multiplication on V . As a result, the study of representations of G can be approached using the mathematical framework of linear algebra, rather than through direct examination of the group itself.

A.1.2 Reducibility

Let V be a representation of a finite group G , and let W be a subspace of V . If W is invariant under G , that is to say, for all $g \in G$ and $w \in W$, the element $g \cdot w$ belongs to W , then W is called a **subrepresentation**. If V possesses exactly two distinct subrepresentations, namely the trivial zero subspace and V itself, then the representation is said to be **irreducible**, otherwise the representation is considered to be **reducible**.

Theorem A.1 (Maschke). *Let V be a representation of a finite group G , and suppose that W is a subrepresentation of V . Then, there exists a complementary subrepresentation W^\perp of V , such that V is the direct sum of W and W^\perp , i.e. $V = W \oplus W^\perp$.*

Proof. Let $\langle \cdot, \cdot \rangle$ be any Hermitian inner product on the complex vector space V . The G -invariant Hermitian inner product $\langle \cdot, \cdot \rangle_G$ on V is defined by

$$\langle v, w \rangle_G := \frac{1}{|G|} \sum_{g \in G} \langle g \cdot v, g \cdot w \rangle.$$

Then $\langle g \cdot v, g \cdot w \rangle_G = \langle v, w \rangle_G$ holds for any $g \in G$.

Let $W^\perp = \{v \in V \mid \langle v, w \rangle_G = 0 \text{ for all } w \in W\}$ be the orthogonal complement of W with respect to the inner product $\langle \cdot, \cdot \rangle_G$. It follows that $V = W \oplus W^\perp$. Furthermore, for any $g \in G$ and for all $v \in W^\perp$ and $w \in W$, the G -invariance of $\langle \cdot, \cdot \rangle_G$ implies

$$\langle g \cdot v, w \rangle_G = \langle v, g^{-1} \cdot w \rangle_G = 0,$$

as $g^{-1} \cdot w \in W$. Therefore $g \cdot v \in W^\perp$ for all $v \in W^\perp$ and $g \in G$, and thus W^\perp is a subrepresentation.

In accordance with Maschke's [Theorem A.1](#), it can be established that if a finite group G possesses a reducible representation, it must necessarily be decomposed into the direct sum of at least two subrepresentations. These subrepresentations may in turn be further decomposed into further direct sums, if they are themselves reducible. This process of decomposition may be repeated until the representation is fully decomposed into the direct sum of irreducible representations of G . This result holds true for any finite group and is referred to as the concept of **complete reducibility** of finite groups.

Remark. Maschke's [Theorem A.1](#) asserts that for any finite group G and its representation (ρ, V) , the matrices $\rho(g)$ for all $g \in G$ are simultaneously block-diagonalizable. Given the decomposition of the underlying complex vector space V into $V = W \oplus W^\perp$, where W is a subrepresentation of V , it follows that for each $g \in G$, the matrix $\rho(g)$ block-diagonalizes as

$$\rho(g) = \begin{pmatrix} \rho_W(g) & 0 \\ 0 & \rho_{W^\perp}(g) \end{pmatrix}.$$

Here, $\rho_W(g)$ and $\rho_{W^\perp}(g)$ are the restrictions of $\rho(g)$ onto W and W^\perp , respectively. This result highlights the crucial property of simultaneously block-diagonalizability of matrices in representation theory.

Maschke's [Theorem A.1](#) plays a crucial role in the field of representation theory of finite groups. The theorem states that every representation of a finite group can be decomposed into a direct sum of irreducible representations. This implies that to comprehend any representation of a finite group, it is sufficient to have a complete understanding of its irreducible representations. This result leads to the fundamental question of determining the number of irreducible representations for a given finite group, which is addressed in [Section A.1.3](#).

Lemma A.2 (Schur). *If V and W are two irreducible representations of a finite group G , and $\phi \in \text{Hom}_G(V, W)$ is a nonzero intertwining map, then ϕ is an isomorphism.*

Proof. Let $v \in \ker \phi$, then for all $g \in G$, since ϕ is an intertwining map it follows that

$$g \cdot \phi(v) = \phi(g \cdot v) = 0.$$

Thus $\ker \phi$ is a subrepresentation of V , and since V is irreducible and ϕ is nonzero, necessarily $\ker \phi = \{0\}$. Therefore ϕ is injective.

Conversely, let $v \in V$, then for all $g \in G$, since ϕ is an intertwining map it follows that

$$g \cdot \phi(v) = \phi(g \cdot v) \in \text{Im } \phi.$$

Thus $\text{Im } \phi$ is a subrepresentation of W , and since W is irreducible and ϕ is nonzero, necessarily $\text{Im } \phi = W$. Hence ϕ is surjective.

Schur's [Lemma A.2](#) is a foundational principle in the field of representation theory. It has two particularly noteworthy implications for a finite group G .

Firstly, if V is an irreducible representation of a G , and ϕ belongs to $\text{Hom}_G(V)$, then ϕ is a homothety, meaning that ϕ is proportional to the identity matrix, with a scalar factor $\lambda \in \mathbb{C}$, i.e.

$$\phi = \lambda \cdot I.$$

Secondly, if V is a representation of G , there exists a unique decomposition of V into a direct sum of non-equivalent irreducible representations, expressed as

$$V \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k},$$

where n_i is referred to as the **multiplicity** of the irreducible representation V_i .

Remark. In the decomposition of complex vector space V as $V \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, both the subspaces V_i and the multiplicities n_i are unique. However, it must be noted that the direct sum decomposition of each $V_i^{\oplus n_i}$ into n_i copies of

■ V_i is not guaranteed to be unique in general.

For any finite group G , there is a unique, up to isomorphism, representation referred to as the **trivial representation**, which is invariant under the action of all elements of G . This representation is characterized by the property that all elements of G act as the identity. Furthermore, this representation is one-dimensional, and thus irreducible.

Let V be a representation of G . The invariant subspace of V with respect to G , denoted as V_G , can be defined as follows:

$$V_G := \{v \in V \mid g \cdot v = v, \forall g \in G\}.$$

Then V_G can be decomposed into a direct sum of irreducible representations. In particular, each summand of this direct sum is isomorphic to the trivial representation.

Proposition A.3. *Let V be a representation of a finite group G . Consider the map $\phi : V \rightarrow V$ defined as follows: for any $v \in V$,*

$$\phi(v) := \frac{1}{|G|} \sum_{g \in G} g \cdot v.$$

Then ϕ is a projector onto V_G .

Proof. Let $v \in V_G$, which implies that $\phi(v) = v$ according to the definition of V_G . As a result, $V_G \subset \text{Im } \phi$. For any $v \in V$, the action of any element $g \in G$ on $\phi(v)$ results in $g \cdot \phi(v) = \phi(v)$, thus $\text{Im } \phi \subset V_G$. Furthermore, for all $v \in V$,

$$\begin{aligned} \phi \circ \phi(v) &= \frac{1}{|G|^2} \sum_{g \in G} \sum_{h \in G} (g \cdot h) \cdot v \\ &= \frac{|G|}{|G|^2} \sum_{g \in G} g \cdot v \\ &= \phi(v), \end{aligned}$$

which implies that $\phi \circ \phi = \phi$.

A representation V of a finite group G is considered to be **unitary** if there exists a Hermitian inner product $\langle \cdot, \cdot \rangle$ defined on V such that the property of unitarity is satisfied, that is, for all $g \in G$, it holds that $\langle g \cdot u, g \cdot v \rangle = \langle u, v \rangle$. Given any Hermitian inner product $\langle \cdot, \cdot \rangle$ in V , a new Hermitian inner product $\langle \cdot, \cdot \rangle_G$ can be

defined on V as follows:

$$\langle u, v \rangle_G := \frac{1}{|G|} \sum_{g \in G} \langle g \cdot u, g \cdot v \rangle.$$

Then for all $g \in G$, it holds that $\langle g \cdot u, g \cdot v \rangle_G = \langle u, v \rangle_G$. This means that every representation of a finite group can be considered to be unitary with respect to $\langle \cdot, \cdot \rangle_G$.

Example. Let \mathfrak{S}_3 be the symmetric group over 3 elements, and V be the complex vector space \mathbb{C}^3 with basis e_1, e_2 and e_3 . The **natural representation** of \mathfrak{S}_3 associates each element σ in \mathfrak{S}_3 with its permutation matrix P_σ . This matrix represents the permutation of the 3 coordinates of V according to σ . Then the invariant subspace $V_{\mathfrak{S}_3}$ is the one-dimensional subspace:

$$\text{Span}_{\mathbb{C}}(1, 1, 1).$$

Let V be an irreducible representation of a finite group G , and consider any nonzero vector v in V . It follows that v generates V under the action of G , that is, $G \cdot v = V$. Otherwise, if the orbit $G \cdot v$ were a proper subspace of V , then by the definition of irreducibility, $G \cdot v$ would be an invariant subspace of V , contradicting the assumption that V is irreducible.

A.1.3 Character theory

Let (ρ, V) be a representation of a finite group G . The **character** of (ρ, V) is the map $\chi_V : G \rightarrow \mathbb{C}$ defined on $g \in G$ as follows:

$$\chi_V(g) := \text{Tr} [\rho(g)].$$

A **class function** on G is complex function f that remains constant on the conjugacy classes of G , i.e. $f(h \cdot g \cdot h^{-1}) = f(g)$, for all $g, h \in G$. The collection of all class functions on G forms a complex vector space with dimension equal to the number of conjugacy classes of G . Due to the cyclic property of the trace, the characters are class functions. The concept of characters plays a central role in the representation theory of finite groups. It provides a means of calculating important quantities, such as the dimension of a representation. For instance, let V be a representation of G , and let ϕ be the projector onto V_G , defined in [Proposition A.3](#), then $\chi_V(1_G) = \dim V$ and $\chi_V(\phi) = \dim V_G$, where $\chi_V(\phi)$ is defined linearly on each summand of ϕ .

Proposition A.4. Consider two representations, (ρ_V, V) and (ρ_W, W) , of a finite group G . Then

$$\chi_{V \oplus W} = \chi_V + \chi_W, \quad \chi_{V \otimes W} = \chi_V \cdot \chi_W, \quad \text{and} \quad \chi_{V^*} = \bar{\chi}_V,$$

where $\bar{\cdot}$ denotes the complex conjugate.

Proof. Let g be in G . By the properties of the trace,

$$\begin{aligned} \chi_{V \oplus W}(g) &= \text{Tr} [\rho_V(g)] + \text{Tr} [\rho_W(g)] \\ \chi_{V \otimes W}(g) &= \text{Tr} [\rho_V(g)] \cdot \text{Tr} [\rho_W(g)]. \end{aligned}$$

Then $\chi_V(g) = \text{Tr} [\rho_V(g)]$ is the sum of the eigenvalues of the diagonalizable $\rho_V(g)$. Furthermore, since the eigenvalues, λ_i , of $\rho_V(g)$ are roots of unity, it follows that the inverse of each eigenvalue is equal to its complex conjugate, i.e. $\lambda_i^{-1} = \bar{\lambda}_i$. Finally

$$\chi_{V^*}(g) = \text{Tr} [\rho_V(g^{-1})] = \bar{\chi}_V(g).$$

Corollary. Let (ρ_V, V) be a representations of a finite group G , then $\bar{\chi}_V(g) = \chi_V(g^{-1})$, for all $g \in G$.

From the character of a tensor product and a dual space, the isomorphism $\text{Hom}(V, W) \simeq V^* \otimes W$ between two finite dimensional complex vector spaces V and W , implies

$$\chi_{\text{Hom}(V, W)} = \bar{\chi}_V \cdot \chi_W.$$

Consider a finite group G and the set of class functions defined on it. Let the Hermitian inner product $\langle \cdot, \cdot \rangle_G$ on this set of class functions be defined as follows: given two class functions f and g on G , then,

$$\langle f, g \rangle_G := \frac{1}{|G|} \sum_{g \in G} \bar{f}(g) \cdot h(g).$$

Theorem A.5. Given a finite group G , the characters of its irreducible representations serve as an orthonormal basis for the space of class functions of G , with respect to the Hermitian inner product $\langle \cdot, \cdot \rangle_G$.

Proof. Let V and W be two irreducible representations of G . By Schur's Lemma A.2, the dimension of $\text{Hom}_G(V, W)$ is either 1 if $V \simeq W$ and 0 otherwise. But the dimension of $\text{Hom}_G(V, W)$ is also equal to the character of the projector onto $(V^* \otimes W)_G$. That is

$$\frac{1}{|G|} \sum_{g \in G} \bar{\chi}_V(g) \cdot \chi_W(g) = \begin{cases} 1 & \text{if } V \simeq W \\ 0 & \text{otherwise.} \end{cases}$$

Thus the characters of the irreducible representations of G are orthonormal with respect to the Hermitian inner product $\langle \cdot, \cdot \rangle_G$.

Hence, the number of irreducible representations of G is finite, and in fact smaller than the number of conjugacy classes of G . Let (ρ_{V_i}, V_i) denotes the irreducible representations of G , and χ_{V_i} the corresponding characters. Define

$$V := \text{Span}_{\mathbb{C}} \{ \chi_{V_i} \},$$

the span of these characters over the field of complex numbers.

Let W be the complex vector space of all functions from G to \mathbb{C} . The basis of this complex vector space are the elements δ_g , where $g \in G$, defined on all $h \in G$ by

$$\delta_g(h) = \begin{cases} 1 & \text{if } g = h \\ 0 & \text{otherwise.} \end{cases}$$

Let $g \in G$ and define the representation of G on W by $(g \cdot f)(h) = f(g^{-1} \cdot h)$, for all $f \in W$ and $h \in G$.

Let $f \in V^\perp$ be a class function in the orthogonal complement of V , for all irreducible representations V_i of G , define the linear map $\phi_i : V_i \rightarrow V_i$ by

$$\phi_i := \frac{1}{|G|} \sum_{g \in G} \bar{f}(g) \cdot \rho_{V_i}(g).$$

Let $h \in G$, then since f is a class function it holds that

$$\begin{aligned} \rho_{V_i}(h^{-1}) \cdot \phi_i \cdot \rho_{V_i}(h) &= \frac{1}{|G|} \sum_{g \in G} \bar{f}(g) \cdot \rho_{V_i}(h^{-1} \cdot g \cdot h) \\ &= \frac{1}{|G|} \sum_{g \in G} \bar{f}(h \cdot g \cdot h^{-1}) \cdot \rho_{V_i}(g) \\ &= \phi_i, \end{aligned}$$

and in particular $\phi_i \in \text{Hom}_G(V_i)$. From Schur's [Lemma A.2](#) there exists $\lambda \in \mathbb{C}$ such that $\phi_i = \lambda \cdot I$, with the equality $\text{Tr} [\phi_i] = \lambda \cdot \dim V_i$, that is

$$\begin{aligned} \lambda &= \frac{1}{|G| \cdot \dim V_i} \sum_{g \in G} \bar{f}(g) \cdot \text{Tr} [\rho_{V_i}(g)] \\ &= \frac{1}{|G| \cdot \dim V_i} \sum_{g \in G} \bar{f}(g) \cdot \chi_{V_i}(g) \\ &= \frac{1}{\dim V_i} \langle f, \chi_{V_i} \rangle_G. \end{aligned}$$

Thus $\lambda = 0$, since $f \in V^\perp$. In particular, from the decomposition into a direct sum of irreducible representations $W \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, it holds that

$$\begin{aligned} \frac{1}{|G|} \sum_{g \in G} \bar{f}(g) \cdot (g \cdot \delta_{1_G}) &= \frac{1}{|G|} \sum_{g \in G} \bar{f}(g^{-1}) \cdot \delta_g \\ &= 0. \end{aligned}$$

But since $\{\delta_g \mid g \in G\}$ forms a basis of W , necessarily $f(g) = 0$ for all $g \in G$. Thus $V^\perp = \{0\}$.

Corollary. *Let V be a representation of a finite group G . The representation V is irreducible if and only if $\langle \chi_V, \chi_V \rangle_G = 1$. Otherwise, V decomposes into a direct sum of irreducible representations $V \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, with*

$$\langle \chi_V, \chi_V \rangle_G = \sum_{i=1}^k n_i^2 \quad \text{and} \quad \langle \chi_{V_i}, \chi_V \rangle_G = n_i.$$

In light of [Theorem A.5](#), it has been established that the cardinality of the set of irreducible representations of a finite group is equal to the cardinality of the set of conjugacy classes of this group. The orthogonality of characters of irreducible representations of the finite group G serves as the foundation for constructing the **character table** of G , which assigns to each irreducible representation of G a unique collection of numbers, namely the characters of each conjugacy class of G . Additionally, given a representation V of a finite group, with its decomposition into a direct sum of irreducible representations $V \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, the character χ_i can be used to determine the multiplicity n_i , i.e., the number of times a particular irreducible representation V_i appears in the representation V .

Despite [Theorem A.5](#), in general for a finite group G , there is no known correspondence between the conjugacy classes of G and the irreducible representations of G . However, in [Section A.1.4](#), this correspondence will be explicitly demonstrated for the specific case of the symmetric group \mathfrak{S}_n .

Consider a finite group G . The **group algebra** of G , denoted by $\mathbb{C}[G]$, is the complex vector space with the elements of G as its basis. The multiplication in $\mathbb{C}[G]$ is defined as the group law of G on the basis. The **regular representation** of G is obtained by considering $\mathbb{C}[G]$ as a representation, where for all $g \in G$ and $x = \sum_{h \in G} c_h \cdot h$ an element of $\mathbb{C}[G]$, the action of g on x becomes,

$$g \cdot x = \sum_{h \in G} c_h \cdot (g \cdot h) = \sum_{h \in G} c_{(g^{-1} \cdot h)} \cdot h.$$

Remark. In this formulation, the group G serves as both the complex vector space of the representation as the group algebra, and the group homomorphism through its group law.

Consider a finite group G . For any element $g \in G$, the action of g on G through the group law results in a bijective mapping. It follows that the regular representation of g has no fixed points except in the case where g is the identity element 1_G of G . Additionally, all elements of G in the regular representation are represented as permutation matrices.

Proposition A.6. *Let $\mathbb{C}[G]$ be the group algebra of a finite group G , and consider $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, its decomposition into a direct sum of irreducible representations V_i with multiplicities n_i . Then for each irreducible representation V_i , its multiplicity in $\mathbb{C}[G]$ is equal the dimension $\dim V_i$.*

Proof. The character of $g \in G$ for the regular representation is equal to the cardinality of the set of fixed points of the action of g on G , thus

$$\chi_{\mathbb{C}[G]}(g) = \begin{cases} |G| & \text{if } g = 1_G \\ 0 & \text{otherwise.} \end{cases}$$

Consider an irreducible representation V_i , which appears in the decomposition of the group algebra $\mathbb{C}[G]$ as follows: $\mathbb{C}[G] \cong V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, then

$$\begin{aligned} n_i &= \langle \chi_{V_i}, \chi_{\mathbb{C}[G]} \rangle_G \\ &= \frac{1}{|G|} \sum_{g \in G} \bar{\chi}_{V_i}(g) \cdot \chi_{\mathbb{C}[G]}(g) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{|G|} \bar{\chi}_{V_i}(1_G) \cdot \chi_{\mathbb{C}[G]}(1_G) \\
&= \dim V_i.
\end{aligned}$$

Corollary. Consider the group algebra $\mathbb{C}[G]$ of a finite group G , and its decomposition into a direct sum of irreducible representations $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$. Then, it follows that:

$$\dim \mathbb{C}[G] = \sum_i \dim(V_i)^2.$$

The decomposition of the group algebra $\mathbb{C}[G]$ of a finite group G , into a direct sum of irreducible representations is a powerful tool for understanding the structure of finite groups, and the equation

$$\mathbb{C}[G] \simeq V_1^{\oplus \dim V_1} \oplus \cdots \oplus V_k^{\oplus \dim V_k},$$

provides important information about the size and multiplicity of irreducible representations in the regular representation.

Theorem. Let $\mathbb{C}[G]$ denote the group algebra of a finite group G , and let $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$ be the decomposition of $\mathbb{C}[G]$ into a direct sum of irreducible representations. For each irreducible representation W of G , there exists an index $i \in \{1, \dots, k\}$ such that W is equivalent to V_i .

Proof. Let U, V and W be three complex vector spaces of finite dimension, and define the inclusion and projection maps

$$V \xrightleftharpoons[\iota_V]{\pi_V} V \oplus W \xrightleftharpoons[\iota_W]{\pi_W} W,$$

on all $v \in V$ and $w \in W$ by

$$\begin{aligned}
\pi_V : (v, w) &\longmapsto v & \iota_V : v &\longmapsto v + 0 \\
\pi_W : (v, w) &\longmapsto w & \iota_W : w &\longmapsto 0 + w.
\end{aligned}$$

Now let G be a finite group, and assume that U, V and W are representations of G . Using the previous inclusion and projection maps, the following isomorphisms holds:

$$\begin{aligned}
\operatorname{Hom}(U, V \oplus W) &\simeq \operatorname{Hom}(U, V) \oplus \operatorname{Hom}(U, W) \\
\operatorname{Hom}_G(U, V \oplus W) &\simeq \operatorname{Hom}_G(U, V) \oplus \operatorname{Hom}_G(U, W).
\end{aligned}$$

Then, in the decomposition of $\mathbb{C}[G]$ into a direct sum of irreducible representations,

$$\mathrm{Hom}_G(\mathbb{C}[G], W) \simeq \bigoplus_{i=1}^k \mathrm{Hom}_G(V_i, W)^{\oplus n_i}.$$

In particular, the equality of dimensions

$$\dim \mathrm{Hom}_G(\mathbb{C}[G], W) = \bigoplus_{i=1}^k n_i \cdot \dim \mathrm{Hom}_G(V_i, W),$$

holds, and from the Schur's [Lemma A.2](#), $\dim \mathrm{Hom}_G(V_i, W) = 1$ if and only if $V_i \simeq W$, otherwise $\dim \mathrm{Hom}_G(V_i, W) = 0$. It follows that the dimension of $\mathrm{Hom}_G(\mathbb{C}[G], W)$ is positive if and only if there exists an index $i \in \{1, \dots, k\}$ such that the irreducible representation W is equivalent to V_i .

Let $f : \mathrm{Hom}_G(\mathbb{C}[G], W) \rightarrow W$ be the linear map defined on h in $\mathrm{Hom}_G(\mathbb{C}[G], W)$ by $f(h) = h(1_G)$, and let such h in $\mathrm{Hom}_G(\mathbb{C}[G], W)$ satisfying $f(h) = 0$, then for all $g \in G$,

$$h(g) = h(g \cdot e) = g \cdot h(e) = 0,$$

thus f is injective. Let $x \in W$ and consider the function $h : G \rightarrow W$ defined on $g \in G$ by $h(g) = g \cdot x$, thus h can be extended linearly on $\mathbb{C}[G]$, and in particular $h \in \mathrm{Hom}_G(\mathbb{C}[G], W)$, but since $f(h) = x$, then f is surjective. Therefore f is an isomorphism and the equality of dimensions

$$\dim \mathrm{Hom}_G(\mathbb{C}[G], W) = \dim W,$$

holds. In particular $\dim \mathrm{Hom}_G(\mathbb{C}[G], W)$ is positive.

The group algebra $\mathbb{C}[G]$, of a finite group G , provides a unified way to study the representations of G by encapsulating all the irreducible representations within its structure.

Example. As established in [Theorem A.5](#), the number of irreducible representations of \mathfrak{S}_3 is equal to the number of its conjugacy classes, in the present case 3. These three irreducible representations consist of the trivial representation, the **sign representation**, and the **standard representation**. The sign representation is the one-dimensional representation on which any $\sigma \in \mathfrak{S}_3$ is sent to $\mathrm{sign}(\sigma)$, where $\mathrm{sign}(\sigma)$ is the **signature** of the permutation σ . The standard

representation is the two-dimensional orthogonal complement of $\text{Span}_{\mathbb{C}}(1, 1, 1)$ in the natural representation. Then

$$\mathbb{C}[\mathfrak{S}_3] \simeq V_{\text{trivial}} \oplus V_{\text{sign}} \oplus V_{\text{standard}}^2.$$

The character table of \mathfrak{S}_3 is defined as the table with the irreducible representations of \mathfrak{S}_3 as its rows, and the conjugacy classes of \mathfrak{S}_3 as its columns. The entries in the table are the character values for the corresponding irreducible representations and conjugacy classes of \mathfrak{S}_3 . The conjugacy classes of \mathfrak{S}_3 are $\{1_{\mathfrak{S}_3}\}$, $\{(1\ 2), (1\ 3), (2\ 3)\}$ and $\{(1\ 2\ 3), (3\ 2\ 1)\}$. The character table of \mathfrak{S}_3 becomes:

\mathfrak{S}_3	$\{1_{\mathfrak{S}_3}\}$	$\{(1\ 2), (1\ 3), (2\ 3)\}$	$\{(1\ 2\ 3), (3\ 2\ 1)\}$
χ_{trivial}	1	1	1
χ_{sign}	1	-1	1
χ_{standard}	2	0	-1

Note that as expected from [Theorem A.5](#), the rows are orthogonal, with respect to Hermitian inner product $\langle \cdot, \cdot \rangle_G$.

Similarly to [Proposition A.3](#), given a finite group G , it is possible to define a projector onto each direct sum of equivalent irreducible representations $V_i^{\oplus n_i}$ that appears in the decomposition of its group algebra, $\mathbb{C}[G]$, into a direct sum of irreducible representations, $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$. The direct sum of equivalent irreducible representations $V_i^{\oplus n_i}$, in the decomposition of its group algebra $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, are called the **isotypic components**.

Theorem A.7. *Given a finite group G and its group algebra $\mathbb{C}[G]$ decomposed into a direct sum of irreducible representations $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, for each irreducible representation V_i , let the map ϕ_i , defined on $\mathbb{C}[G]$ by*

$$\phi_i := \frac{\dim V_i}{|G|} \sum_{g \in G} \bar{\chi}_{V_i}(g) \cdot g.$$

Then ϕ_i is a projector onto $V_i^{\oplus n_i}$.

Proof. Let $h \in G$, then since χ_{V_i} is a class function it holds that

$$h^{-1} \cdot \phi_i \cdot h = \frac{\dim V_i}{|G|} \sum_{g \in G} \bar{\chi}_{V_i}(g) \cdot (h^{-1} \cdot g \cdot h)$$

$$\begin{aligned}
&= \frac{\dim V_i}{|G|} \sum_{g \in G} \bar{\chi}_{V_i}(h \cdot g \cdot h^{-1}) \cdot g \\
&= \phi_i,
\end{aligned}$$

and in particular $\phi_i \in \text{Hom}_G(\mathbb{C}[G])$. Consider the decomposition of the group algebra $\mathbb{C}[G]$ into a direct sum of irreducible representations $\mathbb{C}[G] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$, for all $j \in \{1, \dots, k\}$, from Schur's [Lemma A.2](#), the restriction of ϕ_i to V_j is an homothety $\lambda \cdot I$, with $\lambda \in \mathbb{C}$, such that from [Theorem A.5](#)

$$\begin{aligned}
\lambda &= \frac{\dim V_i}{|G| \cdot \dim V_j} \sum_{g \in G} \bar{\chi}_{V_i}(g) \cdot \chi_{V_j}(g) \\
&= \frac{\dim V_i}{\dim V_j} \langle \chi_{V_i}, \chi_{V_j} \rangle_G \\
&= \begin{cases} 1 & \text{if } V_i \simeq V_j \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

Thus, the restriction of ϕ_i to V_i is the identity if V_j is equivalent to V_i , and the zero map otherwise. But, since V_j is a representation, for all $g \in G$, the action of g on V_j is a subspace of V_j . That is, the map ϕ_i does not cause any intertwining between the representations V_i .

As a consequence, ϕ_i is the identity on $V_i^{\oplus n_i}$ and the zero map on $V_j^{\oplus n_j}$ such that $i \neq j$, and thus ϕ_i is a projector.

A.1.4 Representations of \mathfrak{S}_n

The **symmetric group**, denoted as \mathfrak{S}_n , is the group of order $n!$ consisting of permutations of the set $\{1, \dots, n\}$, as illustrated in previous examples. The cyclic notation convention is employed throughout this thesis. For instance, $(1 \ 2 \ 3)(4 \ 5)$ indicates the permutation

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix},$$

of \mathfrak{S}_5 . According to [Theorem A.5](#), the number of irreducible representations of a finite group G is equal to the number of conjugacy classes in G . For the symmetric group \mathfrak{S}_n , the conjugacy class of a permutation is uniquely determined by the length of its decomposition into a product of disjoint cycles, which is referred to as the


cycle type of the permutation. The cycle type of a permutation can be described by an integer **partition** of n , where each part represents the length of a cycle in the permutation. For instance, the permutation $(1\ 2\ 3)(4\ 5)$ in \mathfrak{S}_5 has cycle type $(3, 2)$. It is worth noting that this justifies the use of cyclic notation in this thesis.

The notation $\lambda \vdash n$ is used to represent an ordered partition of the positive integer n into l parts, denoted by $\lambda = (\lambda_1, \dots, \lambda_l)$. This partition is defined as a collection of non-increasing positive integers that sum up to n , i.e.

$$\lambda_1 \geq \dots \geq \lambda_l \geq 1 \quad \text{and} \quad \sum_{i=1}^l \lambda_i = n.$$

A partition $\lambda \vdash n$ may be represented as a **Young diagram**, which is a collection of n empty boxes arranged in left-justified rows such that the i -th row contains λ_i boxes. This representation is illustrated in [Example A.8](#). The **conjugate** of the partition λ , denoted λ' , is the partition corresponding to transposing the Young diagram representing λ . A **Young tableau** is a Young diagram where each box is assigned an integer between 1 and n . A Young tableau is classified as **standard** if the entries in each row and each column are strictly increasing, **semistandard** if the entries in each row are weakly increasing and the entries in each column are strictly increasing, and **without repetitions** if each entry appears exactly once. Notably, standard Young tableaux are without repetitions. The **canonical Young tableau** is a Young diagram that is filled with consecutive integers from left to right and top to bottom.

Example A.8. Consider the permutation $\sigma \in \mathfrak{S}_9$, expressed as a product of disjoint cycles, arranged in non-increasing order, by $(1\ 2\ 3\ 4)(5\ 6\ 7)(8)(9)$. The cycle type of σ is the partition $\lambda \vdash 9$ given by $\lambda = (4, 3, 1, 1)$, where the i -th entry of λ denotes the length of the i -th cycle in the disjoint cycle decomposition of σ . Moreover, the partition λ can be represented using a Young diagram, which consists of 4 rows with 4, 3, 1, and 1 boxes, respectively



The conjugate of the partition λ , namely λ' is represented using a Young diagram,

which consists of 4 rows with 4, 2, 2 and 1 boxes, respectively:

Two instances of Young tableaux that correspond to the partition λ are provided as examples

1	2	3	4
2	3	4	
3			
4			

and

1	2	3	4
3	4	4	
4			
5			

The previous Young tableau presented on the left satisfies the standard definition, whereas the one on the right solely complies with the semistandard definition, primarily due to the weakly increasing gray box. Neither of them adheres to the requirement of being without repetitions. For the partition λ , the canonical Young tableau can be expressed as follows

1	2	3	4
5	6	7	
8			
9			

It is both standard and thus without repetition.

Consider a partition $\lambda \vdash n$ and a Young tableau T without repetitions, on this partition. An action of the symmetric group \mathfrak{S}_n on the Young tableau T is defined as follows: given a permutation $\sigma \in \mathfrak{S}_n$, the action of σ on T is obtained by permuting the boxes in T according to the permutation induced by the entries of the boxes. For $\sigma \in \mathfrak{S}_n$, the action of σ on T is denoted by $\sigma(T)$.

A permutation σ in \mathfrak{S}_n is said to preserve each row of T if each box of a given row is permuted by σ on that same row. Similarly, a permutation σ in \mathfrak{S}_n is said to preserve each column of T if each box of a given column is permuted by σ on that same column. Two subgroups of \mathfrak{S}_n can be defined as the sets of permutations that

preserve the rows and columns of T , namely,

$$\begin{aligned} R_T &:= \{\sigma \in \mathfrak{S}_n \mid \sigma \text{ preserves each row of } T\} \\ C_T &:= \{\sigma \in \mathfrak{S}_n \mid \sigma \text{ preserves each column of } T\}. \end{aligned}$$

Consider two elements of the group algebra $\mathbb{C}[\mathfrak{S}_n]$, the row and column symmetrizers of a given Young tableau T , defined as follows:

$$\begin{aligned} r_T &:= \sum_{\sigma \in R_T} \sigma \\ c_T &:= \sum_{\sigma \in C_T} \text{sign}(\sigma) \cdot \sigma. \end{aligned}$$

The **Young symmetrizer** associated with a given Young tableau T is an element s_T of $\mathbb{C}[\mathfrak{S}_n]$, defined as the product of the row and column symmetrizers of T , i.e. $s_T := r_T \cdot c_T$. It is worth noting that the row and column symmetrizers satisfy certain properties under the action of the symmetric group \mathfrak{S}_n . Specifically, for any $\sigma \in \mathfrak{S}_n$,

$$\sigma \cdot r_T \cdot \sigma^{-1} = r_{\sigma(T)} \quad \text{and} \quad \sigma \cdot c_T \cdot \sigma^{-1} = c_{\sigma(T)}.$$

As a consequence, it follows that $\sigma \cdot s_T \cdot \sigma^{-1} = s_{\sigma(T)}$.

Example. Consider a partition $\lambda \vdash 3$. When $\lambda = (3)$. In the case where $\lambda = (3)$, the associated canonical Young tableau T is a single row with entries 1, 2, and 3

$$T = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}$$

As the identity permutation is the unique element of \mathfrak{S}_3 that leaves each column of T unchanged, the Young symmetrizer can be expressed as

$$\begin{aligned} s_T &= r_T \cdot c_T \\ &= r_T \\ &= \sum_{\sigma \in \mathfrak{S}_3} \sigma. \end{aligned}$$

In the case where $\lambda = (1, 1, 1)$, the associated canonical Young tableau T is a single column with entries 1, 2, and 3

$$T = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$$

As the identity permutation is the unique element of \mathfrak{S}_3 that leaves each row of T unchanged, the Young symmetrizer can be expressed as

$$\begin{aligned} s_T &= r_T \cdot c_T \\ &= c_T \\ &= \sum_{\sigma \in \mathfrak{S}_3} \text{sign}(\sigma) \cdot \sigma. \end{aligned}$$

The two preceding Young symmetrizers correspond, to within a constant factor of $\frac{1}{6}$, to the projectors onto the irreducible representations V_{trivial} and V_{sign} in the decomposition of $\mathbb{C}[\mathfrak{S}_3]$ into a direct sum of irreducible representations:

$$\mathbb{C}[\mathfrak{S}_3] \simeq V_{\text{trivial}} \oplus V_{\text{sign}} \oplus V_{\text{standard}}^2,$$

as established by [Theorem A.7](#).

In order to obtain a comprehensive collection of inequivalent irreducible representations of the symmetric group \mathfrak{S}_n , it is necessary to first establish several requisite technical lemmas.

Lemma A.9. *Let $\lambda \vdash n$ be a partition of n , let T be a Young tableau on λ without repetitions, and let $x \in \mathbb{C}[\mathfrak{S}_n]$. Then, there exists $\mu \in \mathbb{C}$ such that the following equation holds: $s_T \cdot x \cdot s_T = \mu \cdot s_T$.*

Proof. To begin, it will be proven that if an element $y \in \mathbb{C}[\mathfrak{S}_n]$ satisfies the condition $\sigma \cdot y \cdot \tau = \text{sign}(\tau) \cdot y$, for all permutations $\sigma \in R_T$ and $\tau \in C_T$, then it follows that $y = \mu \cdot s_T$, for some scalar $\mu \in \mathbb{C}$. Consider such an element $y := \sum_{\pi \in \mathfrak{S}_n} c_\pi \cdot \pi$ of the group algebra $\mathbb{C}[\mathfrak{S}_n]$, then for all permutations $\sigma \in R_T$ and $\tau \in C_T$,

$$\sigma \cdot \left(\sum_{\pi \in \mathfrak{S}_n} c_\pi \cdot \pi \right) \cdot \tau = \sum_{\pi \in \mathfrak{S}_n} c_\pi \cdot (\sigma \cdot \pi \cdot \tau) = \text{sign}(\tau) \cdot \sum_{\pi \in \mathfrak{S}_n} c_\pi \cdot \pi.$$

This implies that, $c_{(\sigma \cdot \pi \cdot \tau)} = \text{sign}(\tau) \cdot c_\pi$ for all permutation $\pi \in \mathfrak{S}_n$. Then in particular case if the identity permutation, $c_{(\sigma \cdot \tau)} = \text{sign}(\tau) \cdot c_{1_{\mathfrak{S}_n}}$, and

$$\begin{aligned} \sum_{\substack{\sigma \in R_T \\ \tau \in C_T}} c_{(\sigma \cdot \tau)} \cdot (\sigma \cdot \tau) &= c_{1_{\mathfrak{S}_n}} \cdot \sum_{\substack{\sigma \in R_T \\ \tau \in C_T}} \text{sign}(\tau) \cdot (\sigma \cdot \tau) \\ &= c_{1_{\mathfrak{S}_n}} \cdot s_T. \end{aligned}$$

Assuming that $c_\pi = 0$ when $\pi \notin \{(\sigma \cdot \tau) \mid \sigma \in R_T \text{ and } \tau \in C_T\}$, and given that $R_T \cap C_T = \{1_{\mathfrak{S}_n}\}$, it follows that $y = c_{1_{\mathfrak{S}_n}} \cdot s_T$. Consider a permutation $\pi \notin \{(\sigma \cdot \tau) \mid \sigma \in R_T \text{ and } \tau \in C_T\}$, and suppose that there exist no distinct $i, j \in \{1, \dots, n\}$ such that i and j belong to the same row of T and the same column of $\pi(T)$. Then, all entries in the first row of T must appear in distinct columns of $\pi(T)$. Hence, there exist two permutations $\sigma \in R_T$ and $\tau \in C_{\pi(T)}$ such that the first row of $\sigma(T)$ and $(\tau \cdot \pi)(T)$ are identical. By iterating this process on the remaining rows of T , it follows that there exist two permutations $\sigma \in R_T$ and $\tau \in C_{\pi(T)}$ such that $\sigma(T) = (\tau \cdot \pi)(T)$. Consequently $\sigma = \tau \cdot \pi$ and $\pi \in \{(\sigma \cdot \tau) \mid \sigma \in R_T \text{ and } \tau \in C_T\}$ since $C_{\pi(T)} = \pi \cdot C_T \cdot \pi^{-1}$. Therefore, there exists a transposition $(ij) \in R_T \cap C_{\pi(T)}$, and in particular $(ij) \in C_{\pi(T)}$. Hence, $(ij) = \pi \cdot \tau \cdot \pi^{-1}$ for some permutation $\tau \in C_T$, then equality

$$c_{1_{\mathfrak{S}_n}} = (ij) \cdot (ij) = (ij) \cdot \pi \cdot \tau \cdot \pi^{-1},$$

holds, and finally $\pi = (ij) \cdot \pi \cdot \tau$. Now since $(ij) \in R_T$ and $c_{(\sigma \cdot \pi \cdot \tau)} = \text{sign}(\tau) \cdot c_\pi$ for all permutations $\sigma \in R_T$, $\tau \in C_T$ and $\pi \in \mathfrak{S}_n$, it holds that

$$\begin{aligned} c_\pi &= c_{((ij) \cdot \pi \cdot \tau)} \\ &= \text{sign}((ij)) \cdot c_\pi \\ &= -c_\pi, \end{aligned}$$

which implies that $c_\pi = 0$ when $\pi \notin \{(\sigma \cdot \tau) \mid \sigma \in R_T \text{ and } \tau \in C_T\}$.

Let $\sigma \in R_T$ and $\tau \in C_T$, then, the following equations hold:

$$\sigma \cdot r_T = r_T \cdot \sigma = r_T \quad \text{and} \quad \tau \cdot c_T = c_T \cdot \tau = \text{sign}(\tau) \cdot c_T.$$

Additionally $\sigma \cdot s_T \cdot \tau = \text{sign}(\tau) \cdot s_T$. Furthermore, for y an element of the group algebra $\mathbb{C}[\mathfrak{S}_n]$, it follows that

$$\sigma \cdot r_T \cdot y \cdot c_T \cdot \tau = \text{sign}(\tau) \cdot r_T \cdot y \cdot c_T.$$

It is inferred that there exists a $\mu \in \mathbb{C}$ for which the equality $r_T \cdot y \cdot c_T = \mu \cdot s_T$ holds. Furthermore, when $y = c_T \cdot x \cdot r_T$,

$$s_T \cdot x \cdot s_T = r_T \cdot c_T \cdot x \cdot r_T \cdot c_T = \mu \cdot s_T.$$

Corollary. *Given a partition $\lambda \vdash n$, and an arbitrary Young tableau T on λ without repetitions, the product of the corresponding Young symmetrizer s_T is nonzero, i.e. $s_T \cdot s_T \neq 0$.*

Lemma A.10. *Let $\lambda \vdash n$ and $\mu \vdash n$ be two distinct partitions of n , let T_λ and T_μ be two Young tableaux without repetitions on partitions λ and μ , respectively, and let $x \in \mathbb{C}[\mathfrak{S}_n]$. Then $s_{T_\mu} \cdot x \cdot s_{T_\lambda} = 0$.*

Proof. It is possible to assert, without loss of generality, that by interchanging λ and μ , there is a $k \in \{1, \dots, n\}$ for which $\lambda_k > \mu_k$ holds, and for all $i \in \{1, \dots, k-1\}$, the equalities $\lambda_i = \mu_i$ holds.

Suppose that there exists no pair of distinct indices $i, j \in \{1, \dots, n\}$ such that i and j occupy the same row in T_λ and the same column in T_μ . Under this condition, it follows that every element within the first row of T_λ occupies a unique column within T_μ . When $k > 1$, there exist two permutations $\sigma \in R_{T_\lambda}$ and $\tau \in C_{T_\mu}$ such that $\sigma(T_\lambda)$ and $\tau(T_\mu)$ have identical first rows. By iteratively applying this procedure to the first $k-1$ rows of T_λ and T_μ , it follows that there exist two permutations $\sigma \in R_{T_\lambda}$ and $\tau \in C_{T_\mu}$ such that $\sigma(T_\lambda)$ and $\tau(T_\mu)$ have the same first $k-1$ rows.

The equality $\lambda_k > \mu_k$ holds, otherwise $\lambda = \mu$. Consequently, the k -th row of $\sigma(T_\lambda)$ contains λ_k entries, which appear in λ_k distinct columns of $\tau(T_\mu)$, and are located between the k -th and n -th rows of $\tau(T_\mu)$. However, this arrangement cannot occur since $\lambda_k > \mu_k$. Therefore, there exists a transposition $(i j) \in R_{T_\lambda} \cap C_{T_\mu}$ such that

$$(i j) \cdot r_{T_\lambda} = r_{T_\lambda} \quad \text{and} \quad c_{T_\mu} \cdot (i j) = -c_{T_\mu}.$$

Furthermore, since $(i j) \cdot (i j)$ is the identity element in the symmetric group \mathfrak{S}_n , then

$$c_{T_\mu} \cdot r_{T_\lambda} = c_{T_\mu} \cdot (i j) \cdot (i j) \cdot r_{T_\lambda} = -c_{T_\mu} \cdot r_{T_\lambda},$$

which implies that $c_{T_\mu} \cdot r_{T_\lambda} = 0$.

Let σ be an element of the symmetric group \mathfrak{S}_n . Then $c_{T_\mu} \cdot \sigma \cdot r_{T_\lambda} \cdot \sigma^{-1} = c_{T_\mu} \cdot r_{\sigma(T_\lambda)} = 0$, as the previous equality is independent of the entries of T_λ . Thus, it follows that $c_{T_\mu} \cdot \sigma \cdot r_{T_\lambda} = 0$ for any permutation $\sigma \in \mathfrak{S}_n$. Consequently for all $x \in \mathbb{C}[\mathfrak{S}_n]$ it follows that $c_{T_\mu} \cdot x \cdot r_{T_\lambda} = 0$ and also $s_{T_\mu} \cdot x \cdot s_{T_\lambda} = 0$.

Corollary. *Given two distinct partitions $\lambda \vdash n$ and $\mu \vdash n$, and two arbitrary Young tableaux without repetitions on partitions λ and μ , respectively, the product of the corresponding Young symmetrizers s_{T_μ} and s_{T_λ} is zero, i.e. $s_{T_\mu} \cdot s_{T_\lambda} = 0$.*

Let $\lambda \vdash n$ be a partition of n , and let T be a Young tableau on λ without repetitions. The operation of right multiplication by the Young symmetrizer s_T on the group algebra $\mathbb{C}[\mathfrak{S}_n]$, i.e., $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$, defines a complex vector space that constitutes a representation of the symmetric group \mathfrak{S}_n through its left action on this space.

Lemma A.11. *For any partition $\lambda \vdash n$, and any Young tableau T on λ without repetitions, the representation $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$ is an irreducible representation of the symmetric group \mathfrak{S}_n .*

Proof. Consider two permutations $\sigma \in R_T$ and $\tau \in C_T$. Then, their product $\sigma \cdot \tau$ is equal to the identity element $1_{\mathfrak{S}_n}$ if and only if both σ and τ are equal to $1_{\mathfrak{S}_n}$. This equivalence is due to the fact that $R_T \cap C_T = \{1_{\mathfrak{S}_n}\}$. Consequently $s_T \neq 0$ and $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$ is nonzero.

Consider V as a subrepresentation of $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$, then using [Lemma A.9](#), for all $x \in V$ there exists a scalar $\mu \in \mathbb{C}$ such that $s_T \cdot x = \mu \cdot s_T$. Thus

$$s_T \cdot V \subset \mathbb{C} \cdot s_T.$$

Since $\mathbb{C} \cdot s_T$ has dimension one, the subspace $s_T \cdot V$ is either equal to $\mathbb{C} \cdot s_T$ or to the zero space $\{0\}$. In the former case, the inclusion $\mathbb{C}[\mathfrak{S}_n] \cdot s_T \cdot V \subseteq V$ follows, since V is a representation of the symmetric group \mathfrak{S}_n , and the equality $V = \mathbb{C}[\mathfrak{S}_n] \cdot s_T$ holds. Similarly, in the latter case, since V is a representation of the symmetric group \mathfrak{S}_n , the inclusion $V \cdot V \subseteq V$ holds. Moreover, as V is a subrepresentation of $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$, i.e., $V \subseteq \mathbb{C}[\mathfrak{S}_n] \cdot s_T$, it follows that the product of $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$ and V yields the trivial space $\{0\}$, and the product of V with itself also yields $\{0\}$, namely $\mathbb{C}[\mathfrak{S}_n] \cdot s_T \cdot V = \{0\}$ and $V \cdot V = \{0\}$. Consider an element x in the subrepresentation V , defined as $x := \sum_{\pi \in \mathfrak{S}_n} c_\pi \cdot \pi$. The adjoint of x , denoted as x^* , is defined by

$$x^* := \sum_{\sigma \in \mathfrak{S}_n} \bar{c}_\sigma \cdot \sigma^{-1}.$$

Then both x^* and $x \cdot x^*$ belong to V . As $V \cdot V = \{0\}$ it follows that $x \cdot x^* = 0$. Consequently, for all permutations $\sigma \in \mathfrak{S}_n$ it holds that $c_\sigma \cdot x^* = 0$. In particular

$$c_e \cdot x^* = \sum_{\sigma \in \mathfrak{S}_n} \bar{c}_\sigma \cdot \sigma = 0.$$

Therefore $x = 0$, and thus $V = \{0\}$.

Lemma A.12. *For every partition $\lambda \vdash n$, every Young tableau T on λ without repetitions, and every permutation σ in the symmetric group \mathfrak{S}_n , the representations $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$ and $\mathbb{C}[\mathfrak{S}_n] \cdot s_{\sigma(T)}$ are equivalent.*

Proof. Consider the linear map ϕ from $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$ to $\mathbb{C}[\mathfrak{S}_n] \cdot s_{\sigma(T)}$. Given $x \in \mathbb{C}[\mathfrak{S}_n] \cdot s_T$, with $x = y \cdot s_T$ for some $y \in \mathbb{C}[\mathfrak{S}_n]$, then ϕ is defined on x by $\phi(x) = x \cdot \sigma^{-1}$. Observe that

$$\begin{aligned} \phi(x) &= x \cdot \sigma^{-1} \\ &= y \cdot s_T \cdot \sigma^{-1} \\ &= y \cdot \sigma^{-1} \cdot \sigma \cdot s_T \cdot \sigma^{-1} \\ &= y \cdot \sigma^{-1} \cdot s_{\sigma(T)}. \end{aligned}$$

It follows that $\phi(x)$ belongs to $\mathbb{C}[\mathfrak{S}_n] \cdot s_{\sigma(T)}$, and that ϕ is an isomorphism.

Consider a permutation $\tau \in \mathfrak{S}_n$ and x is an element of $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$. Then, it follows that

$$\begin{aligned} \phi \circ (\tau \cdot x) &= \tau \cdot x \cdot \sigma^{-1} \\ &= \tau \circ \phi(x), \end{aligned}$$

and hence ϕ satisfies the property of being intertwining.

Lemma A.13. *Let $\lambda \vdash n$ and $\mu \vdash n$ be two distinct partitions of n . Let T_λ and T_μ be two Young tableaux without repetitions on partitions λ and μ , respectively. Then, the two representations $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\lambda}$ and $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\mu}$ are inequivalent.*

Proof. According to [Lemma A.9](#), it follows that the left action of the Young symmetrizer s_{T_μ} on $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\mu}$ results in

$$s_{T_\mu} \cdot \mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\mu} = \mathbb{C} \cdot s_{T_\mu}.$$

However, from [Lemma A.10](#), the left action of the Young symmetrizer s_{T_μ} on $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\lambda}$ is zero, i.e.

$$s_{T_\mu} \cdot \mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\lambda} = 0.$$

As a consequence, two irreducible representations $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\lambda}$ and $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_\mu}$ are not isomorphic, hence not equivalent.

From [Lemma A.12](#), for a given partition $\lambda \vdash n$, the representations $\mathbb{C}[\mathfrak{S}_n] \cdot s_T$ of the symmetric group \mathfrak{S}_n , are mutually equivalent across all Young tableaux T on

λ without repetition, thereby solely dependent on λ , and consequently, it is possible to designate any representation of this type as V_λ .

Theorem A.14. *All irreducible representations of \mathfrak{S}_n can be expressed as V_λ for some partition $\lambda \vdash n$.*

Proof. The irreducibility of the representation V_λ is guaranteed for any partition $\lambda \vdash n$, as stated in [Lemma A.11](#). It follows from [Lemma A.12](#) and [Lemma A.13](#) that the number of inequivalent irreducible representations V_λ for some partitions $\lambda \vdash n$ is equivalent to the number of conjugacy classes of the symmetric group \mathfrak{S}_n . This number, as stated in [Theorem A.5](#), corresponds to the number of irreducible representations of \mathfrak{S}_n .

The [Proposition A.6](#) establishes that upon decomposing the group algebra $\mathbb{C}[\mathfrak{S}_n]$ into a direct sum of irreducible representations, namely $\mathbb{C}[\mathfrak{S}_n] \simeq V_1^{\oplus n_1} \oplus \dots \oplus V_k^{\oplus n_k}$, the dimension of each irreducible representation V_i coincides with its multiplicity n_i .

Theorem A.15. *Consider a partition $\lambda \vdash n$. Let V_λ be the corresponding irreducible representation of a symmetric group \mathfrak{S}_n . The dimension of V_λ is equivalent to the cardinality of the set of standard Young tableaux associated with λ .*

Proof. Consider a partition $\lambda \vdash n$ and let $f(\lambda)$ denote the number of standard Young tableaux associated with λ . The standard Young tableaux on this partition can be ordered lexicographically based on their entries. Specifically, let $T_1, T_2, \dots, T_{f(\lambda)}$ be the standard Young tableaux on this partition, ordered such that $T_i < T_j$ if and only if the entries of T_i are smaller in lexicographic order than those of T_j , from left to right and top to bottom. Note that the first tableau in this order, denoted by T_1 , is the canonical Young tableau on this partition.

Let i and j be arbitrary elements of $\{1, \dots, f(\lambda)\}$, with $i < j$. Further, let k and l denote the first row and column, respectively, at which the two standard Young tableaux T_i and T_j differ, proceeding from left to right and top to bottom. Notably, k and l cannot be the first row or column, respectively, since the Young tableaux are standard. Let a denote the entry of T_i located in the k -th row and l -th column, and b denote the entry of T_j located in the same position. Given that $i < j$ and T_i and T_j are standard Young tableaux, it follows that $a < b$. Let m and n denote the row and column, respectively, of the entry a in T_j . By virtue of T_j being a standard Young tableau, it cannot hold that $m > k$ and $n \geq l$, as the entries of T_j in rows greater than k and

columns greater than l are strictly larger than b . Furthermore, it cannot be the case that $m < k$, or that $m = k$ and $n < l$, since T_i and T_j coincide on these rows and columns. Lastly, it cannot hold that $m = k$ and $n > l$, as T_j is a standard Young tableau and $a < b$. Thus $m > k$ and $n < l$ hold necessarily. Notably, the entries located on the k -th row and n -th column of both T_i and T_j are equal and denoted by c . Consequently, the transposition $(a\ c)$ belongs to $R_{T_i} \cap C_{T_j}$, since a and c share the same k -th row of T_i and the same n -th column of T_j . Then, the equations

$$(a\ c) \cdot r_{T_i} = r_{T_i} \quad \text{and} \quad c_{T_j} \cdot (a\ c) = -c_{T_j},$$

hold. Moreover, as $(a\ c) \cdot (a\ c)$ yields the identity element in the symmetric group \mathfrak{S}_n ,

$$\begin{aligned} s_{T_j} \cdot s_{T_i} &= r_{T_j} \cdot c_{T_j} \cdot r_{T_i} \cdot c_{T_i} \\ &= r_{T_j} \cdot c_{T_j} \cdot (a\ c) \cdot (a\ c) \cdot r_{T_i} \cdot c_{T_i} \\ &= -s_{T_j} \cdot s_{T_i}, \end{aligned}$$

implying that $s_{T_j} \cdot s_{T_i} = 0$.

From [Lemma A.12](#), the irreducible representations $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_i}$ and $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_j}$ are equivalent for all $i, j \in \{1, \dots, f(\lambda)\}$. Let $x_1, x_2, \dots, x_{f(\lambda)}$ denote certain elements of the group algebra $\mathbb{C}[\mathfrak{S}_n]$, such that

$$\sum_{i=1}^{f(\lambda)} x_i \cdot s_{T_i} = 0.$$

By applying the result of [Lemma A.9](#) and observing that $T_1 \leq T_i$ for all $i \in \{1, \dots, f(\lambda)\}$, the existence of a nonzero $\mu \in \mathbb{C}$ is established, such that $\sum_{i=1}^{f(\lambda)} x_i \cdot s_{T_i} \cdot s_{T_1} = \mu \cdot x_1 \cdot s_{T_1}$, which further implies that $x_1 = 0$. Right multiplication on both sides of the equation with s_{T_i} yields $x_i = 0$ for all $i \in \{1, \dots, f(\lambda)\}$. Consequently, $\dim V_\lambda \geq f(\lambda)$.

Consider a Young diagram λ . Let μ be another Young diagram obtained by either removing a single box from λ , denoted as $\mu = \lambda - 1$, or adding a single box to λ , denoted as $\mu = \lambda + 1$. Then $f(\lambda)$ can be expressed as the sum of $f(\mu)$ for all possible Young diagrams μ obtained by removing a single box from λ , i.e.

$$f(\lambda) = \sum_{\mu=\lambda-1} f(\mu).$$

The proof of the identity $(n+1)f(\lambda) = \sum_{\mu=\lambda+1} f(\mu)$ shall be established through induction on n . The base case $n = 0$ holds, as $f((0)) = f((1))$. For the inductive step, assume $n > 0$, it follows that

$$\begin{aligned} (n+1)f(\lambda) &= n \cdot f(\lambda) + f(\lambda) \\ &= n \sum_{\mu=\lambda-1} f(\mu) + f(\lambda) \\ &= \sum_{\mu=\lambda-1} \sum_{\nu=\mu+1} f(\nu) + f(\lambda), \end{aligned}$$

wherein the second summand enumerates partitions $\nu \vdash n$. There exist two distinct cases, namely $\nu = \lambda$ and $\nu \neq \lambda$. Consider the sets λ_+ and λ_- , defined as follows:

$$\begin{aligned} \lambda_+ &= \{\mu \mid \mu = \lambda + 1\} \\ \lambda_- &= \{\mu \mid \mu = \lambda - 1\}. \end{aligned}$$

It is worth noting that for each box within a given Young diagram that may be extracted such that it does not alter its validity as a Young diagram, a box can be added to the next row of the Young diagram. Furthermore, it is always possible to add a box to the first row of the Young diagram. As a result, it follows that $\lambda_+ = \lambda_- + 1$. The occurrence of the first case, where $\nu = \lambda$, is equal to the cardinality of the set λ_- , namely $|\lambda_-|$. Thus

$$\begin{aligned} \sum_{\mu=\lambda-1} \sum_{\nu=\mu+1} f(\nu) + f(\lambda) &= \sum_{\mu=\lambda-1} \sum_{\substack{\nu=\mu+1 \\ \nu \neq \lambda}} f(\nu) + (|\lambda_-| + 1)f(\lambda) \\ &= \sum_{\mu=\lambda+1} \sum_{\substack{\nu=\mu-1 \\ \nu \neq \lambda}} f(\nu) + |\lambda_+| \cdot f(\lambda) \\ &= \sum_{\mu=\lambda+1} \sum_{\nu=\mu-1} f(\nu) \\ &= \sum_{\mu=\lambda+1} f(\mu), \end{aligned}$$

concluding the induction proof.

The formula $\sum_{\lambda \vdash n} f(\lambda)^2 = n!$ shall also be established through induction on n . As the base case, it is observed that $f((0)) = 1$, and hence the formula


is verified for $n = 0$. For the inductive step, suppose $n > 0$, then

$$\begin{aligned}
 \sum_{\lambda \vdash n} f(\lambda)^2 &= \sum_{\lambda \vdash n} \sum_{\mu = \lambda - 1} f(\lambda) f(\mu) \\
 &= \sum_{\lambda \vdash (n-1)} \sum_{\mu = \lambda + 1} f(\lambda) f(\mu) \\
 &= n \sum_{\lambda \vdash (n-1)} f(\lambda)^2 \\
 &= n \cdot (n-1)!,
 \end{aligned}$$


concluding the induction proof.

According to [Proposition A.6](#) the following equation is satisfied: $\sum_{\lambda \vdash n} \dim(V_\lambda)^2 = n!$. However, for all partitions $\lambda \vdash n$, it is true that $\dim V_\lambda \geq f(\lambda)$. Hence, $\dim V_\lambda = f(\lambda)$.


Example. There exist three partitions of 3, which are denoted as (3) , $(2, 1)$ and $(1, 1, 1)$. Each partition has a corresponding Young diagram



For the partitions (3) and $(1, 1, 1)$, there exists only one standard Young tableau, which corresponds to the canonical Young tableau



However the partition $(2, 1)$ admits a pair of distinct standard Young tableaux, denoted as follows



From [Theorem A.15](#), it follows that the irreducible representations $V(3)$ and $V_{(1,1,1)}$, of \mathfrak{S}_3 , have dimension 1, while $V_{(2,1)}$ has dimension 2, as expected. The determination of the irreducible representations of \mathfrak{S}_3 corresponding to the canonical Young tableaux $T_{(3)}$, $T_{(2,1)}$, and $T_{(1,1,1)}$ necessitates an examination of their respective Young symmetrizers $s_{T_{(3)}}$, $s_{T_{(1,1,1)}}$ and $s_{T_{(2,1)}}$. This examination is conducted in conjunction with the construction of V_λ 's From [Theorem A.14](#),

$$\begin{aligned}\mathbb{C}[\mathfrak{S}_3] \cdot s_{T_{(3)}} &= \mathbb{C}[\mathfrak{S}_3] \cdot (1_{\mathfrak{S}_3} + (1\ 2) + (1\ 3) + (2\ 3) + (1\ 2\ 3) + (3\ 2\ 1)) \\ \mathbb{C}[\mathfrak{S}_3] \cdot s_{T_{(1,1,1)}} &= \mathbb{C}[\mathfrak{S}_3] \cdot (1_{\mathfrak{S}_3} - (1\ 2) - (1\ 3) - (2\ 3) + (1\ 2\ 3) + (3\ 2\ 1)) \\ \mathbb{C}[\mathfrak{S}_3] \cdot s_{T_{(2,1)}} &= \mathbb{C}[\mathfrak{S}_3] \cdot (1_{\mathfrak{S}_3} + (1\ 2) - (1\ 3) + (3\ 2\ 1)).\end{aligned}$$

The action of the symmetric group \mathfrak{S}_3 on the irreducible representation $\mathbb{C}[\mathfrak{S}_n] \cdot s_{T_{(3)}}$ yields a trivial action. It follows that $V_{(3)}$ is equivalent to V_{trivial} , while the irreducible representations $V_{(1,1,1)}$ and $V_{(2,1)}$ are necessarily equivalent to V_{sign} and V_{standard} , respectively.

Let $\mathbb{C}[\mathfrak{S}_n] \simeq V_1^{\oplus n_1} \oplus \cdots \oplus V_k^{\oplus n_k}$ be the decomposition the group algebra $\mathbb{C}[\mathfrak{S}_n]$ of the symmetric group \mathfrak{S}_n into a direct sum of irreducible representations. By [Theorem A.7](#) the projectors onto the isotypic components $V_i^{\oplus n_i}$ are given by

$$\phi_i := \frac{\dim V_i}{n!} \sum_{\sigma \in \mathfrak{S}_n} \bar{\chi}_{V_i}(\sigma) \cdot \sigma.$$

Theorem. *Let the group algebra $\mathbb{C}[\mathfrak{S}_n]$ of the symmetric group \mathfrak{S}_n and its decomposition into a direct sum of irreducible representations $\mathbb{C}[\mathfrak{S}_n] \simeq \bigoplus_{\lambda \vdash n} V_\lambda^{\oplus n_\lambda}$. For each partition $\lambda \vdash n$, the map ϕ_λ defined on $\mathbb{C}[\mathfrak{S}_n]$ by*

$$\phi_\lambda := \frac{\dim V_\lambda}{n!} \sum_T s_T,$$

where the sum is taken over all Young tableaux without repetitions on λ , is the projector onto $V_\lambda^{\oplus n_\lambda}$.

Proof. Let $\sigma \in \mathfrak{S}_n$, then

$$\begin{aligned}\sigma^{-1} \cdot \sum_T s_T \cdot \sigma &= \sum_T s_{\sigma(T)} \\ &= \sum_T s_T,\end{aligned}$$

and in hence $\sum_T s_T \in \text{Hom}_{\mathfrak{S}_n}(\mathbb{C}[\mathfrak{S}_n])$. Consider the decomposition of the group algebra $\mathbb{C}[\mathfrak{S}_n]$ of the symmetric group \mathfrak{S}_n into a direct sum of irreducible representations $\mathbb{C}[\mathfrak{S}_n] \simeq \bigoplus_{\lambda \vdash n} V_\lambda^{\oplus n_\lambda}$, for all $\mu \vdash n$ partition of n , from Schur's [Lemma A.2](#), the restriction of $\sum_T s_T$ to V_μ is an homothety $\lambda \cdot I$, with $\lambda \in \mathbb{C}$. From [Lemma A.10](#), if λ and μ are distinct, then for all $x \in V_\mu$,

$$\sum_T s_T \cdot x = 0,$$

and thus the restriction of $\sum_T s_T$ to V_μ is the zero map. Let T_λ be any Young tableau without repetitions on λ , then the coefficient of $1_{\mathfrak{S}_n}$ in s_{T_λ} is 1, since $R_T \cap C_T = \{1_{\mathfrak{S}_n}\}$, and thus $\text{Tr} \left[\sum_T s_T \right] = n!$. So the restriction of $\sum_T s_T$ to V_λ is the homothety

$$\frac{n!}{\dim V_\lambda} \cdot I.$$

As a consequence, since $\sum_T s_T$ does not cause any intertwining between the representations V_λ , the map ϕ_λ is the identity on $V_\lambda^{\oplus n_\lambda}$ and the zero map elsewhere, and thus is a projector.

In the previous section, the complex vector spaces associated with the irreducible representations of the symmetric group \mathfrak{S}_n were discussed. However, the corresponding matrices representing the permutation elements were not described. The construction of such matrices is far from trivial, and in fact, there exist a variety of constructions that may be employed based on the desired properties of the resulting matrices, i.e. integer matrix components, rational matrix components, or orthogonal matrices.

A.1.5 Restricted and induced representations

The present Section concerns the correlation existing between a finite group G and a subgroup H . Can a representation of H be derived from a representation of G or vice versa? Furthermore, if the original representation is irreducible, what conclusions can be drawn concerning the derived representation?

Consider a representation (ρ, V) of a finite group G , and let H be a subgroup of G . Given an element $h \in H$, it is worth noting that H is a subgroup of G and hence $h \in G$. Furthermore, as the representation ρ induces an action of G on V , it follows that the action of ρ also restricts to only elements $h \in H$. The **restricted**

representation of (ρ, V) on the subgroup H is denoted $(\text{Res}_H^G(\rho), V)$ and defined on the element $h \in H$ as follows:

$$\text{Res}_H^G(\rho)(h) := \rho(h).$$

In the case where (ρ, V) is an irreducible representation of the group G , it is not always the case that the complex vector space V manifests irreducibility as a representation of a subgroup H . This can occur when the size of V becomes too large to maintain irreducibility with respect to the smaller subgroup H .

If χ is the character of the representation (ρ, V) of G , the **restricted character** of the representation $\text{Res}_H^G(\rho)$, denoted $\text{Res}_H^G(\chi)$, becomes $\text{Res}_H^G(\chi)(h) = \chi(h)$, for all $h \in H$.

Example. Let \mathfrak{S}_4 be the group of all permutations of $\{1, 2, 3, 4\}$. From [Section A.1.4](#), the symmetric group \mathfrak{S}_4 has 5 conjugacy classes, and consequently 5 irreducible representations, namely

$$V_{(4)}, \quad V_{(3,1)}, \quad V_{(2,2)}, \quad V_{(2,1,1)} \quad \text{and} \quad V_{(1,1,1,1)}.$$

Let D_4 be the **dihedral group** of order 8, whose elements are the symmetries of the square, generated by the $\frac{\pi}{4}$ counterclockwise rotation r and the vertical reflection s :

$$D_4 = \langle r, s \rangle = \{1_{D_4}, r, r^2, r^3, s, sr, sr^2, sr^3\}.$$

It has 5 conjugacy classes, which are given by


$$\{1_{D_4}\}, \quad \{r, r^3\}, \quad \{r^2\}, \quad \{sr, sr^3\} \quad \text{and} \quad \{s, sr^2\},$$

and as many irreducible representations: W_1, W_2, W_3, W_4 and W_5 . Then character table of D_4 can be expressed as follows

D_4	$\{1_{D_4}\}$	$\{r, r^3\}$	$\{r^2\}$	$\{sr, sr^3\}$	$\{s, sr^2\}$
χ_{W_1}	1	1	1	1	1
χ_{W_2}	1	-1	1	-1	1
χ_{W_3}	1	-1	1	1	-1
χ_{W_4}	1	1	1	-1	-1
χ_{W_5}	2	0	-2	0	0

As can be inferred from [Theorem A.5](#), the rows exhibit orthogonality with respect to the Hermitian inner product $\langle \cdot, \cdot \rangle_{D_4}$. By indexing the vertices of the square

counterclockwise with 1, 2, 3 and 4, starting from upper left corner, i.e.



the group D_4 can be made a subgroup of \mathfrak{S}_4 , where the 8 symmetries of the square become,

$(1)(2)(3)(4)$	identity
$(1, 2, 3, 4)$	$\frac{\pi}{4}$ counterclockwise rotation
$(1, 3)(2, 4)$	$\frac{2\pi}{4}$ counterclockwise rotation
$(4, 3, 2, 1)$	$\frac{3\pi}{4}$ counterclockwise rotation
$(1, 2)(3, 4)$	vertical reflection
$(1, 4)(2, 3)$	horizontal reflection
$(1, 3)(2)(4)$	main diagonal reflection
$(2, 4)(1)(3)$	secondary diagonal reflection

In the general case, the restricted representations $\text{Res}_{D_4}^{\mathfrak{S}_4}(V_\lambda)$ of D_4 , for every partition $\lambda \vdash 4$, do not constitute irreducible representations. According to Maschke's [Theorem A.1](#), these representations can be decomposed into a direct sum of irreducible representations. Specifically,

$$\begin{aligned}
 \text{Res}_{D_4}^{\mathfrak{S}_4}(V_{(4)}) &\simeq W_1 \\
 \text{Res}_{D_4}^{\mathfrak{S}_4}(V_{(3,1)}) &\simeq W_3 \oplus W_5 \\
 \text{Res}_{D_4}^{\mathfrak{S}_4}(V_{(2,2)}) &\simeq W_1 \oplus W_2 \\
 \text{Res}_{D_4}^{\mathfrak{S}_4}(V_{(2,1,1)}) &\simeq W_4 \oplus W_5 \\
 \text{Res}_{D_4}^{\mathfrak{S}_4}(V_{(1,1,1,1)}) &\simeq W_2.
 \end{aligned}$$

The passage from a representation (ρ, V) of a subgroup H of a finite group G to the group G necessitates a more intricate approach. This arises due to the inability of ρ to induce an action of $G \setminus H$ on V . Nevertheless, it is feasible to look at G/H , the **quotient group**, comprising elements in the form of $g \cdot h$ with $g \in G$ and $h \in H$, where $\rho(h)$ exhibits well-defined behavior.

Let V, W be two complex vector space of finite dimension. The complex tensor product between V and W , namely $V \otimes W$, is a condensed notation for the explicit

notation $V \otimes_{\mathbb{C}} W$. It is worth noting that for any $v \in V$ and $w \in W$, the relation

$$(c \cdot v) \otimes w = v \otimes (c \cdot w),$$

holds for all $c \in \mathbb{C}$.

Given a subgroup H of a finite group G , suppose that (ρ, V) is a representation of H . In this case, it is possible to define the tensor product complex vector space $\mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$ as follow: for all $g \in \mathbb{C}[G]$ and $v \in V$, the following relation,

$$(g \cdot h) \otimes v = g \otimes (h \cdot v),$$

holds for all $h \in H$. The **induced representation** of (ρ, V) on the group G is denoted $(\text{Ind}_H^G(\rho), \mathbb{C}[G] \otimes_{\mathbb{C}[H]} V)$ and defined on element $g \in G$ by

$$\text{Ind}_H^G(\rho)(g)(x) := g \cdot x,$$

for all $x \in \mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$.

In order to gain a more comprehensive understanding of the operation of group G on $\mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$, consider a **coset representation** of G , i.e. $G = g_1 \cdot H \uplus \dots \uplus g_k \cdot H$, where \uplus denotes the disjoint union, with $g_i \in G$. For any element $g \in G$, there exists $i \in \{1, \dots, k\}$ and $h \in H$ such that $g = g_i \cdot h$. Consequently, all elements $g \otimes v \in \mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$ satisfy

$$g \otimes v = (g_i \cdot h) \otimes v = g_i \otimes (h \cdot v).$$

Thus the action of the group G on the tensor product $\mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$ is exclusively defined for the elements $g_i \otimes v \in \mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$. Specifically, for all $g \in G$, the representation $\text{Ind}_H^G(\rho)(g)$ takes the form of a $k \times k$ block matrix that corresponds to the coset representation $G = g_1 \cdot H \uplus \dots \uplus g_k \cdot H$. The action of an element $g \in G$ upon $g_i \otimes v \in \mathbb{C}[G] \otimes_{\mathbb{C}[H]} V$ is given by

$$\begin{aligned} g \cdot (g_i \otimes v) &= (g \cdot g_i) \otimes v \\ &= (g_j \cdot h) \otimes v = g_j \otimes (h \cdot v), \end{aligned}$$

where h is the element of H such that $g \cdot g_i = g_j \cdot h$ in the coset representation $G = g_1 \cdot H \uplus \dots \uplus g_k \cdot H$. In other words, h is defined as $h := g_j^{-1} \cdot g \cdot g_i$. Additionally, it follows that h acts on a vector $v \in V$ via the representation $\rho(g_j^{-1} \cdot g \cdot g_i)(v)$. Finally,

$$\text{Ind}_H^G(\rho)(g) = \begin{pmatrix} \rho(g_1^{-1} \cdot g \cdot g_1) & \rho(g_1^{-1} \cdot g \cdot g_2) & \dots & \rho(g_1^{-1} \cdot g \cdot g_k) \\ \rho(g_1^{-2} \cdot g \cdot g_1) & \rho(g_1^{-2} \cdot g \cdot g_2) & \dots & \rho(g_1^{-2} \cdot g \cdot g_k) \\ \vdots & \vdots & \ddots & \vdots \\ \rho(g_1^{-k} \cdot g \cdot g_1) & \rho(g_1^{-k} \cdot g \cdot g_2) & \dots & \rho(g_1^{-k} \cdot g \cdot g_k) \end{pmatrix}.$$

If χ is the character of the representation (ρ, V) of H , the **induced character** of the representation $\text{Ind}_H^G(\rho)$, denoted $\text{Ind}_H^G(\chi)$ becomes, for all $g \in G$,

$$\text{Ind}_H^G(\chi) = \text{Tr} [\text{Ind}_H^G(\rho)(g)] = \sum_{i=1}^k \chi(g_i^{-1} \cdot g \cdot g_i),$$

in the coset representation $G = g_1 \cdot H \uplus \cdots \uplus g_k \cdot H$.

Example. Let $\mathbb{Z}/4\mathbb{Z}$ be the cyclic group of order 4 generated by the element z . Formally, $\mathbb{Z}/4\mathbb{Z}$ is defined by $\mathbb{Z}/4\mathbb{Z} := \{1_{\mathbb{Z}/4\mathbb{Z}}, z, z^2, z^3\}$. Let $\mathbb{Z}/2\mathbb{Z}$ the subgroup of $\mathbb{Z}/4\mathbb{Z}$ of order 2 generated by z^2 , i.e. $\mathbb{Z}/2\mathbb{Z} := \{1_{\mathbb{Z}/2\mathbb{Z}}, z^2\}$. Consider the one-dimensional irreducible representation (ρ, \mathbb{C}) of $\mathbb{Z}/2\mathbb{Z}$, defined as follows:

$$\rho(1_{\mathbb{Z}/4\mathbb{Z}}) = 1 \quad \text{and} \quad \rho(z^2) = -1.$$

The induced representation $\text{Ind}_{\mathbb{Z}/2\mathbb{Z}}^{\mathbb{Z}/4\mathbb{Z}}(\rho)$ of $\mathbb{Z}/4\mathbb{Z}$ is defined on the tensor product $\mathbb{C}[\mathbb{Z}/4\mathbb{Z}] \otimes_{\mathbb{C}[\mathbb{Z}/2\mathbb{Z}]} \mathbb{C}$ generated by the vectors

$$1_{\mathbb{Z}/4\mathbb{Z}} \otimes 1, \quad z \otimes 1, \quad z^2 \otimes 1 \quad \text{and} \quad z^3 \otimes 1.$$

But by definition of this tensor product over the group algebra $\mathbb{C}[\mathbb{Z}/2\mathbb{Z}]$, the two following relations hold:

$$z^2 \otimes 1 = 1_{\mathbb{Z}/4\mathbb{Z}} \otimes (z^2 \cdot 1) = -1 \cdot (1_{\mathbb{Z}/4\mathbb{Z}} \otimes 1),$$

and

$$z^3 \otimes 1 = z \otimes (z^2 \cdot 1) = -1 \cdot (z \otimes 1).$$

Thus $\mathbb{C}[\mathbb{Z}/4\mathbb{Z}] \otimes_{\mathbb{C}[\mathbb{Z}/2\mathbb{Z}]} \mathbb{C}$ is a 2-dimensional complex vector space with basis $1_{\mathbb{Z}/4\mathbb{Z}} \otimes 1$ and $z \otimes 1$. The action of the elements of $\mathbb{Z}/4\mathbb{Z}$ on the tensor product $\mathbb{C}[\mathbb{Z}/4\mathbb{Z}] \otimes_{\mathbb{C}[\mathbb{Z}/2\mathbb{Z}]} \mathbb{C}$, defined with respect to the generator z , is given by

$$\text{Ind}_{\mathbb{Z}/2\mathbb{Z}}^{\mathbb{Z}/4\mathbb{Z}}(\rho)(z) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Let $C(G)$ denotes the complex vector space of class functions of a finite group G , and $C(H)$ denotes the complex vector space of class functions of a subgroup H of G . There is an linear map $\phi : C(G) \rightarrow C(H)$ given by restriction of class functions. As linear map between Hermitian inner product complex vector spaces of

finite dimension, there exists a unique adjoint map $\phi^* : C(H) \rightarrow C(G)$ satisfying

$$\langle \phi^*(g), f \rangle_G = \langle g, \phi(f) \rangle_H,$$

for all $f \in C(G)$ and $g \in C(H)$.

Theorem A.16 (Frobenius reciprocity). *Consider a finite group G and a subgroup H of G . Let (ρ, V) and (σ, W) be representations of G and H , respectively. Then*

$$\langle \text{Ind}_H^G(\chi_W), \chi_V \rangle_G = \langle \chi_W, \text{Res}_H^G(\chi_V) \rangle_H.$$

Proof. By definition of the Hermitian inner product of class functions on finite groups, the properties of the characters of [Proposition A.4](#), and the restricted and induced characters in the coset representation $G = g_1 \cdot H \uplus \cdots \uplus g_k \cdot H$,

$$\begin{aligned} \langle \text{Ind}_H^G(\chi_W), \chi_V \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \text{Ind}_H^G(\bar{\chi}_W)(g) \cdot \chi_V(g) \\ &= \frac{1}{|G|} \sum_{g \in G} \sum_{i=1}^k \bar{\chi}_W(g_i^{-1} \cdot g \cdot g_i) \cdot \chi_V(g) \\ &= \frac{1}{|G|} \frac{1}{|H|} \sum_{g \in G} \sum_{h \in H} \bar{\chi}_W(h^{-1} \cdot g \cdot h) \cdot \chi_V(g) \\ &= \frac{1}{|G|} \frac{1}{|H|} \sum_{g \in G} \sum_{h \in H} \chi_W((h^{-1} \cdot g \cdot h)^{-1}) \cdot \chi_V(g) \\ &= \frac{1}{|G|} \frac{1}{|H|} \sum_{g \in G} \sum_{h \in H} \chi_W(h^{-1} \cdot g^{-1} \cdot h) \cdot \chi_V(g). \end{aligned}$$

Through a variable substitution, and the explicit definition χ_W as the zero class function on $G \setminus H$,

$$\begin{aligned} \frac{1}{|G|} \frac{1}{|H|} \sum_{g \in G} \sum_{h \in H} \chi_W(g^{-1}) \cdot \chi_V(h \cdot g \cdot h^{-1}) &= \frac{1}{|G|} \frac{1}{|H|} \sum_{g \in G} \sum_{h \in H} \chi_W(g^{-1}) \cdot \chi_V(g) \\ &= \frac{1}{|H|} \sum_{g \in G} \chi_W(g^{-1}) \cdot \chi_V(g) \\ &= \frac{1}{|H|} \sum_{g \in H} \chi_W(g^{-1}) \cdot \chi_V(g) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{|H|} \sum_{g \in H} \bar{\chi}_W(g) \cdot \text{Res}_H^G(\chi_V)(g) \\
&= \langle \chi_W, \text{Res}_H^G(\chi_V) \rangle_H.
\end{aligned}$$

It is worth noting that the induced representation is not the inverse operation of the restricted representation. However, they are regarded as adjoint in the sense made precise by the Frobenius reciprocity [Theorem A.16](#).

In the general case, the relationship between the irreducible representations of a finite group G and those of its subgroups H is not established. Nevertheless, a distinct circumstance emerges when H is a **normal** subgroup of G .

Proposition. *Consider a finite group G and a normal subgroup H of G . Let (ρ, V) be a representation of the quotient group G/N . Define a representation (σ, V) of G as follows: for all $g \in G$, let \bar{g} denote the representative of g in G/N , and define $\sigma(g) := \rho(\bar{g})$. Then σ is irreducible if and only if ρ is irreducible.*

Proof. Let χ be the character of σ , then by definition of σ ,

$$\begin{aligned}
\langle \chi, \chi \rangle_G &= \frac{1}{|G|} \sum_{g \in G} \bar{\chi}(g) \cdot \chi(g) \\
&= \frac{1}{|G|} \sum_{\bar{g} \in G/N} \sum_{h \in H} \bar{\chi}(\bar{g} \cdot h) \cdot \chi(\bar{g} \cdot h) \\
&= \frac{|H|}{|G|} \sum_{\bar{g} \in G/N} \bar{\chi}(\bar{g}) \cdot \chi(\bar{g}) \\
&= \frac{1}{|G/N|} \sum_{\bar{g} \in G/N} \bar{\chi}(\bar{g}) \cdot \chi(\bar{g}),
\end{aligned}$$

where the last equality is the Hermitian inner product of the character of ρ .

A.2 Matrix groups

A.2.1 Representations of connected compact matrix groups

The preceding Sections were specifically devoted to the study of the representation theory of finite groups. Specifically, the concepts of complete reducibility

of group representations and the interplay between irreducible representations and characters were thoroughly examined. Moving forward, this section aims to extend the understanding of representation theory by examining the representations of infinite matrix groups, which are closed subgroups of the group consisting of invertible matrices.

A representation of a matrix group G is a pair (ρ, V) , where V is a complex vector space with dimension d , and $\rho : G \rightarrow \text{GL}(V)$ is a group continuous homomorphism. The continuity of ρ is equivalent to each of the component maps $g \mapsto (\rho(g))_{i,j}$ being continuous, with $i, j \in \{1, \dots, d\}$. If the component maps are rational functions of the matrix components, then the representation (ρ, V) is referred to as being **ratio-nal**, while if the component maps are polynomial functions of the matrix components, then the representation (ρ, V) is referred to as being **polynomial**.

Theorem. *The rational representations of a matrix group have component maps polynomial in the matrix components and the inverse of the determinant.*

When considering finite groups, the normalized summation $\frac{1}{|G|} \sum_{g \in G}$ is frequently used as a means of averaging over the group. This technique is especially pertinent in the context of reducibility, in [Section A.1.2](#) and character theory, in [Section A.1.3](#).

Generalising this practice to infinite matrix groups is not possible. However, for certain matrix groups, namely the compact matrix group, it is possible to perform an integration over a normalized **Haar measure** as $\int_G dg$.

Consider a compact matrix group G , and let $L^2(G)$ denote the Hilbert space of all complex function on G , with finite 2-norm, for the Hermitian inner product defined on any $\phi, \psi \in L^2(G)$ by

$$\langle \phi, \psi \rangle := \int_G \phi(g) \bar{\psi}(g) dg.$$

In the case where G is a finite group equipped with the **counting measure** as its Haar measure, an isomorphism between the Hilbert space $L^2(G)$ and the group algebra $\mathbb{C}[G]$ is established via the map

$$f \mapsto \sum_{g \in G} f(g) \cdot g.$$

The collection of results presented herein, alongside their corollaries, are commonly referred to as the **Peter-Weyl Theorem**.

Theorem A.17 (Peter-Weyl). *Let G be a compact matrix group, then the following hold.*

1. *For any non-identity matrix g in G , there exists an irreducible represen-*

- tation of G which maps g to a non-identity matrix.
2. The matrix components of inequivalent irreducible representations of G , scaled by a factor of the square root of the dimension of the corresponding irreducible representation, constitute an orthonormal basis of the Hilbert space $L^2(G)$.
 3. The Hilbert space $L^2(G)$ is isomorphic to a Hilbert space direct sum of irreducible representations of G , where the multiplicities correspond to the dimensions of the corresponding irreducible representations.

Corollary. *The compact matrix groups exhibit the properties of complete reducibility and character orthogonality.*

The third assertion of the Peter–Weyl’s [Theorem A.17](#) constitutes the compact matrix group analog of [Proposition A.6](#).

Example. Let GL_2 be the **complex general linear group**, consisting of invertible matrices of degree 2, and which is locally compact but not compact, resulting in a non-finite Haar measure. Let the representation (ρ, \mathbb{C}^2) of GL_2 defined on $g \in GL_2$ as follows:

$$\rho(g) := \begin{pmatrix} 1 & \log |\det g| \\ 0 & 1 \end{pmatrix}.$$

As this representation admits only one-dimensional invariant subspace, namely $\text{Span}_{\mathbb{C}}(1, 0)$, the representation is not completely reducible. Indeed, if for all $g \in GL_2$, there exists $\lambda_g \in \mathbb{C}$ and $(c_1, c_2) \in \mathbb{C}^2$ such that $g \cdot (c_1, c_2) = \lambda_g \cdot (c_1, c_2)$, then it follows from

$$\begin{aligned} c_1 + \log(|\det g|) \cdot c_2 &= \lambda_g \cdot c_1 \\ c_2 &= \lambda_g \cdot c_2, \end{aligned}$$

that $c_2 = 0$.

A **torus** T of a compact matrix group G is defined as a compact and connected abelian subgroup of G . A torus of G is said to be **maximal** if there is no other torus of G that contains it as a proper subgroup.

Let \mathbb{S}^1 denotes the **unit 1-sphere** defined by $\mathbb{S}^1 := \{e^{i\theta} \mid \theta \in [0, 2\pi)\}$ on the complex plane, and isomorphic to the unit circle on the real plane.

Theorem A.18. *For all tori T , there exists some $k \in \mathbb{N}$ such that T is isomorphic to a direct product of k copies of the unit 1-sphere \mathbb{S}^1 .*

In what follows, a given element t in a torus T is implicitly expressed by means of the isomorphism in [Theorem A.18](#), whereby t is written as (z_1, \dots, z_k) with $z_i \in \mathbb{S}^1$, given $k \in \mathbb{N}$, the number of direct products of copies of \mathbb{C}^* via the isomorphism.

Example. Let SO_{2n} be the **special orthogonal group**, which consists of the special orthogonal matrices of even degree $2n$. An instance of SO_{2n} 's maximal tori is the subgroup T consisting of the matrices of the following structure

$$\begin{pmatrix} \cos \theta_1 & \sin \theta_1 & \cdots & 0 & 0 \\ -\sin \theta_1 & \cos \theta_1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \cos \theta_n & \sin \theta_n \\ 0 & 0 & \cdots & -\sin \theta_n & \cos \theta_n \end{pmatrix},$$

where θ_i belongs to the interval $[0, 2\pi)$. In other words, T is the set of all block-diagonal matrices with 2×2 rotation matrix blocks. Each block is isomorphic to an element of \mathbb{S}^1 .

Let the automorphism of a compact matrix group G , given by the map $h \mapsto g^{-1} \cdot h \cdot g$ for all $h \in G$. This automorphism induces a transformation on the set of maximal tori of G through conjugation. More specifically, it maps a torus T to $g^{-1} \cdot T \cdot g$, which is itself a maximal torus of G .

Theorem A.19 (Eli Cartan). *Let G be connected compact matrix group, then the following hold.*

1. *Each element of G is an element of some maximal torus of G .*
2. *All maximal tori of G are conjugate to each other in G .*

Corollary. *Let T be a maximal torus of a connected compact matrix group G , then*

$$G = \bigcup_{g \in G} g^{-1} \cdot T \cdot g.$$

Consider a connected compact matrix group G and let T be a maximal torus of G . According to Eli Cartan's [Theorem A.19](#), for all $g \in G$ and $t \in T$ there exists $h \in G$ such that $g = u^{-1} \cdot t \cdot u$. Let (ρ, V) be a representation of G , then

$$\rho(g) = \rho(u^{-1}) \cdot \rho(t) \cdot \rho(u),$$

which implies that $\rho(g)$ and $\rho(t)$ have the same spectrum.

Let $N_G(T)$ denote the **normalizer** of T . The **Weyl group** of G with respect to T is defined as the quotient of $N_G(T)$ by T , denoted by $W(T) := N_G(T)/T$. The Weyl group acts on T through conjugation, where for any $w \in W(T)$ and $t \in T$, the action is given by $w \cdot t := w^{-1} \cdot t \cdot w$.

Proposition. *Consider a connected and compact matrix group G and let T be a maximal torus of G . The Weyl group $W(T)$ is a finite group. Furthermore, two elements $g, h \in T$ are conjugate in G if and only if there exists an element $w \in W(T)$ satisfying $w \cdot g = h$.*

Corollary. *Let G be a connected compact matrix group and let T be a maximal torus of G . The space of continuous class functions on G is isomorphic to the space of continuous complex functions on T which are invariant under the action $W(T)$. Specifically every continuous complex functions on T which are invariant under the action $W(T)$, extends uniquely to a continuous class functions on G .*

Corollary (Weyl integration formula). *Let T be a maximal torus of a connected compact matrix group G , and let f be a continuous class function on G . Then there exists a continuous real function h on T such that*

$$\int_G f(g) \, dg = \int_T f(t) h(t) \, dt.$$

Let G be a connected compact matrix group, with a maximal torus T . The Weyl group $W(T)$ can be identified to the group of automorphisms of T that are induced by inner automorphisms of G . Given any element $g \in G$, it holds that $N_G(g^{-1} \cdot T \cdot g) = g^{-1} \cdot N_G(T) \cdot g$. Consequently, the Weyl groups of G with respect to distinct maximal tori of G are all isomorphic.

Example. Let U_n be the unitary group of degree n . The group of diagonal unitary matrices forms a maximal torus of U_n , denoted by T . Any unitary matrix can be decomposed into a diagonal form via conjugation by another unitary matrix, so U_n consists of diagonalizable matrices. The quotient of U_n by the conjugacy relation yields the conjugacy classes of diagonal matrices with diagonal entries being roots of unity, which are the eigenvalues of the matrices. In other words, each conjugacy class is represented by a diagonal matrix of the form

$$\text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n}),$$

where θ_i is a real number in the interval $[0, 2\pi)$. The spectrum of a matrix is preserved under conjugation, but representatives of these conjugacy classes are unique only up to a permutation of the diagonal components. The permutation group acting here is precisely the Weyl group $W(T)$. In the case of U_n , this is

the entire symmetric group S_n . Hence, then quotient of the unitary group U_n by the conjugacy relation is equal to the quotient of the maximal torus T by the Weyl group $W(T)$.

Let G denote a compact and connected matrix group, and let T be a maximal torus of G . Then, T is an abelian group. The irreducible representations of T are one-dimensional. More specifically, they are continuous homomorphisms $\rho : T \rightarrow \mathbb{C}^*$, that map each element $z \in (\mathbb{S}^1)^k$ of T , to a product of the form

$$\rho(z_1, \dots, z_k) = \prod_{i=1}^k z_i^{\lambda_i},$$

for some λ in \mathbb{Z}^k , referred to as a **weight**. It follows that for each weight λ in \mathbb{Z}^k , there exists an irreducible representation of T of this form.

Consider a representation (ρ, V) of a group G , and let $(\text{Res}_T^G(\rho), V)$ denote the restricted representation of G on V with respect to the subgroup T . By the Peter–Weyl’s [Theorem A.17](#), the restricted representation $(\text{Res}_T^G(\rho), V)$ can be decomposed into a direct sum of one-dimensional irreducible representations of T . This decomposition takes the form of

$$V \simeq \bigoplus_{\lambda \in \mathbb{Z}^k} V_\lambda^{\oplus n_\lambda},$$

where each V_λ is referred to as the **weight space** corresponding to the weight λ .

Example. Let SU_2 be the **special unitary group**, which consists of the special unitary matrices of degree 2. Consider a maximal torus of SU_2 , denoted by T , which is a subgroup of SU_2 consisting of diagonal matrices of the form:

$$\begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix},$$

where $\theta \in [0, 2\pi)$. Thus, T is isomorphic to \mathbb{S}^1 . Let (ρ, V) be a representation of SU_2 of dimension d , and consider the restricted representation $(\text{Res}_T^{SU_2}(\rho), V)$ and its decomposition into a direct sum of weight spaces:

$$V \simeq \bigoplus_{i=1}^d V_i.$$

In this basis, the action of the restricted representation $\text{Res}_T^{SU_2}(\rho)$ to an element

of the torus T is given by

$$\rho \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} = \text{diag}(e^{i\lambda_1\theta}, \dots, e^{i\lambda_d\theta}),$$

for some weights $\lambda_i \in \mathbb{Z}$.

Suppose G is a connected compact group of $n \times n$ matrices having a maximal torus T , and consider the decomposition of \mathbb{C}^n into a direct sum of weight spaces, for the representation of T that maps each element $t \in T$ to itself:

$$\mathbb{C}^n \simeq \bigoplus_{\lambda \in \mathbb{Z}^k} V_{\lambda}^{\oplus n_{\lambda}}.$$

The Weyl group $W(T)$ acts on this decomposition by permuting the weights λ .

Example. Let SU_3 be the special unitary group of degree 3. An instance of SU_3 's maximal tori is the subgroup T consisting of the diagonal matrices,

$$\begin{pmatrix} e^{i\theta_1} & 0 & 0 \\ 0 & e^{i\theta_2} & 0 \\ 0 & 0 & e^{-i(\theta_1+\theta_2)} \end{pmatrix},$$

where θ_1 and θ_2 belong to the interval $[0, 2\pi)$. Therefore, T is isomorphic to \mathbb{S}^2 , via the map

$$\begin{pmatrix} e^{i\theta_1} & 0 & 0 \\ 0 & e^{i\theta_2} & 0 \\ 0 & 0 & e^{-i(\theta_1+\theta_2)} \end{pmatrix} \mapsto (e^{i\theta_1}, e^{i\theta_2}).$$

Using this isomorphism, the action of each $(z_1, z_2) \in \mathbb{S}^2$, to the elements e_1, e_2 and e_3 forming the standard basis of \mathbb{C}^3 , is given by

$$\begin{aligned} (z_1, z_2) \cdot e_1 &= z_1^1 \cdot z_2^0 \\ (z_1, z_2) \cdot e_2 &= z_1^0 \cdot z_2^1 \\ (z_1, z_2) \cdot e_3 &= z_1^{-1} \cdot z_2^{-1}, \end{aligned}$$

This action yields the 3 weights $(1, 0)$, $(0, 1)$ and $(-1, -1)$, and their corresponding weight spaces. The Weyl group $W(T)$, being isomorphic to the symmetric group S_3 , acts on these weight spaces by permuting the weights via the above action.

Concluding the section, a summary of the various matrix groups of degree n that

have been used, and their properties is presented, assuming that the degree n is larger than 1:

Matrix group	Topology
Complex general linear group $\mathrm{GL}_n := \{g \in M_n(\mathbb{C}) \mid \det g \neq 0\}$	not compact connected / not simply connected
Complex special linear group $\mathrm{SL}_n := \{g \in M_n(\mathbb{C}) \mid \det g = 1\}$	not compact simply connected
Unitary group $\mathrm{U}_n := \{g \in M_n(\mathbb{C}) \mid gg^* = I\}$	compact connected / not simply connected
Special unitary group $\mathrm{SU}_n := \{g \in M_n(\mathbb{C}) \mid gg^* = I \text{ and } \det g = 1\}$	compact simply connected
Orthogonal group $\mathrm{O}_n := \{g \in M_n(\mathbb{R}) \mid gg^T = I\}$	compact not connected
Special orthogonal group $\mathrm{SO}_n := \{g \in M_n(\mathbb{R}) \mid gg^T = I \text{ and } \det g = 1\}$	compact connected / not simply connected

With the inclusions $\mathrm{O}_n \subseteq \mathrm{U}_n \subseteq \mathrm{GL}_n$ and $\mathrm{SO}_n \subseteq \mathrm{SU}_n \subseteq \mathrm{SL}_n$.

A.2.2 Representations of U_d

The **unitary group** of degree d , denoted U_d , is the group of $d \times d$ unitary matrices, i.e., matrices U with complex entries such that $UU^* = U^*U = I$, where U^* denotes the conjugate transpose of U .

An equivalent definition of the unitary group is as the matrix group that preserves the standard Hermitian inner product $\langle \cdot, \cdot \rangle$ defined on the complex vector space \mathbb{C}^d . That is, for all vectors $x, y \in \mathbb{C}^d$ and for all $U \in \mathrm{U}_d$, it holds that

$$\langle Ux, Uy \rangle = \langle x, y \rangle.$$

Let T be the group of diagonal unitary matrices, a maximal torus of the unitary group U_d . Let (ρ, V) be a representation of U_d , and $(\mathrm{Res}_T^{\mathrm{U}_d}(\rho), V)$ the restricted

representation of U_d on V with respect to the subgroup T , such that

$$V \simeq \bigoplus_{\lambda \in \mathbb{Z}^d} V_\lambda^{\oplus n_\lambda},$$

is the decomposition of V into a direct sum of one-dimensional irreducible representations of T , with some weights $\lambda \in \mathbb{Z}^d$. The maximal element in the set of these weights, with respect to the lexicographic order, is denoted the **highest weight**.

Example. Let U_d be the unitary group of degree d , and let T be the group of diagonal unitary matrices, an instance of U_d 's maximal tori. Let (ρ, \mathbb{C}^d) be the representation of U_d that maps each element $U \in U_d$ to itself, i.e. $\rho(U) = U$ for all $U \in U_d$. Introduce the element $\text{diag}(e^{i\theta_1}, \dots, e^{i\theta_d})$ of T , and let e_1, \dots, e_d constitute the standard basis of the vector space \mathbb{C}^d . The relationship between these elements is given by the equation:

$$\text{diag}(e^{i\theta_1}, \dots, e^{i\theta_d}) \cdot e_i = e^{i\theta_i} \cdot e_i.$$

The decomposition of \mathbb{C}^d into a direct sum of one-dimensional irreducible representations of T can be expressed as

$$V \simeq \bigoplus_{i=1}^d V_i,$$

where the index i denotes the weight $\lambda \in \mathbb{Z}^d$, such that $\lambda_i = 1$ and $\lambda_j = 0$ for every $j \neq i$. Consequently, the highest weight is $(1, 0, \dots, 0)$.

The unitary group U_d has a trivial irreducible representation of dimension 1, which assigns the scalar value 1 to every unitary matrix $U \in U_d$. Additionally, there exists a d -dimensional irreducible representation of U_d , wherein each unitary matrix $U \in U_d$ is mapped to itself. However, no irreducible representation of intermediate dimension, i.e., between 1 and n , can be found. In order to discover new irreducible representations, it is necessary to increase the dimension. One possible approach to achieving this is to consider a complex tensor product space.

Theorem A.20. *For each weight $\lambda \in \mathbb{Z}^d$ such that $\lambda_1 \geq \dots \geq \lambda_d$, there exists a unique irreducible representation V_λ of the unitary group U_d , with highest weight λ . These are all inequivalent and they are all the irreducible representations of U_d .*

Consider the complex tensor product space $(\mathbb{C}^d)^{\otimes n}$. Define a representation of

the unitary group U_d on this space by

$$U \mapsto U^{\otimes n}.$$

Additionally, define a representation of the symmetric group \mathfrak{S}_n on this space by permuting the tensor positions, and extend this action linearly to the group algebra $\mathbb{C}[\mathfrak{S}_n]$.

Example. Consider the complex tensor product space $(\mathbb{C}^d)^{\otimes 3}$. Let $v_1, v_2, v_3 \in \mathbb{C}^d$ be arbitrary vectors, and $\sigma \in \mathfrak{S}_n$ be arbitrary permutation, then

$$\sigma \cdot (v_1 \otimes v_2 \otimes v_3) = v_{\sigma^{-1}(1)} \otimes v_{\sigma^{-1}(2)} \otimes v_{\sigma^{-1}(3)}.$$

Define T as the canonical Young tableau corresponding to the partition (3) of 3. The action of the Young symmetrizer s_T on the complex tensor product space $(\mathbb{C}^d)^{\otimes 3}$ gives rise to a new complex tensor product space, $s_T \cdot (\mathbb{C}^d)^{\otimes 3}$, which is referred to as the **symmetric subspace** of $(\mathbb{C}^d)^{\otimes 3}$, and denoted by $\vee^3 \mathbb{C}^d$. For instance,

$$\begin{aligned} s_T \cdot (v_1 \otimes v_2 \otimes v_3) &= \sum_{\sigma \in \mathfrak{S}_3} \sigma \cdot (v_1 \otimes v_2 \otimes v_3) \\ &= (v_1 \otimes v_2 \otimes v_3) + (v_2 \otimes v_1 \otimes v_3) + (v_3 \otimes v_2 \otimes v_1) + \\ &\quad (v_1 \otimes v_3 \otimes v_2) + (v_3 \otimes v_1 \otimes v_2) + (v_2 \otimes v_3 \otimes v_1). \end{aligned}$$

The symmetric group \mathfrak{S}_3 acts trivially on $\vee^3 \mathbb{C}^d$, i.e. for all $\sigma \in \mathfrak{S}_3$ and $x \in \vee^3 \mathbb{C}^d$, then $\sigma \cdot x = x$. Let T be the canonical Young tableau associated with the partition (1, 1, 1) of 3. The complex tensor product space $s_T \cdot (\mathbb{C}^d)^{\otimes 3}$, is referred to as the **antisymmetric subspace** of $(\mathbb{C}^d)^{\otimes 3}$, and denoted by $\wedge^3 \mathbb{C}^d$. For instance,

$$\begin{aligned} s_T \cdot (v_1 \otimes v_2 \otimes v_3) &= \sum_{\sigma \in \mathfrak{S}_3} \sigma \cdot (v_1 \otimes v_2 \otimes v_3) \\ &= (v_1 \otimes v_2 \otimes v_3) - (v_2 \otimes v_1 \otimes v_3) - (v_3 \otimes v_2 \otimes v_1) - \\ &\quad (v_1 \otimes v_3 \otimes v_2) + (v_3 \otimes v_1 \otimes v_2) + (v_2 \otimes v_3 \otimes v_1). \end{aligned}$$

The symmetric group \mathfrak{S}_3 acts on $\wedge^3 \mathbb{C}^d$ through multiplication by the signature, i.e. for all $\sigma \in \mathfrak{S}_3$ and $x \in \wedge^3 \mathbb{C}^d$, then $\sigma \cdot x = \text{sign}(\sigma) \cdot x$.

Let U_d be the unitary group of degree d . Given a partition $\lambda \vdash n$ consisting of l parts, it is possible to associate λ with a weight in \mathbb{Z}^d , provided that $l \leq d$, by

appending $d - l$ zeros to the right of the partition.

Theorem. Consider $\lambda \vdash n$ a partition of n with l parts. Let T a Young tableau without repetitions on λ . Then, the complex tensor product space $s_T \cdot (\mathbb{C}^d)^{\otimes n}$ is the zero space, if $l > d$, and an irreducible representation of U_d , for the representation $U \mapsto U^{\otimes n}$, with highest weight λ , if $l \leq d$.

Corollary. All irreducible representations of the unitary group U_n , highest weights $\lambda \in \mathbb{N}^d$, is equivalent to $s_T \cdot (\mathbb{C}^d)^{\otimes n}$, with T a Young tableau without repetitions on λ .

From [Theorem A.20](#), for a given partition $\lambda \vdash n$ with at most d part, the irreducible representations $s_T \cdot (\mathbb{C}^d)^{\otimes n}$ of the of the unitary group U_d , are mutually equivalent across all Young tableaux without repetition T on λ , thereby solely dependent on λ , and consequently, it is possible to designate any representation of this type as V_λ^d .

Theorem. Consider a partition $\lambda \vdash n$ with at most d parts. Let V_λ^d be the corresponding irreducible representation of the unitary group U_d . The dimension of V_λ^d is equivalent to the cardinality of the set of semistandard Young tableaux associated with λ , with entries in $\{1, \dots, d\}$

Since the entries in each columns of a semistandard Young tableau are strictly increasing, there is no semistandard Young tableau with entries in $\{1, \dots, d\}$, and with more than d rows.

Example. Let U_2 be the unitary group of degree 2. Consider λ the partition $(1, 1, 1)$ of 3, then the antisymmetric subspace $V_\lambda^3 \simeq \wedge^3 \mathbb{C}^2$ is the zero space. Let λ be the partition $(2, 1)$ of 3, then the irreducible representation V_λ^2 of U_2 has dimension 2, since the semistandard Young tableaux associated with λ , with entries in $\{1, 2\}$ are

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \quad \text{and} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}$$

For all $k \in \mathbb{Z}$ there is a 1-dimensional irreducible representation of U_d defined by $U \mapsto (\det U)^k$, and denoted \det^k . This irreducible representation is rational for all $k \in \mathbb{Z}$ and polynomial for $k \in \mathbb{N}$.

Theorem. All irreducible representations of U_d with highest weight $\lambda \in \mathbb{Z}^d$ are equivalent to the representation

$$\det^{-k} \cdot V_\mu^d,$$

where $\mu \vdash n$ is a partition of n with at most d parts, and $k \in \mathbb{N}$ is an integer such that $\lambda_i = \mu_i - k$.

The set of unitary matrices U_d is dense, with respect to the Zariski topology, within the set of complex invertible matrices GL_d .

Theorem. *The irreducible rational representations of GL_d are the same as the irreducible representations of U_d .*

Bibliography

- [Ada82] John Frank Adams. *Lectures on Lie groups*. University of Chicago Press, 1982 (page 107).
- [All+23] Rene Allerstorfer, Matthias Christandl, Dmitry Grinko, Ion Nechita, Maris Ozols, Denis Rochette, and Philip Verduyn Lunel. *Monogamy of highly symmetric states*. 2023. arXiv: 2309.16655 [quant-ph] (page 87).
- [Ami+08] Luigi Amico, Rosario Fazio, Andreas Osterloh, and Vlatko Vedral. « Entanglement in many-body systems ». In: *Reviews of modern physics* 80.2 (2008), p. 517 (page 87).
- [AS17] Guillaume Aubrun and Stanisław J Szarek. *Alice and Bob meet Banach*. Vol. 223. American Mathematical Soc., 2017 (page 31).
- [AST17] Henning Haahr Andersen, Catharina Stroppel, and Daniel Tubbenhauer. « Semisimplicity of Hecke and (walled) Brauer algebras ». In: *Journal of the Australian Mathematical Society* 103.1 (2017), pp. 1–44 (page 18).
- [Aub18] Guillaume Aubrun. *A naive look at Schur–Weyl duality*. 2018. URL: <http://math.univ-lyon1.fr/~aubrun/recherche/schur-weyl.pdf> (page 24).
- [Bak03] Andrew Baker. *Matrix groups: An introduction to Lie group theory*. Springer Science & Business Media, 2003 (page 107).
- [BB84] Charles H. Bennett and Gilles Brassard. « Quantum cryptography: Public key distribution and coin tossing ». In: *Theoretical Computer Science* 560 (2014), pp. 7–11 (pages 46, 51).
- [BBV04] Stephen Boyd, Stephen P Boyd, and Lieven Vandenberghe. *Convex optimization*. Cambridge university press, 2004 (page 92).
- [BC17] Jacob C Bridgeman and Christopher T Chubb. « Hand-waving and interpretive dance: an introductory course on tensor networks ». In: *Journal of Physics A: Mathematical and Theoretical* 50.22 (2017), p. 223001 (page 40).

- [Ben+94] Georgia Benkart, Manish Chakrabarti, Thomas Halverson, Robert Leduc, Chanyoung Y Lee, and Jeffrey Stroome. « Tensor product representations of general linear groups and their connections with Brauer algebras ». In: *Journal of Algebra* 166.3 (1994), pp. 529–567 (page 28).
- [BH96] Vladimir Bužek and Mark Hillery. « Quantum copying: Beyond the no-cloning theorem ». In: *Physical Review A* 54.3 (1996), p. 1844 (page 45).
- [BL20] Anne Broadbent and Sébastien Lord. « Uncloneable Quantum Encryption via Oracles ». In: *15th Conference on the Theory of Quantum Computation, Communication and Cryptography (TQC 2020)*. Schloss Dagstuhl-Leibniz-Zentrum für Informatik. 2020 (page 46).
- [Boe+70] Hermann Boerner et al. *Representations of groups: with special consideration for the needs of modern physics*. Vol. 5. North-Holland, 1970 (page 107).
- [Bra37] Richard Brauer. « On algebras which are connected with the semisimple continuous groups ». In: *Annals of Mathematics* (1937), pp. 857–872 (page 27).
- [Bru+00] Dagmar Bruß, Mirko Cinchetti, G Mauro D’Ariano, and Chiara Macchiavello. « Phase-covariant quantum cloning ». In: *Physical Review A* 62.1 (2000), p. 012302 (page 45).
- [Bru+98] Dagmar Bruß, David P DiVincenzo, Artur Ekert, Christopher A Fuchs, Chiara Macchiavello, and John A Smolin. « Optimal universal and state-dependent quantum cloning ». In: *Physical Review A* 57.4 (1998), p. 2368 (page 45).
- [BT13] Theodor Bröcker and Tammo Tom Dieck. *Representations of compact Lie groups*. Vol. 98. Springer Science & Business Media, 2013 (page 107).
- [Bul20] Daria V Bulgakova. « Some aspects of representation theory of walled Brauer algebras ». PhD thesis. Aix Marseille Université, 2020 (page 20).
- [Bum+04] Daniel Bump et al. *Lie groups*. Vol. 8. Springer, 2004 (page 107).
- [BV07] Pierfrancesco Buonsante and Alessandro Vezzani. « Ground-state fidelity and bipartite entanglement in the Bose-Hubbard model ». In: *Physical review letters* 98.11 (2007), p. 110601 (page 87).
- [Cer+04] NJ Cerf, O Krueger, P Navez, RF Werner, and MM Wolf. « The optimal cloning of quantum coherent states is non-Gaussian ». In: *arXiv preprint quant-ph/0410058* (2004) (page 45).

- [Cer+05] NJ Cerf, O Krüger, Patrick Navez, RF Werner, and Michael M Wolf. « Non-Gaussian cloning of quantum coherent states is optimal ». In: *Physical review letters* 95.7 (2005), p. 070501 (page 45).
- [Cer00] Nicolas J Cerf. « Asymmetric quantum cloning in any dimension ». In: *Journal of modern optics* 47.2-3 (2000), pp. 187–209 (page 45).
- [Cer98] Nicolas J Cerf. « Asymmetric quantum cloning machines ». In: *J. Mod. Opt.* 47.KRL-MAP-224 (1998), p. 187 (page 45).
- [Chr+21] Matthias Christandl, Felix Leditzky, Christian Majenz, Graeme Smith, Florian Speelman, and Michael Walter. « Asymptotic performance of port-based teleportation ». In: *Communications in Mathematical Physics* 381 (2021), pp. 379–451 (page 56).
- [ČHS12] Piotr Źwikliński, Michał Horodecki, and Michał Studziński. « Region of fidelities for a $1 \rightarrow N$ universal qubit quantum cloner ». In: *Physics Letters A* 376.32 (2012), pp. 2178–2187 (page 45).
- [CK17] Bob Coecke and Aleks Kissinger. *Picturing quantum processes*. Cambridge University Press, 2017 (page 40).
- [Cox+08] Anton Cox, Maud De Visscher, Stephen Doty, and Paul Martin. « On the blocks of the walled Brauer algebra ». In: *Journal of Algebra* 320.1 (2008), pp. 169–212 (page 20).
- [CST10] Tullio Ceccherini-Silberstein, Fabio Scarabotti, and Filippo Tolli. *Representation theory of the symmetric groups: the Okounkov-Vershik approach, character formulas, and partition algebras*. Vol. 121. Cambridge University Press, 2010 (page 107).
- [DD04a] Thomas Durt and Jianfeng Du. « About economic qubit cloning' ». In: *Phys. Rev. A* 69 (2004), p. 62316 (page 45).
- [DD04b] Thomas Durt and Jiangfeng Du. « Characterization of low-cost one-to-two qubit cloning ». In: *Physical Review A* 69.6 (2004), p. 062316 (page 45).
- [DFC05] Thomas Durt, Jaromír Fiurášek, and Nicolas J Cerf. « Economical quantum cloning in any dimension ». In: *Physical Review A* 72.5 (2005), p. 052322 (page 45).
- [DG98] Lu-Ming Duan and Guang-Can Guo. « Probabilistic cloning and identification of linearly independent quantum states ». In: *Physical review letters* 80.22 (1998), p. 4999 (page 45).

- [Die82] Dennis Dieks. « Communication by EPR devices ». In: *Physics Letters A* 92.6 (1982), pp. 271–272 (page 46).
- [DLS19] Stephen Doty, Aaron Lauve, and George H Seelinger. « Canonical idempotents of multiplicity-free families of algebras ». In: *L'Enseignement Mathématique* 64.1 (2019), pp. 23–63 (pages 95, 96).
- [DM03] Giacomo Mauro D'Ariano and Chiara Macchiavello. « Optimal phase-covariant cloning for qubits and qutrits ». In: *Physical Review A* 67.4 (2003), p. 042306 (page 45).
- [DMP05] Giacomo Mauro D'Ariano, Chiara Macchiavello, and Paolo Perinotti. « Superbroadcasting of mixed states ». In: *Physical review letters* 95.6 (2005), p. 060503 (page 47).
- [DPS04] Andrew C Doherty, Pablo A Parrilo, and Federico M Spedalieri. « Complete family of separability criteria ». In: *Physical Review A* 69.2 (2004), p. 022308 (page 37).
- [DWH99] William F Doran IV, David B Wales, and Philip J Hanlon. « On the semisimplicity of the Brauer centralizer algebras ». In: *Journal of Algebra* 211.2 (1999), pp. 647–685 (page 17).
- [Eve57] Hugh Everett III. « " Relative state" formulation of quantum mechanics ». In: *Reviews of modern physics* 29.3 (1957), p. 454 (page 34).
- [FFC05] Jaromír Fiurášek, Radim Filip, and Nicolas J Cerf. « Highly asymmetric quantum cloning in arbitrary dimension ». In: *Quantum Information & Computation* 5.7 (2005), pp. 583–592 (page 45).
- [FH13] William Fulton and Joe Harris. *Representation theory: a first course*. Vol. 129. Springer Science & Business Media, 2013 (pages 12, 23, 107).
- [Fiu01] Jaromír Fiurášek. « Extremal equation for optimal completely positive maps ». In: *Physical Review A* 64.6 (2001), p. 062310 (page 56).
- [FST22] Omar Fawzi, Ala Shayeghi, and Hoang Ta. « A hierarchy of efficient bounds on quantum capacities exploiting symmetry ». In: *IEEE Transactions on Information Theory* 68.11 (2022), pp. 7346–7360 (page 49).
- [GM97] Nicolas Gisin and Serge Massar. « Optimal quantum cloning machines ». In: *Physical review letters* 79.11 (1997), p. 2153 (page 45).
- [GO22] Dmitry Grinko and Maris Ozols. « Linear programming with unitary-equivariant constraints ». In: *arXiv preprint arXiv:2207.05713* (2022) (page 49).

- [Gur03] Leonid Gurvits. « Classical deterministic complexity of Edmonds' problem and quantum entanglement ». In: *Proceedings of the thirty-fifth annual ACM symposium on Theory of computing*. 2003, pp. 10–19 (page 37).
- [GW09] Roe Goodman and Nolan R Wallach. *Symmetry, representations, and invariants*. Vol. 255. Springer, 2009 (pages 23, 107).
- [Haa+21] Erkka Haapasalo, Tristan Kraft, Nikolai Miklin, and Roope Uola. « Quantum marginal problem and incompatibility ». In: *Quantum* 5 (2021), p. 476 (page 46).
- [Har13] Aram W Harrow. « The church of the symmetric subspace ». In: *arXiv preprint arXiv:1308.6595* (2013) (page 54).
- [Has17] A-L Hashagen. « Universal Asymmetric Quantum Cloning Revisited ». In: *Quantum Information & Computation* 17.9-10 (2017), pp. 0747–0778 (page 45).
- [HH13] Brian C Hall and Brian C Hall. *Lie groups, Lie algebras, and representations*. Springer, 2013 (page 107).
- [HH99] Michał Horodecki and Paweł Horodecki. « Reduction criterion of separability and limits for a class of distillation protocols ». In: *Physical Review A* 59.6 (1999), p. 4206 (page 36).
- [HHH01] Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki. « Separability of n-particle mixed states: necessary and sufficient conditions in terms of linear maps ». In: *Physics Letters A* 283.1-2 (2001), pp. 1–7 (page 35).
- [HJ20] Tom Halverson and Theodore N Jacobson. « Set-partition tableaux and representations of diagram algebras ». In: *Algebraic Combinatorics* 3.2 (2020), pp. 509–538 (page 9).
- [HLA22] Chung-Yun Hsieh, Matteo Lostaglio, and Antonio Acín. « Quantum channel marginal problem ». In: *Physical Review Research* 4.1 (2022), p. 013249 (page 46).
- [HR05] Tom Halverson and Arun Ram. « Partition algebras ». In: *European Journal of Combinatorics* 26.6 (2005), pp. 869–921 (pages 15, 27).
- [IAG06] Sofyan Iblisdir, Antonio Acín, and Nicolas Gisin. « Generalised asymmetric quantum cloning machines ». In: *Quantum Information & Computation* 6.4 (2006), pp. 410–435 (page 45).

- [Ibl+05] Sofyan Iblisdir, Antonio Acín, Nicolas J Cerf, Radim Filip, Jaromir Fiurášek, and Nicolas Gisin. « Multipartite asymmetric quantum cloning ». In: *Physical Review A* 72.4 (2005), p. 042328 (page 45).
- [Jam06] Gordon Douglas James. *The representation theory of the symmetric groups*. Vol. 682. Springer, 2006 (page 107).
- [Kay14] Alastair Kay. « Optimal universal quantum cloning: Asymmetries and fidelity measures ». In: *arXiv preprint arXiv:1407.4951* (2014) (page 45).
- [Kay16] Alastair Kay. « Optimal Universal Quantum Cloning: Asymmetries and Fidelity Measures ». In: *Quantum Information and Computation* 16.11 & 12 (2016), pp. 0991–1028 (page 45).
- [Koe08] Steffen Koenig. « A panorama of diagram algebras ». In: *Trends in representation theory of algebras and related topics, EMS Ser. Congr. Rep* (2008), pp. 491–540 (page 9).
- [K RK12] Alastair Kay, Ravishankar Ramanathan, and Dagomir Kaszlikowski. « Optimal asymmetric quantum cloning ». In: *arXiv preprint arXiv:1208.5574* (2012) (page 45).
- [KW99] Michael Keyl and Reinhard F Werner. « Optimal cloning of pure states, testing single clones ». In: *Journal of Mathematical Physics* 40.7 (1999), pp. 3283–3299 (pages 45, 70).
- [Lan12] Joseph M Landsberg. « Tensors: geometry and applications ». In: *Representation theory* 381.402 (2012), p. 3 (page 107).
- [Led22] Felix Leditzky. « Optimality of the pretty good measurement for port-based teleportation ». In: *Letters in Mathematical Physics* 112.5 (2022), p. 98 (page 56).
- [LN21] Faedi Loulidi and Ion Nechita. « The compatibility dimension of quantum measurements ». In: *Journal of Mathematical Physics* 62.4 (2021), p. 042205 (page 46).
- [Mar00] Paul P Martin. « The partition algebra and the Potts model transfer matrix spectrum in high dimensions ». In: *Journal of Physics A: Mathematical and General* 33.19 (2000), p. 3669 (page 27).
- [MHS14] Marek Możejko, Michał Horodecki, and Michał Studziński. « Structure and properties of the algebra of partially transposed permutation operators ». In: *Journal of Mathematical Physics* 55.3 (2014), p. 032202 (page 45).

- [Moz+18] Marek Mozrzykmas, Michał Studziński, Sergii Strelchuk, and Michał Horodecki. « Optimal port-based teleportation ». In: *New Journal of Physics* 20.5 (2018), p. 053006 (page 56).
- [MR98] Paul P Martin and Geneviève Rollet. « The Potts model representation and a Robinson–Schensted correspondence for the partition algebra ». In: *Compositio Mathematica* 112.2 (1998), pp. 237–254 (page 27).
- [MS94] Paul Martin and Hubert Saleur. « Algebras in higher-dimensional statistical mechanics—the exceptional partition (mean field) algebras ». In: *letters in mathematical physics* 30 (1994), pp. 179–185 (page 15).
- [MSH18] Marek Mozrzykmas, Michał Studziński, and Michał Horodecki. « A simplified formalism of the algebra of partially transposed permutation operators with applications ». In: *Journal of Physics A: Mathematical and Theoretical* 51.12 (2018), p. 125202 (page 45).
- [NC02] Michael A Nielsen and Isaac Chuang. *Quantum computation and quantum information*. 2002 (page 31).
- [NPR21] Ion Nechita, Clément Pellegrini, and Denis Rochette. « A geometrical description of the universal $1 \rightarrow 2$ asymmetric quantum cloning region ». In: *Quantum Information Processing* 20.10 (2021), p. 333 (pages 45, 57, 58).
- [NPR22] Ion Nechita, Clément Pellegrini, and Denis Rochette. « The Asymmetric Quantum Cloning Region ». In: *arXiv preprint arXiv:2209.11999* (2022) (pages 45, 65, 69).
- [NS21] Ion Nechita and Satvik Singh. « A graphical calculus for integration over random diagonal unitary matrices ». In: *Linear Algebra and its Applications* 613 (2021), pp. 46–86 (page 57).
- [OEIS] *The On-Line Encyclopedia of Integer Sequences*. OEIS. URL: <https://oeis.org> (pages 14, 16, 17).
- [Oka16] Soichi Okada. « Pieri Rules for Classical Groups and Equinumeration between Generalized Oscillating Tableaux and Semistandard Tableaux ». In: *The Electronic Journal of Combinatorics* 23.4 (2016), P4–43 (page 95).
- [Ore+21] Rosa Orellana, Franco Saliola, Anne Schilling, and Mike Zabrocki. « Plethysm and the algebra of uniform block permutations ». In: *arXiv preprint arXiv:2112.13909* (2021) (page 16).

- [Pen71] Roger Penrose. « Applications of negative dimensional tensors ». In: *Combinatorial mathematics and its applications* 1 (1971), pp. 221–244 (page 40).
- [Per95] Asher Peres. « Higher order Schmidt decompositions ». In: *Physics Letters A* 1.202 (1995), pp. 16–17 (page 35).
- [Per96] Asher Peres. « Separability criterion for density matrices ». In: *Physical Review Letters* 77.8 (1996), p. 1413 (page 35).
- [RS06] Hebing Rui and Mei Si. « A criterion on the semisimple Brauer algebras II ». In: *Journal of Combinatorial Theory, Series A* 113.6 (2006), pp. 1199–1203 (page 18).
- [Rui05] Hebing Rui. « A criterion on the semisimple Brauer algebras ». In: *Journal of Combinatorial Theory, Series A* 111.1 (2005), pp. 78–88 (page 18).
- [Sag13] Bruce E Sagan. *The symmetric group: representations, combinatorial algorithms, and symmetric functions*. Vol. 203. Springer Science & Business Media, 2013 (page 107).
- [Sch07] Erhard Schmidt. « Zur Theorie der linearen und nichtlinearen Integralgleichungen ». In: *Mathematische Annalen* 63.4 (1907), pp. 433–476 (page 34).
- [Sch27] Issai Schur. *Über die rationalen Darstellungen der allgemeinen linearen Gruppe*. 1927 (pages 9, 24).
- [Ser+77] Jean-Pierre Serre et al. *Linear representations of finite groups*. Vol. 42. Springer, 1977 (pages 12, 23, 107).
- [SHM13] Michał Studziński, Michał Horodecki, and Marek Mozrzyński. « Com-mutant structure of $U^{\otimes(n-1)} \otimes U^*$ transformations ». In: *Journal of Physics A: Mathematical and Theoretical* 46.39 (2013), p. 395303 (page 45).
- [Sim96] Barry Simon. *Representations of finite and compact groups*. 10. American Mathematical Soc., 1996 (page 107).
- [SPS01] J-M Sixdeniers, KA Penson, and AI Solomon. « Extended Bell and Stir-ling Numbers From Hypergeometric Exponentiation ». In: *Journal of Integer Sequences* 4.2 (2001), p. 3 (page 16).
- [Ste+16] Benjamin Steinberg et al. *Representation theory of finite monoids*. Springer, 2016 (page 16).
- [Ste95] Shlomo Sternberg. *Group theory and physics*. Cambridge University Press, 1995 (page 107).

- [Stu+14] Michał Studziński, Piotr Ćwikliński, Michał Horodecki, and Marek Mozrzy-
mas. « Group-representation approach to $1 \rightarrow N$ universal quantum
cloning machines ». In: *Physical Review A* 89.5 (2014), p. 052322 (page 45).
- [Stu+17] Michał Studziński, Sergii Strelchuk, Marek Mozrzymas, and Michał Horodecki.
« Port-based teleportation in arbitrary dimension ». In: *Scientific reports*
7.1 (2017), p. 10871 (page 56).
- [Tab+22] Gelo Noel M Tabia, Kai-Siang Chen, Chung-Yun Hsieh, Yu-Chun Yin,
and Yeong-Cherng Liang. « Entanglement transitivity problems ». In:
npj Quantum Information 8.1 (2022), p. 98 (page 87).
- [Tan97] Kenichiro Tanabe. « On the centralizer algebra of the unitary reflec-
tion group $G(m, p, n)$ ». In: *Nagoya mathematical journal* 148 (1997),
pp. 113–126 (page 27).
- [Tun85] Wu-Ki Tung. *Group theory in physics*. Vol. 1. World Scientific, 1985
(page 107).
- [VO05] Anatoli M Vershik and A Yu Okounkov. « A new approach to the rep-
resentation theory of the symmetric groups. II ». In: *Journal of Mathe-
matical Sciences* 131 (2005), pp. 5471–5494 (page 107).
- [Wat18] John Watrous. *The theory of quantum information*. Cambridge univer-
sity press, 2018 (pages 31, 38, 39).
- [WBC15] Christopher J Wood, Jacob D Biamonte, and David G Cory. « Tensor
networks and graphical calculus for open quantum systems ». In: *Quan-
tum Information & Computation* 15.9-10 (2015), pp. 759–811 (page 40).
- [Wen88] Hans Wenzl. « On the structure of Brauer’s centralizer algebras ». In:
Annals of Mathematics 128.1 (1988), pp. 173–193 (pages 17, 95).
- [Wer89] Reinhard F Werner. « Quantum states with Einstein-Podolsky-Rosen
correlations admitting a hidden-variable model ». In: *Physical Review A*
40.8 (1989), p. 4277 (page 37).
- [Wer98] Reinhard F Werner. « Optimal cloning of pure states ». In: *Physical
Review A* 58.3 (1998), p. 1827 (pages 45, 47).
- [Wey46] Hermann Weyl. *The classical groups: their invariants and representa-
tions*. 1946 (pages 9, 24).
- [WVI03] Michael M Wolf, Frank Verstraete, and J IGNACIO CIRAC. « Entan-
glement and frustration in ordered systems ». In: *International Journal
of Quantum Information* 1.04 (2003), pp. 465–477 (page 87).

- [WZ82] William K Wootters and Wojciech H Zurek. « A single quantum cannot be cloned ». In: *Nature* 299 (1982), pp. 802–803 (page [46](#)).
- [Zhe73] Dmitrii Petrovich Zhelobenko. *Compact Lie groups and their representations*. Vol. 40. American Mathematical Soc., 1973 (page [107](#)).